



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 10:03 am BST

PDB ID : 5GKS  
Title : Crystal structure of SLE patient-derived anti-DNA antibody  
Authors : Arimori, T.; Sakakibara, S.; Kikutani, H.; Takagi, J.  
Deposited on : 2016-07-05  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	218	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>5%</div> <div>6%</div> </div> </div>
1	C	218	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>•</div> <div>8%</div> </div> </div>
2	B	216	<div> <div>6%</div> <div> <div></div> <div>94%</div> <div>•</div> <div>•</div> </div> </div>
2	D	216	<div> <div>10%</div> <div> <div></div> <div>93%</div> <div>•</div> <div>•</div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6431 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IgG2, Fab (heavy chain).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	206	Total	C	N	O	S	0	0	0
			1564	998	256	306	4			
1	C	200	Total	C	N	O	S	0	0	0
			1529	978	252	295	4			

- Molecule 2 is a protein called lambda, Fab (light chain).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	211	Total	C	N	O	S	0	0	0
			1574	985	262	321	6			
2	D	209	Total	C	N	O	S	0	0	0
			1560	978	259	317	6			

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	65	Total	O	0	0
			65	65		
4	B	31	Total	O	0	0
			31	31		
4	C	64	Total	O	0	0
			64	64		
4	D	34	Total	O	0	0
			34	34		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: IgG2, Fab (heavy chain)





- Molecule 1: IgG2, Fab (heavy chain)



- Molecule 2: lambda, Fab (light chain)



- Molecule 2: lambda, Fab (light chain)



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	139.36Å 41.17Å 156.00Å 90.00° 113.07° 90.00°	Depositor
Resolution (Å)	29.65 – 2.05 29.65 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.65-2.05) 99.9 (29.65-2.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.70 (at 2.04Å)	Xtriage
Refinement program	REFMAC 5.8.0124	Depositor
R, $R_{free}$	0.217 , 0.255 0.222 , 0.262	Depositor DCC
$R_{free}$ test set	2618 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.6	Xtriage
Anisotropy	0.656	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 37.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6431	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 72.81 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.0598e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	0/1607	0.75	2/2198 (0.1%)
1	C	0.58	0/1571	0.76	2/2147 (0.1%)
2	B	0.50	0/1614	0.66	0/2202
2	D	0.46	0/1599	0.65	0/2180
All	All	0.53	0/6391	0.71	4/8727 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	66	ARG	NE-CZ-NH1	6.07	123.33	120.30
1	C	66	ARG	NE-CZ-NH2	-6.05	117.28	120.30
1	A	66	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	A	66	ARG	NE-CZ-NH2	-5.13	117.74	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1564	0	1526	5	0
1	C	1529	0	1494	1	0
2	B	1574	0	1518	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1560	0	1506	4	0
3	A	5	0	0	0	0
3	C	5	0	0	0	0
4	A	65	0	0	1	0
4	B	31	0	0	0	0
4	C	64	0	0	0	0
4	D	34	0	0	0	0
All	All	6431	0	6044	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:78:LEU:HD21	2:D:106:VAL:HG22	1.68	0.74
2:B:131:ALA:HB3	2:B:181:LEU:HD12	1.80	0.64
1:A:160:THR:O	1:A:163:VAL:HG12	2.06	0.55
2:B:168:GLN:HE21	2:B:174:ALA:HB2	1.72	0.55
1:A:18:LEU:HD13	1:A:109:VAL:HG11	1.88	0.54
1:A:159:LEU:HD21	1:A:182:VAL:HG21	1.91	0.52
2:D:21:ILE:HD11	2:D:104:VAL:CG2	2.42	0.49
1:A:77:GLN:NE2	4:A:601:HOH:O	2.46	0.48
2:B:165:PRO:HA	2:B:174:ALA:O	2.19	0.42
1:C:29:ILE:HG21	1:C:73:THR:HG22	2.01	0.42
2:B:181:LEU:HD11	2:B:186:TRP:HB2	2.02	0.42
2:D:36:TYR:HE1	2:D:46:VAL:HG22	1.85	0.41
1:A:143:LYS:NZ	1:A:171:GLN:HE22	2.19	0.41
2:D:54:ARG:HD3	2:D:62:PHE:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	202/218 (93%)	191 (95%)	11 (5%)	0	100	100
1	C	194/218 (89%)	187 (96%)	7 (4%)	0	100	100
2	B	209/216 (97%)	202 (97%)	7 (3%)	0	100	100
2	D	205/216 (95%)	197 (96%)	8 (4%)	0	100	100
All	All	810/868 (93%)	777 (96%)	33 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/190 (95%)	179 (99%)	1 (1%)	86	87
1	C	175/190 (92%)	173 (99%)	2 (1%)	73	73
2	B	177/182 (97%)	175 (99%)	2 (1%)	73	73
2	D	175/182 (96%)	174 (99%)	1 (1%)	86	87
All	All	707/744 (95%)	701 (99%)	6 (1%)	81	82

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	TYR
2	B	18	SER
2	B	145	VAL
1	C	14	SER
1	C	50	TYR
2	D	29	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	77	GLN
1	A	95	HIS
2	B	39	HIS
2	B	168	GLN
2	D	109	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PO4	A	501	-	4,4,4	0.80	0	6,6,6	0.49	0
3	PO4	C	501	-	4,4,4	0.91	0	6,6,6	0.78	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	206/218 (94%)	0.34	10 (4%) 29 31	18, 27, 53, 74	0
1	C	200/218 (91%)	0.28	11 (5%) 25 27	18, 26, 44, 66	0
2	B	211/216 (97%)	0.33	12 (5%) 23 25	21, 39, 55, 65	0
2	D	209/216 (96%)	0.60	22 (10%) 6 6	22, 46, 76, 86	0
All	All	826/868 (95%)	0.39	55 (6%) 17 19	18, 33, 65, 86	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	184	GLU	5.8
1	C	184	VAL	5.3
2	D	56	SER	4.8
2	D	76	SER	4.5
1	A	189	LEU	4.4
2	B	188	SER	4.3
2	D	190	ARG	4.2
1	A	191	THR	4.1
2	D	57	GLY	3.8
1	A	41	PRO	3.7
2	D	187	LYS	3.7
1	C	213	ARG	3.6
1	C	136	ALA	3.5
1	A	74	SER	3.4
1	C	185	PRO	3.4
2	B	95	TYR	3.3
1	C	31	SER	3.3
1	A	190	GLY	3.2
2	D	2	SER	3.2
2	D	184	GLU	3.2
2	D	95	TYR	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	167	LYS	3.1
1	C	26	GLY	3.0
1	A	42	GLY	3.0
1	A	192	GLN	3.0
2	B	60	ASP	2.9
2	D	77	GLY	2.9
2	B	153	GLY	2.9
2	B	32	TYR	2.9
2	D	30(A)	GLY	2.8
1	C	42	GLY	2.7
1	A	93	ALA	2.7
1	A	161	SER	2.7
2	D	168	GLN	2.6
2	D	108	GLY	2.6
1	C	97	ASN	2.6
2	B	2	SER	2.6
2	B	169	SER	2.6
2	D	110	PRO	2.5
2	B	56	SER	2.4
2	D	32	TYR	2.4
2	D	112	ALA	2.3
2	D	31	ASN	2.3
2	B	189	HIS	2.3
2	D	60	ASP	2.2
2	D	109	GLN	2.2
1	A	187	SER	2.2
2	D	8	ARG	2.2
2	B	190	ARG	2.2
1	C	29	ILE	2.1
1	C	102	TYR	2.1
1	C	193	THR	2.1
2	D	141	TYR	2.1
2	D	78	LEU	2.0
2	B	181	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	C	501	5/5	0.86	0.21	30,33,33,34	5
3	PO4	A	501	5/5	0.92	0.17	29,29,31,32	5

### 6.5 Other polymers [i](#)

There are no such residues in this entry.