



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 15, 2020 – 12:01 am BST

PDB ID : 3GL2
Title : Crystal structure of dicamba monooxygenase bound to dicamba
Authors : Wilson, M.A.; Dumitru, R.; Jiang, W.Z.; Weeks, D.P.
Deposited on : 2009-03-11
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>13%</div> <div>••</div> </div> </div>
1	B	349	<div> <div>9%</div> <div> <div></div> <div>74%</div> <div>17%</div> <div>••</div> </div> </div>
1	C	349	<div> <div>7%</div> <div> <div></div> <div>75%</div> <div>13%</div> <div>• 10%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DdmC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	340	Total	C	N	O	S	0	0	0
			2633	1659	473	488	13			
1	B	334	Total	C	N	O	S	0	0	0
			2590	1634	463	480	13			
1	C	315	Total	C	N	O	S	0	1	0
			2440	1545	428	454	13			

There are 33 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INSERTION	UNP Q5S3I3
A	2	ALA	-	INSERTION	UNP Q5S3I3
A	341	ARG	-	INSERTION	UNP Q5S3I3
A	342	LEU	-	EXPRESSION TAG	UNP Q5S3I3
A	343	GLU	-	EXPRESSION TAG	UNP Q5S3I3
A	344	HIS	-	EXPRESSION TAG	UNP Q5S3I3
A	345	HIS	-	EXPRESSION TAG	UNP Q5S3I3
A	346	HIS	-	EXPRESSION TAG	UNP Q5S3I3
A	347	HIS	-	EXPRESSION TAG	UNP Q5S3I3
A	348	HIS	-	EXPRESSION TAG	UNP Q5S3I3
A	349	HIS	-	EXPRESSION TAG	UNP Q5S3I3
B	1	MET	-	INSERTION	UNP Q5S3I3
B	2	ALA	-	INSERTION	UNP Q5S3I3
B	341	ARG	-	INSERTION	UNP Q5S3I3
B	342	LEU	-	EXPRESSION TAG	UNP Q5S3I3
B	343	GLU	-	EXPRESSION TAG	UNP Q5S3I3
B	344	HIS	-	EXPRESSION TAG	UNP Q5S3I3
B	345	HIS	-	EXPRESSION TAG	UNP Q5S3I3
B	346	HIS	-	EXPRESSION TAG	UNP Q5S3I3
B	347	HIS	-	EXPRESSION TAG	UNP Q5S3I3
B	348	HIS	-	EXPRESSION TAG	UNP Q5S3I3
B	349	HIS	-	EXPRESSION TAG	UNP Q5S3I3
C	1	MET	-	INSERTION	UNP Q5S3I3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	2	ALA	-	INSERTION	UNP Q5S3I3
C	341	ARG	-	INSERTION	UNP Q5S3I3
C	342	LEU	-	EXPRESSION TAG	UNP Q5S3I3
C	343	GLU	-	EXPRESSION TAG	UNP Q5S3I3
C	344	HIS	-	EXPRESSION TAG	UNP Q5S3I3
C	345	HIS	-	EXPRESSION TAG	UNP Q5S3I3
C	346	HIS	-	EXPRESSION TAG	UNP Q5S3I3
C	347	HIS	-	EXPRESSION TAG	UNP Q5S3I3
C	348	HIS	-	EXPRESSION TAG	UNP Q5S3I3
C	349	HIS	-	EXPRESSION TAG	UNP Q5S3I3

- Molecule 2 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe S 4 2 2	0	0
2	B	1	Total Fe S 4 2 2	0	0
2	C	1	Total Fe S 4 2 2	0	0

- Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Fe 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		

- Molecule 4 is 3,6-dichloro-2-methoxybenzoic acid (three-letter code: D3M) (formula: $C_8H_6Cl_2O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	Cl	O	0	0
			13	8	2	3		
4	B	1	Total	C	Cl	O	0	0
			13	8	2	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	155	Total	O	0	0
			155	155		
5	B	142	Total	O	0	0
			142	142		
5	C	156	Total	O	0	0
			156	156		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

V325	L202	M203	A2	MET
R329	M203	A204	T3	
E332	K205	F4		
Q336	F206			
L337	L207			
E338	ARG			
A339	GLY			
A340	GLY			
ARG	ASN			
LEU	ASN			
GLU	THR			
HIS	P213			
HIS	V214			
HIS	D215			
HIS	A216			
HIS	D217			
HIS	W218			
HIS	D219			
HIS	A233			
HIS	V234			
	A235			
	P236			
	E237			
	P240			
	K241			
	E242			
	Q243			
	S244			
	I245			
	T250			
	E258			
	Y263			
	F264			
	F265			
	I273			
	P276			
	D279			
	S284			
	A287			
	V291			
	K293			
	E293			
	D294			
	K295			
	P315			
	A316			
	M317			
	L157			
	L158			
	GLY			
	HIS			
	ALA			
	GLN			
	TYR			
	VAL			
	HIS			
	ARG			
	ALA			
	ASN			
	ALA			
	GLN			
	THR			
	ASP			
	ALA			
	PHE			
	ASP			
	ARG			
	LEU			
	E178			
	R179			
	E180			
	M192			
	K193			
	I194			
	P195			
	G196			
	I197			
	T198			
	P199			
	S200			
	W201			

4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	81.32Å 81.32Å 159.28Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.00 – 2.10 36.21 – 2.11	Depositor EDS
% Data completeness (in resolution range)	98.4 (36.00-2.10) 98.4 (36.21-2.11)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.190 , 0.247 0.194 , 0.250	Depositor DCC
R_{free} test set	3338 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	49.9	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l 0.034 for h,-h-k,-l 0.013 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8156	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: D3M, FE, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/2698	0.53	0/3674
1	B	0.47	0/2654	0.55	0/3612
1	C	0.45	0/2502	0.54	0/3405
All	All	0.46	0/7854	0.54	0/10691

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
1	C	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	340	ALA	Peptide
1	B	205	LYS	Peptide
1	B	213	PRO	Peptide
1	C	339	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2633	0	2566	35	0
1	B	2590	0	2521	76	0
1	C	2440	0	2386	40	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	13	0	5	0	0
4	B	13	0	5	0	0
5	A	155	0	0	3	0
5	B	142	0	0	4	0
5	C	156	0	0	7	0
All	All	8156	0	7483	148	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 148 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:PRO:O	1:A:239:THR:HG23	1.42	1.20
1:C:158:LEU:CD1	1:C:158:LEU:C	2.19	1.10
1:B:174:PHE:HA	1:B:177:LEU:CD2	1.85	1.07
1:B:174:PHE:HA	1:B:177:LEU:HD21	1.38	1.05
1:B:291:VAL:HG12	1:B:291:VAL:O	1.54	1.05

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/349 (97%)	324 (96%)	14 (4%)	0	100	100
1	B	330/349 (95%)	303 (92%)	20 (6%)	7 (2%)	7	3
1	C	310/349 (89%)	292 (94%)	16 (5%)	2 (1%)	25	21
All	All	978/1047 (93%)	919 (94%)	50 (5%)	9 (1%)	17	12

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	166	ARG
1	B	340	ALA
1	B	237	GLU
1	B	292	LYS
1	C	180	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/284 (97%)	266 (97%)	9 (3%)	38	40
1	B	271/284 (95%)	254 (94%)	17 (6%)	18	15
1	C	258/284 (91%)	247 (96%)	11 (4%)	29	29
All	All	804/852 (94%)	767 (95%)	37 (5%)	27	26

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	202	LEU
1	B	276	PRO
1	C	241	LYS
1	B	214	VAL
1	B	217	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	D3M	A	600	-	11,13,13	1.35	3 (27%)	13,18,18	0.99	0
2	FES	B	500	1	0,4,4	0.00	-	-		
2	FES	A	500	1	0,4,4	0.00	-	-		
2	FES	C	500	1	0,4,4	0.00	-	-		
4	D3M	B	600	-	11,13,13	1.45	3 (27%)	13,18,18	1.22	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	D3M	A	600	-	-	0/2/6/6	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FES	B	500	1	-	-	0/1/1/1
2	FES	A	500	1	-	-	0/1/1/1
4	D3M	B	600	-	-	0/2/6/6	0/1/1/1
2	FES	C	500	1	-	-	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	600	D3M	C5-CL1	2.77	1.80	1.73
4	B	600	D3M	C1-C7	2.69	1.50	1.47
4	A	600	D3M	C5-CL1	2.28	1.79	1.73
4	B	600	D3M	C2-CL2	2.23	1.78	1.73
4	A	600	D3M	C1-C7	2.19	1.49	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	600	D3M	C8-O3-C6	2.97	122.92	114.78
4	B	600	D3M	C6-C5-CL1	2.20	121.28	118.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/349 (97%)	-0.06	9 (2%) 56 61	41, 56, 86, 110	0
1	B	334/349 (95%)	0.28	32 (9%) 8 10	39, 57, 112, 138	0
1	C	315/349 (90%)	0.17	23 (7%) 15 19	39, 55, 116, 145	0
All	All	989/1047 (94%)	0.13	64 (6%) 18 23	39, 56, 107, 145	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	201	VAL	6.8
1	C	198	THR	6.3
1	B	214	VAL	5.8
1	B	194	ILE	5.2
1	C	207	LEU	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	D3M	B	600	13/13	0.94	0.11	57,72,76,79	0
4	D3M	A	600	13/13	0.97	0.09	46,53,60,69	0
2	FES	A	500	4/4	0.99	0.10	46,47,49,53	0
2	FES	C	500	4/4	0.99	0.16	41,43,44,50	0
3	FE	A	501	1/1	0.99	0.17	48,48,48,48	0
3	FE	B	501	1/1	0.99	0.16	64,64,64,64	0
2	FES	B	500	4/4	1.00	0.08	41,44,44,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.