



wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 05:34 pm BST

PDB ID : 1GPD
Title : STUDIES OF ASYMMETRY IN THE THREE-DIMENSIONAL STRUCTURE OF LOBSTER D-GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE
Authors : Moras, D.; Olsen, K.W.; Sabesan, M.N.; Buehner, M.; Ford, G.C.; Rossmann, M.G.
Deposited on : 1975-07-01
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	G	334	
1	R	334	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	R	338	-	-	X	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5112 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	334	Total	C	N	O	S	0	0	0
			2510	1593	419	483	15			
1	R	334	Total	C	N	O	S	0	0	0
			2510	1593	419	483	15			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).






Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	R	1	Total	P	0	0
			1	1		
3	R	1	Total	P	0	0
			1	1		
3	R	1	Total	P	0	0
			1	1		
3	R	1	Total	P	0	0
			1	1		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: D-GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE



• Molecule 1: D-GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE



K302	T303	F304	V305	K306	V307	V308	S309	W310	Y311	D312	N313	E314	F315	G316	Y317	S318	Q319	R320	V321	I322	D323	I324	L325	K326	H327	M328	Q329	K330	V331	D332	S333	A334
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	149.00Å 139.10Å 80.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.90	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	unknown	Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5112	wwPDB-VP
Average B, all atoms (Å ²)	0.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, NAD, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	G	1.18	30/2554 (1.2%)	2.05	168/3453 (4.9%)
1	R	1.18	30/2554 (1.2%)	1.88	130/3453 (3.8%)
All	All	1.18	60/5108 (1.2%)	1.97	298/6906 (4.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	5
1	R	0	7
All	All	0	12

The worst 5 of 60 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	84	TRP	NE1-CE2	-7.38	1.27	1.37
1	R	84	TRP	NE1-CE2	-7.35	1.27	1.37
1	R	310	TRP	NE1-CE2	-7.34	1.28	1.37
1	R	193	TRP	NE1-CE2	-7.33	1.28	1.37
1	G	310	TRP	NE1-CE2	-7.31	1.28	1.37

The worst 5 of 298 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	277	ASP	N-CA-C	13.71	148.02	111.00
1	G	300	LEU	N-CA-C	13.61	147.75	111.00
1	R	198	GLY	N-CA-C	12.19	143.56	113.10
1	R	195	GLY	N-CA-C	12.14	143.44	113.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	198	GLY	N-CA-C	12.06	143.25	113.10

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	13	ARG	Sidechain
1	G	194	ARG	Sidechain
1	G	231	ARG	Sidechain
1	G	245	ARG	Sidechain
1	G	320	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2510	0	2519	1064	7
1	R	2510	0	2522	1004	7
2	G	44	0	26	20	0
2	R	44	0	26	8	0
3	R	4	0	0	8	0
All	All	5112	0	5093	2047	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 201.

The worst 5 of 2047 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:117:VAL:HG21	1:G:328:MET:SD	1.23	1.70
1:R:324:LEU:HD11	1:R:328:MET:SD	1.24	1.66
1:G:173:THR:HG22	1:G:228:MET:SD	1.39	1.60
1:R:137:TYR:CD1	1:R:331:VAL:HG11	1.47	1.47
1:R:5:ILE:CG1	1:R:30:VAL:HG13	1.02	1.47

The worst 5 of 7 symmetry-related close contacts are listed below. The label for Atom-2 includes

the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:135:GLU:CB	1:R:220:GLU:OE2[4_555]	1.21	0.99
1:G:135:GLU:CG	1:R:220:GLU:OE2[4_555]	1.26	0.94
1:G:135:GLU:N	1:R:220:GLU:OE1[4_555]	1.32	0.88
1:G:135:GLU:CA	1:R:220:GLU:OE1[4_555]	1.53	0.67
1:G:135:GLU:CB	1:R:220:GLU:CD[4_555]	1.61	0.59

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	332/334 (99%)	178 (54%)	73 (22%)	81 (24%)	0	0
1	R	332/334 (99%)	167 (50%)	72 (22%)	93 (28%)	0	0
All	All	664/668 (99%)	345 (52%)	145 (22%)	174 (26%)	0	0

5 of 174 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	2	LYS
1	G	10	ARG
1	G	13	ARG
1	G	25	ALA
1	G	28	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	271/271 (100%)	149 (55%)	122 (45%)	0	0
1	R	271/271 (100%)	140 (52%)	131 (48%)	0	0
All	All	542/542 (100%)	289 (53%)	253 (47%)	0	0

5 of 253 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	313	ASN
1	R	47	ASP
1	R	291	ILE
1	G	324	LEU
1	R	15	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	327	HIS
1	R	26	GLN
1	R	299	GLN
1	G	313	ASN
1	G	319	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are modelled with single atom - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAD	G	335	-	42,48,48	0.73	1 (2%)	50,73,73	0.88	3 (6%)
2	NAD	R	339	-	42,48,48	0.72	1 (2%)	50,73,73	1.70	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	G	335	-	-	9/26/62/62	0/5/5/5
2	NAD	R	339	-	-	12/26/62/62	0/5/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	R	339	NAD	C8A-N7A	-2.55	1.30	1.34
2	G	335	NAD	C8A-N7A	-2.53	1.30	1.34

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	R	339	NAD	O2A-PA-O5B	7.85	144.21	107.75
2	R	339	NAD	O2A-PA-O1A	-7.30	76.17	112.24
2	G	335	NAD	O2A-PA-O5B	2.61	119.85	107.75
2	G	335	NAD	O4D-C1D-C2D	-2.41	103.41	106.93
2	R	339	NAD	O4D-C1D-C2D	-2.26	103.62	106.93

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	G	335	NAD	C5B-O5B-PA-O1A
2	G	335	NAD	C5B-O5B-PA-O2A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	G	335	NAD	C5D-O5D-PN-O3
2	G	335	NAD	C5D-O5D-PN-O1N
2	G	335	NAD	O4D-C4D-C5D-O5D


There are no ring outliers.

2 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	335	NAD	20	0
2	R	339	NAD	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.