



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 08:35 pm BST

PDB ID : 2GQN  
Title : Cystathionine Beta-Lyase (CBL) from Escherichia Coli in complex with N-Hydrazinocarbonylmethyl-2-Nitro-Benzamide  
Authors : Summerfield, R.; Junop, M.S.  
Deposited on : 2006-04-21  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	415	 80% 13% 6%
1	B	415	 81% 13% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BLP	A	700	-	-	X	-
2	BLP	B	700	-	-	X	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cystathionine beta-lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	391	Total	C	N	O	S	0	0	0
			3012	1912	524	562	14			
1	B	392	Total	C	N	O	S	0	0	0
			3021	1918	526	563	14			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	CLONING ARTIFACT	UNP P06721
A	-18	GLY	-	CLONING ARTIFACT	UNP P06721
A	-17	SER	-	CLONING ARTIFACT	UNP P06721
A	-16	SER	-	CLONING ARTIFACT	UNP P06721
A	-15	HIS	-	EXPRESSION TAG	UNP P06721
A	-14	HIS	-	EXPRESSION TAG	UNP P06721
A	-13	HIS	-	EXPRESSION TAG	UNP P06721
A	-12	HIS	-	EXPRESSION TAG	UNP P06721
A	-11	HIS	-	EXPRESSION TAG	UNP P06721
A	-10	HIS	-	EXPRESSION TAG	UNP P06721
A	-9	SER	-	CLONING ARTIFACT	UNP P06721
A	-8	SER	-	CLONING ARTIFACT	UNP P06721
A	-7	GLY	-	CLONING ARTIFACT	UNP P06721
A	-6	LEU	-	CLONING ARTIFACT	UNP P06721
A	-5	VAL	-	CLONING ARTIFACT	UNP P06721
A	-4	PRO	-	CLONING ARTIFACT	UNP P06721
A	-3	ARG	-	CLONING ARTIFACT	UNP P06721
A	-2	GLY	-	CLONING ARTIFACT	UNP P06721
A	-1	SER	-	CLONING ARTIFACT	UNP P06721
A	0	HIS	-	CLONING ARTIFACT	UNP P06721
B	-19	MET	-	CLONING ARTIFACT	UNP P06721
B	-18	GLY	-	CLONING ARTIFACT	UNP P06721
B	-17	SER	-	CLONING ARTIFACT	UNP P06721
B	-16	SER	-	CLONING ARTIFACT	UNP P06721
B	-15	HIS	-	EXPRESSION TAG	UNP P06721

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	EXPRESSION TAG	UNP P06721
B	-13	HIS	-	EXPRESSION TAG	UNP P06721
B	-12	HIS	-	EXPRESSION TAG	UNP P06721
B	-11	HIS	-	EXPRESSION TAG	UNP P06721
B	-10	HIS	-	EXPRESSION TAG	UNP P06721
B	-9	SER	-	CLONING ARTIFACT	UNP P06721
B	-8	SER	-	CLONING ARTIFACT	UNP P06721
B	-7	GLY	-	CLONING ARTIFACT	UNP P06721
B	-6	LEU	-	CLONING ARTIFACT	UNP P06721
B	-5	VAL	-	CLONING ARTIFACT	UNP P06721
B	-4	PRO	-	CLONING ARTIFACT	UNP P06721
B	-3	ARG	-	CLONING ARTIFACT	UNP P06721
B	-2	GLY	-	CLONING ARTIFACT	UNP P06721
B	-1	SER	-	CLONING ARTIFACT	UNP P06721
B	0	HIS	-	CLONING ARTIFACT	UNP P06721

- Molecule 2 is (5-HYDROXY-6-METHYL-4-((2-(2-(2-NITROBENZAMIDO)ACETYL)HYDRAZINYL)METHYL)PYRIDIN-3-YL)METHYL DIHYDROGEN PHOSPHATE (three-letter code: BLP) (formula: C<sub>17</sub>H<sub>20</sub>N<sub>5</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			32	17	5	9	1		
2	B	1	Total	C	N	O	P	0	0
			32	17	5	9	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	474	Total 474	O 474	0	0
3	B	473	Total 473	O 473	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cystathionine beta-lyase



- Molecule 1: Cystathionine beta-lyase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.64Å 153.01Å 151.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.80 44.78 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.6 (50.00-1.80) 99.6 (44.78-1.80)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.51 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.174 , 0.227 0.201 , 0.253	Depositor DCC
$R_{free}$ test set	3250 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.4	Xtriage
Anisotropy	0.679	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 34.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7044	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.04 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.9753e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.85	2/3075 (0.1%)	0.84	1/4169 (0.0%)
1	B	0.84	1/3084 (0.0%)	0.83	2/4180 (0.0%)
All	All	0.84	3/6159 (0.0%)	0.84	3/8349 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	363	GLU	CG-CD	6.00	1.60	1.51
1	A	363	GLU	CB-CG	5.08	1.61	1.52
1	B	298	GLU	CB-CG	5.02	1.61	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	285	ARG	NE-CZ-NH2	-7.86	116.37	120.30
1	B	285	ARG	NE-CZ-NH1	5.67	123.13	120.30
1	A	142	LYS	CB-CA-C	-5.00	100.39	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	31	ALA	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3012	0	2988	47	0
1	B	3021	0	3001	53	0
2	A	32	0	17	12	0
2	B	32	0	18	16	0
3	A	474	0	0	11	0
3	B	473	0	0	8	0
All	All	7044	0	6024	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:338:TYR:HB3	2:B:700:BLP:C9	1.60	1.31
1:B:338:TYR:HB3	2:B:700:BLP:H92	1.16	1.08
1:B:280:HIS:HD2	1:B:282:GLN:H	1.19	0.87
1:A:326:ASN:HB2	3:A:898:HOH:O	1.75	0.87
1:B:338:TYR:CB	2:B:700:BLP:H92	2.05	0.86
1:B:338:TYR:CB	2:B:700:BLP:C9	2.51	0.85
1:B:61:THR:H	1:B:64:HIS:HD2	1.22	0.84
1:B:111:TYR:CZ	2:B:700:BLP:H72	2.13	0.83
1:A:339:SER:H	2:A:700:BLP:H92	1.45	0.82
1:A:64:HIS:HE1	1:A:244:VAL:O	1.64	0.80
1:B:288:HIS:HD2	1:B:290:ALA:H	1.26	0.80
1:B:339:SER:H	2:B:700:BLP:H92	1.44	0.80
1:A:97:PHE:O	1:A:149:LYS:HE2	1.83	0.79
1:B:338:TYR:HB3	2:B:700:BLP:H91	1.64	0.79
1:B:61:THR:H	1:B:64:HIS:CD2	2.02	0.78
1:A:210:LYS:HZ1	2:A:700:BLP:H71	1.49	0.77
1:A:61:THR:H	1:A:64:HIS:HD2	1.31	0.77

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:HIS:HD2	1:A:290:ALA:H	1.34	0.74
1:B:131:TRP:HD1	1:B:361:GLN:HE21	1.36	0.74
1:B:149:LYS:HD2	3:B:1008:HOH:O	1.87	0.74
1:B:285:ARG:HD3	3:B:911:HOH:O	1.89	0.73
1:B:64:HIS:HE1	1:B:244:VAL:O	1.72	0.72
1:A:61:THR:H	1:A:64:HIS:CD2	2.08	0.71
1:A:210:LYS:NZ	2:A:700:BLP:H71	2.05	0.71
1:A:280:HIS:HD2	1:A:282:GLN:H	1.36	0.71
1:B:339:SER:H	2:B:700:BLP:C9	2.04	0.69
1:A:301:LYS:HE3	3:A:1103:HOH:O	1.93	0.69
1:B:339:SER:N	2:B:700:BLP:H92	2.09	0.68
1:A:139:ASP:HB3	3:A:917:HOH:O	1.93	0.67
1:A:356:ALA:HA	1:A:364:ILE:HD13	1.77	0.67
1:A:285:ARG:HD2	3:A:822:HOH:O	1.96	0.66
1:B:280:HIS:CD2	1:B:282:GLN:H	2.09	0.64
1:A:339:SER:N	2:A:700:BLP:H92	2.11	0.63
1:A:64:HIS:CE1	1:A:244:VAL:O	2.52	0.61
1:B:288:HIS:CD2	1:B:290:ALA:H	2.14	0.61
1:B:338:TYR:CD1	2:B:700:BLP:H91	2.36	0.61
1:B:338:TYR:HB3	2:B:700:BLP:N4	2.14	0.61
1:A:338:TYR:HB3	2:A:700:BLP:HN4	1.65	0.60
1:B:267:HIS:CD2	1:B:307:SER:H	2.19	0.60
1:A:130:SER:HB2	1:A:361:GLN:HE22	1.66	0.60
1:A:97:PHE:HE2	1:A:233:LEU:HD22	1.65	0.60
1:A:359:ARG:HG3	1:A:364:ILE:HD11	1.83	0.60
2:A:700:BLP:O8	2:A:700:BLP:O6	2.20	0.59
1:B:338:TYR:CB	2:B:700:BLP:H91	2.28	0.59
1:B:47:ARG:HD3	1:B:47:ARG:O	2.02	0.58
1:B:225:CYS:SG	1:B:233:LEU:HD23	2.44	0.58
1:A:339:SER:H	2:A:700:BLP:C9	2.16	0.57
1:B:361:GLN:HA	3:B:821:HOH:O	2.04	0.57
1:B:339:SER:H	2:B:700:BLP:C8	2.16	0.57
1:A:338:TYR:CD1	2:A:700:BLP:N4	2.73	0.57
1:A:355:ILE:HG22	1:A:364:ILE:HD11	1.87	0.57
1:B:64:HIS:CE1	1:B:244:VAL:O	2.57	0.55
2:B:700:BLP:O8	2:B:700:BLP:O6	2.23	0.55
1:A:280:HIS:CD2	1:A:282:GLN:H	2.22	0.55
1:B:111:TYR:OH	2:B:700:BLP:H72	2.06	0.55
1:A:174:ARG:NH1	3:A:1113:HOH:O	2.41	0.55
1:B:143:HIS:CD2	3:B:787:HOH:O	2.60	0.54
1:A:39:VAL:O	1:A:43:LYS:HD2	2.08	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:111:TYR:CD2	1:B:113:PRO:HD2	2.43	0.53
1:B:95:LEU:HD12	1:B:98:ILE:HD11	1.91	0.53
1:B:15:ARG:NH2	3:B:967:HOH:O	2.41	0.53
1:A:359:ARG:HG3	1:A:364:ILE:CD1	2.40	0.51
1:A:338:TYR:HD1	2:A:700:BLP:N4	2.09	0.51
1:B:289:PRO:O	1:B:297:HIS:HD2	1.94	0.50
1:A:318:LYS:HE2	1:A:353:GLU:OE2	2.12	0.50
1:B:21:LEU:HD12	1:B:27:VAL:HG11	1.93	0.50
1:B:102:ASP:OD2	1:B:149:LYS:NZ	2.40	0.49
1:A:289:PRO:O	1:A:297:HIS:HD2	1.94	0.49
1:A:111:TYR:CZ	2:A:700:BLP:H72	2.48	0.49
1:B:370:LEU:HD23	1:B:370:LEU:C	2.32	0.49
1:B:143:HIS:HD2	3:B:787:HOH:O	1.95	0.48
1:A:285:ARG:HD3	3:A:1147:HOH:O	2.13	0.48
1:B:338:TYR:HD1	2:B:700:BLP:H91	1.77	0.48
1:A:372:ARG:HH22	2:A:700:BLP:H91	1.78	0.48
1:B:47:ARG:HA	1:B:47:ARG:HE	1.79	0.48
1:B:77:GLY:HA3	1:B:224:VAL:CG1	2.44	0.48
1:A:95:LEU:HD12	1:A:98:ILE:HD11	1.94	0.47
1:B:120:LYS:HE2	3:B:980:HOH:O	2.13	0.47
1:A:372:ARG:HH22	2:A:700:BLP:C9	2.28	0.47
1:A:131:TRP:CD1	1:A:360:PRO:HB2	2.52	0.45
3:A:909:HOH:O	1:B:29:GLN:HG3	2.16	0.45
1:A:65:PHE:HA	1:A:68:GLN:HE21	1.82	0.45
1:A:326:ASN:CB	3:A:898:HOH:O	2.48	0.44
1:B:141:VAL:HG13	1:B:172:ALA:HB1	1.99	0.44
1:A:153:LEU:HB3	1:A:164:HIS:CD2	2.53	0.43
1:A:270:SER:OG	1:A:288:HIS:HE1	2.02	0.43
1:A:15:ARG:HD2	1:A:24:VAL:O	2.18	0.43
1:B:106:MET:O	1:B:131:TRP:HA	2.18	0.43
1:A:21:LEU:HD12	1:A:27:VAL:HG11	2.00	0.42
1:A:43:LYS:N	1:A:43:LYS:HE3	2.34	0.42
1:B:211:TYR:CE1	1:B:341:GLY:HA2	2.53	0.42
1:A:301:LYS:CE	3:A:1103:HOH:O	2.60	0.42
1:B:208:ALA:HB3	1:B:220:ILE:HG23	2.00	0.42
1:A:354:HIS:HB3	3:A:912:HOH:O	2.19	0.42
1:B:131:TRP:CD1	1:B:131:TRP:N	2.84	0.42
1:B:131:TRP:HD1	1:B:361:GLN:NE2	2.10	0.41
1:B:131:TRP:H	1:B:361:GLN:NE2	2.17	0.41
1:B:142:LYS:HE3	3:B:1155:HOH:O	2.19	0.41
1:B:130:SER:HB2	1:B:361:GLN:HE22	1.85	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106:MET:O	1:A:131:TRP:HA	2.20	0.40
1:B:104:VAL:O	1:B:129:THR:HA	2.20	0.40
1:A:51:ASN:ND2	3:A:1126:HOH:O	2.47	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	389/415 (94%)	376 (97%)	12 (3%)	1 (0%)	41	27
1	B	390/415 (94%)	374 (96%)	15 (4%)	1 (0%)	41	27
All	All	779/830 (94%)	750 (96%)	27 (4%)	2 (0%)	41	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	LYS
1	B	210	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/339 (94%)	310 (97%)	9 (3%)	43	30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	320/339 (94%)	311 (97%)	9 (3%)	43	30
All	All	639/678 (94%)	621 (97%)	18 (3%)	43	30

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	43	LYS
1	A	136	ILE
1	A	150	ILE
1	A	186	ASN
1	A	233	LEU
1	A	251	ILE
1	A	363	GLU
1	A	395	VAL
1	B	4	LYS
1	B	5	LYS
1	B	18	LYS
1	B	186	ASN
1	B	233	LEU
1	B	239	LEU
1	B	317	LYS
1	B	363	GLU
1	B	395	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	ASN
1	A	51	ASN
1	A	64	HIS
1	A	68	GLN
1	A	143	HIS
1	A	186	ASN
1	A	242	GLN
1	A	267	HIS
1	A	280	HIS
1	A	282	GLN
1	A	287	ASN
1	A	288	HIS
1	A	297	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	361	GLN
1	B	51	ASN
1	B	64	HIS
1	B	68	GLN
1	B	143	HIS
1	B	186	ASN
1	B	242	GLN
1	B	267	HIS
1	B	280	HIS
1	B	288	HIS
1	B	297	HIS
1	B	361	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BLP	B	700	-	31,33,33	1.78	7 (22%)	39,46,46	2.46	13 (33%)
2	BLP	A	700	-	31,33,33	2.28	8 (25%)	39,46,46	2.49	11 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BLP	B	700	-	-	4/23/25/25	0/2/2/2
2	BLP	A	700	-	-	7/23/25/25	0/2/2/2

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	700	BLP	N3-N2	6.53	1.46	1.41
2	A	700	BLP	C8-N3	6.29	1.42	1.34
2	B	700	BLP	C8-N3	5.61	1.41	1.34
2	A	700	BLP	O7-N5	3.96	1.29	1.22
2	A	700	BLP	C16-N5	-3.61	1.39	1.45
2	B	700	BLP	C16-N5	-3.60	1.39	1.45
2	B	700	BLP	O7-N5	3.10	1.28	1.22
2	B	700	BLP	C2-N1	3.00	1.39	1.33
2	A	700	BLP	C9-C8	-2.81	1.43	1.52
2	B	700	BLP	C1-C2	2.57	1.54	1.50
2	B	700	BLP	N3-N2	2.17	1.43	1.41
2	A	700	BLP	O6-C10	2.15	1.27	1.23
2	B	700	BLP	C9-N4	2.13	1.50	1.45
2	A	700	BLP	C10-N4	2.10	1.38	1.33
2	A	700	BLP	C17-C6	2.01	1.43	1.40

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	700	BLP	O5-C8-N3	9.01	133.73	122.81
2	A	700	BLP	C8-N3-N2	-6.09	112.28	120.03
2	A	700	BLP	C8-C9-N4	-5.83	96.80	113.17
2	B	700	BLP	C9-C8-N3	-5.79	108.47	115.20
2	A	700	BLP	O5-C8-N3	5.11	129.00	122.81
2	A	700	BLP	C9-N4-C10	4.70	132.52	121.33
2	B	700	BLP	C9-N4-C10	4.51	132.07	121.33
2	A	700	BLP	O5-C8-C9	-4.44	111.56	120.75
2	A	700	BLP	C7-C6-C17	4.21	124.55	120.04
2	B	700	BLP	O3-P1-O1	-3.90	96.35	106.73
2	A	700	BLP	C9-C8-N3	3.44	119.20	115.20
2	B	700	BLP	O3-P1-O2	3.26	120.09	107.64
2	A	700	BLP	C3-C4-C6	3.16	120.35	118.12

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	700	BLP	C8-N3-N2	-2.55	116.79	120.03
2	B	700	BLP	O6-C10-N4	-2.52	117.58	122.61
2	B	700	BLP	O1-P1-O4	-2.51	99.43	106.47
2	A	700	BLP	C13-C12-C11	2.47	124.48	119.81
2	A	700	BLP	O3-P1-O4	2.46	120.31	110.68
2	B	700	BLP	C4-C3-N1	-2.43	119.78	123.82
2	B	700	BLP	O3-P1-O4	2.42	120.15	110.68
2	B	700	BLP	C3-N1-C2	2.21	123.26	119.17
2	A	700	BLP	O6-C10-N4	-2.15	118.33	122.61
2	B	700	BLP	C17-C2-N1	-2.06	118.10	120.77
2	B	700	BLP	C11-C10-N4	2.04	121.36	117.36

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	700	BLP	C17-C6-C7-N2
2	B	700	BLP	C4-C6-C7-N2
2	A	700	BLP	C4-C6-C7-N2
2	B	700	BLP	O5-C8-C9-N4
2	A	700	BLP	O5-C8-C9-N4
2	B	700	BLP	N3-C8-C9-N4
2	A	700	BLP	C6-C4-C5-O1
2	A	700	BLP	N3-C8-C9-N4
2	A	700	BLP	C7-N2-N3-C8
2	A	700	BLP	C3-C4-C5-O1
2	A	700	BLP	C17-C6-C7-N2


There are no ring outliers.

2 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	700	BLP	16	0
2	A	700	BLP	12	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	391/415 (94%)	-0.05	1 (0%) 94 92	11, 17, 26, 34	0
1	B	392/415 (94%)	-0.04	6 (1%) 73 70	11, 17, 27, 42	0
All	All	783/830 (94%)	-0.05	7 (0%) 84 82	11, 17, 27, 42	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	4	LYS	2.9
1	A	326	ASN	2.5
1	B	395	VAL	2.5
1	B	220	ILE	2.4
1	B	251	ILE	2.2
1	B	325	ALA	2.1
1	B	44	HIS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BLP	A	700	32/32	0.75	0.32	5,17,23,27	0
2	BLP	B	700	32/32	0.77	0.29	5,16,20,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around BLP B 700:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.