



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 02:19 pm BST

PDB ID : 3GQT  
Title : Crystal structure of glutaryl-CoA dehydrogenase from Burkholderia pseudomallei with fragment (1,4-dimethyl-1,2,3,4-tetrahydroquinoxalin-6-yl)methylamine  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2009-03-24  
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	399	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>9%</div> <div></div> </div> <div></div> </div>
1	B	399	<div> <div>%</div> <div> <div></div> <div>85%</div> <div>8%</div> <div>7%</div> </div> <div></div> </div>
1	C	399	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>8%</div> <div></div> </div> <div></div> </div>
1	D	399	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>9%</div> <div>6%</div> </div> <div></div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11853 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutaryl-CoA dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	383	Total	C	N	O	S	0	0	0
			2903	1833	510	544	16			
1	B	372	Total	C	N	O	S	0	0	0
			2837	1798	498	525	16			
1	C	385	Total	C	N	O	S	0	0	0
			2910	1839	512	543	16			
1	D	376	Total	C	N	O	S	0	0	0
			2875	1820	508	531	16			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP Q3JP94
A	-2	PRO	-	EXPRESSION TAG	UNP Q3JP94
A	-1	GLY	-	EXPRESSION TAG	UNP Q3JP94
A	0	SER	-	EXPRESSION TAG	UNP Q3JP94
B	-3	GLY	-	EXPRESSION TAG	UNP Q3JP94
B	-2	PRO	-	EXPRESSION TAG	UNP Q3JP94
B	-1	GLY	-	EXPRESSION TAG	UNP Q3JP94
B	0	SER	-	EXPRESSION TAG	UNP Q3JP94
C	-3	GLY	-	EXPRESSION TAG	UNP Q3JP94
C	-2	PRO	-	EXPRESSION TAG	UNP Q3JP94
C	-1	GLY	-	EXPRESSION TAG	UNP Q3JP94
C	0	SER	-	EXPRESSION TAG	UNP Q3JP94
D	-3	GLY	-	EXPRESSION TAG	UNP Q3JP94
D	-2	PRO	-	EXPRESSION TAG	UNP Q3JP94
D	-1	GLY	-	EXPRESSION TAG	UNP Q3JP94
D	0	SER	-	EXPRESSION TAG	UNP Q3JP94

- Molecule 2 is 1-(1,4-dimethyl-1,2,3,4-tetrahydroquinoxalin-6-yl)methanamine (three-letter code: UFO) (formula: C<sub>11</sub>H<sub>17</sub>N<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			14	11	3		
2	B	1	Total	C	N	0	0
			14	11	3		
2	D	1	Total	C	N	0	0
			14	11	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	86	Total	O	0	0
			86	86		
3	B	74	Total	O	0	0
			74	74		
3	C	67	Total	O	0	0
			67	67		
3	D	59	Total	O	0	0
			59	59		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Glutaryl-CoA dehydrogenase




#### • Molecule 1: Glutaryl-CoA dehydrogenase




#### • Molecule 1: Glutaryl-CoA dehydrogenase



#### • Molecule 1: Glutaryl-CoA dehydrogenase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.42Å 106.37Å 144.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.99 46.68 – 1.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-1.99) 99.9 (46.68-1.99)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.35 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, $R_{free}$	0.200 , 0.238 0.198 , 0.237	Depositor DCC
$R_{free}$ test set	5175 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.0	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 38.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11853	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UFO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.66	0/2957	0.69	1/4000 (0.0%)
1	B	0.69	0/2890	0.72	0/3908
1	C	0.72	1/2966 (0.0%)	0.70	0/4015
1	D	0.71	0/2927	0.69	0/3953
All	All	0.69	1/11740 (0.0%)	0.70	1/15876 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	192	GLU	CB-CG	5.07	1.61	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	87	ARG	NE-CZ-NH2	-6.67	116.97	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2903	0	2870	22	0
1	B	2837	0	2826	20	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2910	0	2876	22	0
1	D	2875	0	2878	26	0
2	A	14	0	17	3	0
2	B	14	0	17	0	0
2	D	14	0	17	1	0
3	A	86	0	0	1	0
3	B	74	0	0	5	0
3	C	67	0	0	2	0
3	D	59	0	0	0	0
All	All	11853	0	11501	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:247:PHE:HZ	1:A:324:VAL:HG11	1.18	1.00
1:B:331:LYS:NZ	1:B:335:CYS:SG	2.42	0.93
1:A:247:PHE:CZ	1:A:324:VAL:HG11	2.07	0.88
1:B:16:GLN:HG2	3:B:432:HOH:O	1.74	0.88
1:C:146:ASP:CB	3:C:434:HOH:O	2.32	0.77
1:B:291:LEU:HD23	1:C:291:LEU:HD23	1.67	0.75
3:B:410:HOH:O	1:D:372:THR:HG21	1.87	0.73
1:D:331:LYS:NZ	1:D:335:CYS:SG	2.60	0.70
1:C:99:MET:CE	1:C:219:SER:HA	2.21	0.70
1:B:216:LEU:H	1:B:366:ASN:HD22	1.41	0.69
1:C:99:MET:HE3	1:C:219:SER:HA	1.76	0.68
1:A:43:GLU:OE2	1:A:46:ARG:NH1	2.29	0.66
1:B:99:MET:HE3	3:B:451:HOH:O	1.95	0.66
1:A:117:ASP:O	1:A:121:GLU:HG2	1.97	0.64
1:B:216:LEU:H	1:B:366:ASN:ND2	1.96	0.63
1:D:50:THR:HG21	1:D:220:ILE:HB	1.80	0.63
3:B:410:HOH:O	1:D:372:THR:CG2	2.45	0.62
1:C:54:ILE:HD12	1:C:54:ILE:H	1.70	0.56
1:A:171:THR:O	1:A:172:ASN:HB2	2.05	0.56
1:D:372:THR:O	1:D:372:THR:HG23	2.04	0.56
1:B:157:VAL:HB	1:B:158:PRO:HD2	1.89	0.54
1:D:66:PRO:HA	1:D:75:PRO:HG2	1.89	0.54
1:B:21:ASP:O	1:B:25:VAL:HG23	2.07	0.54
1:D:247:PHE:CE2	1:D:324:VAL:HG11	2.42	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:247:PHE:HZ	1:A:324:VAL:CG1	2.07	0.54
1:C:54:ILE:HG22	1:C:58:MET:HE2	1.90	0.54
1:D:20:ASP:O	1:D:24:MET:HG3	2.07	0.53
1:D:51:ASP:OD1	1:D:53:ALA:HB3	2.08	0.53
1:B:102:VAL:HG21	1:B:171:THR:HB	1.89	0.53
1:C:99:MET:HE1	1:C:219:SER:HA	1.90	0.53
1:A:61:ILE:HG13	1:A:63:LEU:HG	1.91	0.53
1:B:43:GLU:OE1	1:B:46:ARG:NH1	2.42	0.53
1:D:169:TRP:HE3	1:D:221:THR:HG21	1.74	0.52
1:A:168:MET:HG3	1:A:169:TRP:CD1	2.44	0.52
1:B:61:ILE:HG13	1:B:63:LEU:HG	1.93	0.51
1:C:54:ILE:HG22	1:C:58:MET:CE	2.41	0.50
1:C:168:MET:HG3	1:C:169:TRP:CD1	2.46	0.50
1:B:126:LYS:HD2	1:B:132:TRP:CE2	2.47	0.50
1:B:51:ASP:O	1:B:54:ILE:HD12	2.11	0.50
1:D:315:ARG:O	1:D:319:GLU:HG2	2.12	0.50
1:C:126:LYS:HD2	1:C:132:TRP:CE2	2.47	0.49
1:A:290:GLN:HE22	1:D:289:ASN:HD21	1.61	0.48
1:B:290:GLN:HE22	1:C:289:ASN:HB3	1.79	0.48
1:A:21:ASP:O	1:A:25:VAL:HG23	2.14	0.48
1:A:352:ASN:HD22	2:A:1000:UFO:HN14	1.63	0.47
1:D:247:PHE:CZ	1:D:324:VAL:HG11	2.50	0.47
1:C:156:LYS:NZ	1:C:233:GLU:OE1	2.48	0.47
1:D:43:GLU:OE2	1:D:46:ARG:NH1	2.48	0.46
1:A:294:LYS:HE3	1:D:291:LEU:HD12	1.98	0.46
1:A:43:GLU:OE1	1:A:46:ARG:NH1	2.49	0.46
1:B:36:LYS:O	1:B:40:ARG:HG3	2.16	0.46
1:D:352:ASN:HD22	2:D:1000:UFO:HN14	1.64	0.45
1:C:14:LEU:CD1	1:C:311:LEU:HD22	2.46	0.45
1:D:21:ASP:O	1:D:25:VAL:HG23	2.17	0.45
1:C:157:VAL:HB	1:C:158:PRO:CD	2.47	0.45
1:C:331:LYS:NZ	1:C:335:CYS:SG	2.63	0.45
1:A:186:ASP:HA	1:A:191:ASP:HA	1.98	0.45
1:A:355:SER:HB3	1:A:357:GLU:OE1	2.17	0.45
1:D:179:PHE:N	1:D:179:PHE:CD2	2.85	0.45
2:A:1000:UFO:HN1A	1:C:213:LYS:HD2	1.82	0.44
1:A:66:PRO:HG2	1:A:82:TYR:CD2	2.53	0.44
1:A:290:GLN:NE2	1:D:289:ASN:HD21	2.16	0.43
1:D:51:ASP:OD1	1:D:53:ALA:CB	2.66	0.43
1:B:329:ILE:HG13	1:B:384:LEU:HD21	1.99	0.43
1:D:295:LYS:O	1:D:299:MET:HG3	2.18	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:170:ILE:HG21	1:C:224:ILE:HD11	1.99	0.43
1:C:99:MET:HE3	3:C:436:HOH:O	2.18	0.43
1:A:43:GLU:CD	1:A:46:ARG:NH1	2.71	0.43
2:A:1000:UFO:H11	2:A:1000:UFO:H1	1.84	0.43
1:A:102:VAL:HG21	1:A:171:THR:HB	2.00	0.43
1:B:99:MET:CE	1:B:219:SER:HA	2.49	0.43
1:A:244:ARG:NH1	3:A:452:HOH:O	2.52	0.42
1:B:54:ILE:HG22	1:B:58:MET:CE	2.49	0.42
1:C:106:LEU:HB3	1:C:136:PHE:CG	2.54	0.42
1:D:171:THR:O	1:D:172:ASN:HB2	2.20	0.42
1:C:14:LEU:HD12	1:C:311:LEU:HD22	1.99	0.42
1:D:102:VAL:HG21	1:D:171:THR:HB	2.02	0.42
1:A:240:VAL:HG21	1:A:245:GLY:HA2	2.02	0.41
1:C:135:CYS:SG	1:C:179:PHE:CD1	3.14	0.41
1:A:280:LYS:HA	1:A:285:PRO:HA	2.03	0.41
1:D:90:GLU:HG2	1:D:264:ALA:HB2	2.03	0.41
1:B:214:VAL:HG23	1:D:353:GLY:CA	2.50	0.41
1:C:156:LYS:HE3	1:C:160:GLY:O	2.20	0.41
1:B:24:MET:CB	3:B:458:HOH:O	2.69	0.40
1:D:13:LEU:HB3	1:D:16:GLN:HG3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	377/399 (94%)	366 (97%)	11 (3%)	0	100	100
1	B	364/399 (91%)	356 (98%)	8 (2%)	0	100	100
1	C	381/399 (96%)	374 (98%)	6 (2%)	1 (0%)	41	37
1	D	368/399 (92%)	357 (97%)	11 (3%)	0	100	100
All	All	1490/1596 (93%)	1453 (98%)	36 (2%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	141	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	296/314 (94%)	290 (98%)	6 (2%)	55	58
1	B	290/314 (92%)	283 (98%)	7 (2%)	49	51
1	C	295/314 (94%)	289 (98%)	6 (2%)	55	58
1	D	295/314 (94%)	287 (97%)	8 (3%)	44	46
All	All	1176/1256 (94%)	1149 (98%)	27 (2%)	50	53

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	GLU
1	A	100	MET
1	A	149	SER
1	A	166	SER
1	A	277	LEU
1	A	286	LEU
1	B	16	GLN
1	B	21	ASP
1	B	24	MET
1	B	166	SER
1	B	241	LYS
1	B	243	LEU
1	B	277	LEU
1	C	36	LYS
1	C	141	PRO
1	C	188	ASP
1	C	243	LEU
1	C	277	LEU
1	C	286	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	21	ASP
1	D	71	GLN
1	D	100	MET
1	D	291	LEU
1	D	331	LYS
1	D	354	ILE
1	D	372	THR
1	D	392	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	290	GLN
1	B	17	GLN
1	B	290	GLN
1	B	308	GLN
1	B	366	ASN
1	C	251	ASN
1	C	290	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	UFO	A	1000	-	15,15,15	0.86	0	19,21,21	1.61	3 (15%)
2	UFO	B	1001	-	15,15,15	0.90	0	19,21,21	1.00	2 (10%)
2	UFO	D	1000	-	15,15,15	0.75	0	19,21,21	1.42	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UFO	A	1000	-	-	0/2/15/15	0/2/2/2
2	UFO	B	1001	-	-	0/2/15/15	0/2/2/2
2	UFO	D	1000	-	-	0/2/15/15	0/2/2/2

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	UFO	C8-C9-N10	4.22	115.94	110.77
2	A	1000	UFO	C9-C8-N7	3.45	115.00	110.77
2	D	1000	UFO	C8-C9-N10	3.36	114.88	110.77
2	D	1000	UFO	C4-C5-N7	-3.00	118.50	122.03
2	A	1000	UFO	C4-C5-N7	-2.83	118.71	122.03
2	D	1000	UFO	C9-C8-N7	2.38	113.68	110.77
2	B	1001	UFO	C8-C9-N10	-2.33	107.91	110.77
2	B	1001	UFO	C12-N7-C5	2.01	122.63	119.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1000	UFO	3	0
2	D	1000	UFO	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	383/399 (95%)	0.23	12 (3%) 49 48	26, 38, 58, 71	0
1	B	372/399 (93%)	0.14	3 (0%) 86 85	24, 36, 51, 57	0
1	C	385/399 (96%)	0.08	5 (1%) 77 76	25, 36, 52, 64	0
1	D	376/399 (94%)	0.27	18 (4%) 30 29	25, 39, 56, 64	0
All	All	1516/1596 (94%)	0.18	38 (2%) 57 56	24, 37, 55, 71	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	52	ALA	4.1
1	D	50	THR	4.1
1	D	354	ILE	3.7
1	D	282	PHE	3.5
1	A	394	PHE	3.3
1	A	283	GLY	3.3
1	D	169	TRP	3.2
1	D	247	PHE	3.0
1	A	149	SER	3.0
1	D	291	LEU	2.9
1	D	243	LEU	2.9
1	C	282	PHE	2.9
1	D	24	MET	2.8
1	A	377	HIS	2.8
1	A	50	THR	2.7
1	D	212	GLY	2.6
1	C	230	PHE	2.5
1	D	373	TYR	2.5
1	A	159	GLY	2.5
1	D	53	ALA	2.4
1	A	247	PHE	2.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	377	HIS	2.4
1	D	356	ASP	2.4
1	C	145	SER	2.4
1	B	247	PHE	2.3
1	A	147	PRO	2.3
1	D	155	ARG	2.3
1	B	243	LEU	2.2
1	A	148	GLY	2.2
1	C	189	GLY	2.2
1	D	210	ILE	2.2
1	D	118	ALA	2.2
1	C	187	GLU	2.2
1	B	358	PHE	2.2
1	D	51	ASP	2.2
1	A	281	GLN	2.1
1	A	282	PHE	2.1
1	A	379	ILE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	UFO	B	1001	14/14	0.86	0.24	53,54,55,56	0
2	UFO	A	1000	14/14	0.93	0.14	31,33,36,37	0
2	UFO	D	1000	14/14	0.93	0.20	35,41,45,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.