



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 06:59 am BST

PDB ID : 5GQQ
Title : Structure of ALG-2/HEBP2 Complex
Authors : Liu, X.; Ma, J.; Zhang, H.; Feng, Y.
Deposited on : 2016-08-08
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	<div> <div>7%</div> <div> <div></div> <div>77%</div> <div>19%</div> <div>..</div> </div> </div>
1	B	178	<div> <div>8%</div> <div> <div></div> <div>76%</div> <div>20%</div> <div>..</div> </div> </div>
2	C	170	<div> <div>4%</div> <div> <div></div> <div>75%</div> <div>22%</div> <div>..</div> </div> </div>
2	D	170	<div> <div>4%</div> <div> <div></div> <div>79%</div> <div>18%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CL	C	204	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5948 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heme-binding protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	0	0
			1418	903	230	281	4			
1	B	178	Total	C	N	O	S	0	0	0
			1418	903	230	281	4			

- Molecule 2 is a protein called Programmed cell death protein 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	170	Total	C	N	O	S	0	0	0
			1409	894	244	267	4			
2	D	167	Total	C	N	O	S	0	0	0
			1388	880	241	263	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	22	GLY	-	expression tag	UNP O75340
C	23	SER	-	expression tag	UNP O75340
D	22	GLY	-	expression tag	UNP O75340
D	23	SER	-	expression tag	UNP O75340

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	3	Total	Ca	0	0
			3	3		
3	C	3	Total	Ca	0	0
			3	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Cl	0	0
			1	1		
4	C	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	91	Total	O	0	0
			91	91		
5	B	74	Total	O	0	0
			74	74		
5	C	78	Total	O	0	0
			78	78		
5	D	64	Total	O	0	0
			64	64		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heme-binding protein 2



- Molecule 1: Heme-binding protein 2




- Molecule 2: Programmed cell death protein 6



- Molecule 2: Programmed cell death protein 6





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	47.28 Å 78.12 Å 87.17 Å 113.71° 105.64° 93.95°	Depositor
Resolution (Å)	25.93 – 2.20 26.71 – 2.20	Depositor EDS
% Data completeness (in resolution range)	94.9 (25.93-2.20) 95.0 (26.71-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 2.20 Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.194 , 0.225 0.194 , 0.225	Depositor DCC
R_{free} test set	2633 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.011 for -h,-k,h+k+l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5948	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/1455	0.68	1/1976 (0.1%)
1	B	0.40	0/1455	0.60	0/1976
2	C	0.44	0/1441	0.56	0/1946
2	D	0.43	0/1420	0.53	0/1917
All	All	0.44	0/5771	0.59	1/7815 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	179	TYR	N-CA-C	-5.27	96.77	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1418	0	1370	27	0
1	B	1418	0	1370	32	0
2	C	1409	0	1340	27	0
2	D	1388	0	1315	18	0
3	C	3	0	0	0	0
3	D	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	1	0	0	2	0
4	D	1	0	0	1	0
5	A	91	0	0	4	0
5	B	74	0	0	11	0
5	C	78	0	0	2	1
5	D	64	0	0	3	1
All	All	5948	0	5395	97	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:180:ASN:OD1	5:B:201:HOH:O	1.98	0.79
1:B:35:PRO:O	1:B:37:SER:N	2.18	0.77
1:B:115:GLN:NE2	5:B:205:HOH:O	2.18	0.76
1:A:163:ARG:NH2	5:A:203:HOH:O	2.17	0.76
1:B:20:VAL:N	5:B:204:HOH:O	2.17	0.75
1:B:163:ARG:NH1	5:B:206:HOH:O	2.18	0.74
1:B:133:ALA:O	5:B:202:HOH:O	2.06	0.71
2:D:167:ARG:NH1	5:D:302:HOH:O	1.96	0.70
1:B:188:ARG:NH2	5:B:211:HOH:O	2.24	0.69
1:A:98:GLY:HA3	1:A:101:SER:OG	1.93	0.68
1:A:37:SER:O	5:A:201:HOH:O	2.13	0.67
1:B:164:GLU:OE2	5:B:203:HOH:O	2.14	0.65
2:D:89:TRP:NE1	5:D:301:HOH:O	1.85	0.65
2:D:160:ARG:NH1	5:D:303:HOH:O	2.12	0.64
1:B:56:MET:HE1	2:D:34:ARG:HH11	1.62	0.63
2:C:177:GLN:NE2	5:C:305:HOH:O	2.31	0.62
1:A:94:GLU:OE2	5:A:202:HOH:O	2.16	0.62
2:C:188:PHE:HB3	2:D:137:LYS:HG2	1.82	0.62
1:A:32:GLY:N	1:A:33:PRO:HD3	2.17	0.59
1:A:140:ARG:HB2	1:A:158:LEU:HG	1.85	0.59
1:B:83:ILE:HD11	1:B:120:PRO:HG2	1.85	0.59
2:C:137:LYS:HG2	2:D:188:PHE:HB3	1.87	0.55
2:C:135:ILE:HG12	2:C:146:ILE:HD11	1.89	0.55
1:B:51:THR:HG21	1:B:68:LYS:HD2	1.89	0.54
1:A:158:LEU:HD13	1:A:194:LEU:HD12	1.88	0.53
1:A:180:ASN:HA	1:A:184:LYS:HA	1.92	0.52
2:D:139:ASP:OD1	2:D:141:GLN:N	2.36	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:PRO:HB2	1:A:34:GLN:OE1	2.10	0.51
1:A:70:ASN:O	1:A:74:GLN:HG2	2.11	0.51
1:B:140:ARG:HB2	1:B:158:LEU:HG	1.92	0.50
2:C:60:PHE:CE1	2:C:149:ASP:HB3	2.47	0.50
2:C:66:ARG:HH21	2:C:70:SER:HB3	1.76	0.50
1:A:35:PRO:HD2	1:A:37:SER:H	1.77	0.49
2:C:75:GLU:HB2	2:C:77:LYS:HG3	1.94	0.49
2:C:187:VAL:HG21	2:D:187:VAL:HG11	1.94	0.49
2:D:77:LYS:HB2	2:D:77:LYS:HE3	1.58	0.49
1:B:142:PHE:CD1	1:B:150:LYS:HE2	2.48	0.48
2:D:57:TRP:CZ3	2:D:58:THR:HG22	2.48	0.48
1:A:83:ILE:HG23	1:A:115:GLN:HE21	1.78	0.48
1:A:31:ALA:HB1	1:A:32:GLY:C	2.34	0.48
1:A:91:SER:O	1:A:175:TYR:HA	2.14	0.48
1:B:113:SER:O	1:B:116:GLN:HB2	2.14	0.48
1:A:94:GLU:HA	1:A:95:PRO:HD3	1.58	0.48
2:C:48:LEU:HD23	2:C:65:VAL:HG13	1.96	0.47
1:B:180:ASN:HB2	1:B:184:LYS:HE3	1.96	0.47
1:B:94:GLU:O	1:B:103:SER:HB2	2.16	0.46
1:B:80:GLU:HB3	5:B:259:HOH:O	2.16	0.46
1:A:30:ASP:HB3	1:A:31:ALA:H	1.53	0.46
2:C:165:PHE:HB2	2:D:183:TYR:CE2	2.50	0.46
1:A:73:ILE:HB	1:A:83:ILE:HB	1.97	0.46
2:C:73:ASP:OD1	2:C:77:LYS:NZ	2.49	0.46
1:B:153:GLU:OE2	4:C:204:CL:CL	2.71	0.45
2:C:190:ILE:HG22	2:C:191:VAL:H	1.82	0.45
2:D:173:ASP:OD2	2:D:174:GLY:N	2.46	0.45
1:A:34:GLN:HA	1:A:35:PRO:HD3	1.69	0.45
2:C:127:SER:O	2:C:131:HIS:ND1	2.48	0.45
2:C:96:GLN:OE1	2:C:100:ARG:NH2	2.50	0.45
2:C:164:ILE:HG21	2:C:186:MET:HG3	1.98	0.45
1:B:185:LEU:H	1:B:185:LEU:HG	1.66	0.44
2:C:93:THR:O	2:C:96:GLN:HG2	2.17	0.44
1:B:183:VAL:HB	1:B:184:LYS:H	1.63	0.44
1:B:188:ARG:NH2	5:B:223:HOH:O	2.51	0.44
1:B:76:LYS:HE2	1:B:124:GLU:HG3	1.99	0.43
1:A:56:MET:HG3	1:A:102:GLU:HA	2.00	0.43
2:D:38:ASP:OD1	4:D:204:CL:CL	2.73	0.43
1:A:89:VAL:HG22	1:A:109:LEU:HD21	2.01	0.43
1:B:65:GLY:HA3	1:B:107:ILE:HD13	1.99	0.43
2:D:139:ASP:OD1	2:D:140:ARG:N	2.52	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:142:PHE:HD1	1:B:150:LYS:HE2	1.83	0.43
1:B:163:ARG:NH1	5:B:219:HOH:O	2.45	0.43
2:C:75:GLU:OE2	5:C:301:HOH:O	2.21	0.43
1:A:115:GLN:NE2	5:A:213:HOH:O	2.41	0.42
1:A:56:MET:CE	2:C:34:ARG:HD2	2.48	0.42
1:B:162:LEU:HB3	1:B:167:LYS:HB2	2.01	0.42
1:B:183:VAL:O	1:B:184:LYS:HB2	2.18	0.42
2:C:164:ILE:CG2	2:C:186:MET:HG3	2.50	0.42
1:B:164:GLU:CD	5:B:203:HOH:O	2.55	0.42
1:A:181:SER:HA	1:A:182:PRO:HA	1.91	0.42
1:B:56:MET:HE2	1:B:56:MET:HB2	1.78	0.41
2:C:38:ASP:OD1	4:C:204:CL:CL	2.74	0.41
2:C:48:LEU:HG	2:C:52:LEU:HD22	2.02	0.41
2:C:28:LEU:HA	2:C:28:LEU:HD23	1.82	0.41
2:D:80:VAL:HA	2:D:84:GLU:OE1	2.21	0.41
1:A:66:PHE:CZ	1:A:183:VAL:HG11	2.56	0.41
1:B:98:GLY:N	1:B:101:SER:OG	2.40	0.41
1:B:86:THR:O	1:B:185:LEU:HD22	2.21	0.41
2:C:45:ASP:OD2	2:C:66:ARG:NH1	2.53	0.41
1:A:66:PHE:CE1	1:A:183:VAL:HG21	2.55	0.41
2:C:93:THR:HG23	2:C:96:GLN:HE21	1.86	0.41
1:A:56:MET:HE2	2:C:34:ARG:HD2	2.03	0.41
2:D:164:ILE:O	2:D:167:ARG:HB2	2.21	0.41
2:D:173:ASP:CG	2:D:174:GLY:H	2.23	0.41
1:A:107:ILE:HD12	1:A:107:ILE:N	2.35	0.40
1:B:51:THR:HG23	1:B:127:VAL:HG22	2.03	0.40
2:D:134:LEU:HD22	2:D:138:PHE:CE2	2.56	0.40
2:C:116:LYS:NZ	2:C:128:ASP:OD1	2.52	0.40
2:C:188:PHE:HA	2:C:189:SER:HA	1.70	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:378:HOH:O	5:D:364:HOH:O[1_655]	1.89	0.31

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	176/178 (99%)	163 (93%)	7 (4%)	6 (3%)	3	1
1	B	176/178 (99%)	165 (94%)	6 (3%)	5 (3%)	5	2
2	C	168/170 (99%)	155 (92%)	12 (7%)	1 (1%)	25	26
2	D	165/170 (97%)	160 (97%)	5 (3%)	0	100	100
All	All	685/696 (98%)	643 (94%)	30 (4%)	12 (2%)	8	5

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	30	ASP
1	A	33	PRO
1	A	35	PRO
1	B	33	PRO
1	B	36	GLY
1	B	184	LYS
1	B	185	LEU
1	A	180	ASN
1	A	181	SER
2	C	128	ASP
1	A	34	GLN
1	B	35	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/158 (100%)	149 (94%)	9 (6%)	20	24
1	B	158/158 (100%)	148 (94%)	10 (6%)	18	20
2	C	154/154 (100%)	141 (92%)	13 (8%)	11	11
2	D	151/154 (98%)	138 (91%)	13 (9%)	10	10
All	All	621/624 (100%)	576 (93%)	45 (7%)	14	15

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	THR
1	A	53	VAL
1	A	62	ILE
1	A	63	GLN
1	A	97	SER
1	A	149	GLN
1	A	155	LEU
1	A	158	LEU
1	A	180	ASN
1	B	51	THR
1	B	56	MET
1	B	62	ILE
1	B	113	SER
1	B	126	ASP
1	B	134	GLU
1	B	155	LEU
1	B	158	LEU
1	B	183	VAL
1	B	185	LEU
2	C	34	ARG
2	C	52	LEU
2	C	59	PRO
2	C	76	ASN
2	C	77	LYS
2	C	104	ARG
2	C	120	SER
2	C	125	ARG
2	C	128	ASP
2	C	143	ARG
2	C	158	LEU
2	C	178	VAL
2	C	184	LEU
2	D	45	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	58	THR
2	D	125	ARG
2	D	128	ASP
2	D	135	ILE
2	D	136	ARG
2	D	153	GLN
2	D	158	LEU
2	D	159	GLN
2	D	172	GLN
2	D	177	GLN
2	D	178	VAL
2	D	188	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	115	GLN
1	A	149	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	178/178 (100%)	0.11	12 (6%) 17 16	16, 31, 85, 122	0
1	B	178/178 (100%)	0.23	15 (8%) 11 9	17, 40, 92, 115	0
2	C	170/170 (100%)	-0.14	6 (3%) 44 42	13, 37, 82, 116	0
2	D	167/170 (98%)	-0.14	6 (3%) 42 41	14, 40, 90, 128	0
All	All	693/696 (99%)	0.02	39 (5%) 24 23	13, 36, 89, 128	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	31	ALA	6.1
1	B	32	GLY	5.6
1	A	33	PRO	5.5
2	D	143	ARG	4.4
1	B	117	PHE	4.2
1	B	33	PRO	4.1
1	A	32	GLY	4.1
1	B	186	LEU	4.0
1	B	29	GLU	3.8
1	A	31	ALA	3.7
1	B	185	LEU	3.4
1	B	109	LEU	3.4
1	B	89	VAL	3.1
1	B	30	ASP	3.1
1	A	29	GLU	2.9
2	C	128	ASP	2.9
2	D	129	GLN	2.9
1	B	35	PRO	2.8
1	A	35	PRO	2.8
2	C	129	GLN	2.7
2	C	116	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	30	ASP	2.6
1	B	108	SER	2.5
2	D	136	ARG	2.5
2	C	138	PHE	2.5
1	A	89	VAL	2.5
2	C	172	GLN	2.5
1	A	177	ALA	2.4
2	D	172	GLN	2.4
2	D	125	ARG	2.4
2	C	143	ARG	2.3
1	A	145	PHE	2.2
1	A	36	GLY	2.1
1	B	90	THR	2.1
1	B	123	LEU	2.1
2	D	167	ARG	2.1
1	A	186	LEU	2.1
1	A	192	VAL	2.1
1	B	91	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	D	203	1/1	0.59	0.14	107,107,107,107	0
3	CA	C	203	1/1	0.87	0.16	114,114,114,114	0
3	CA	D	202	1/1	0.94	0.06	47,47,47,47	0
4	CL	C	204	1/1	0.97	0.07	41,41,41,41	0
3	CA	C	202	1/1	0.98	0.07	40,40,40,40	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CL	D	204	1/1	0.99	0.05	46,46,46,46	0
3	CA	C	201	1/1	1.00	0.07	16,16,16,16	0
3	CA	D	201	1/1	1.00	0.08	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.