



Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 05:23 pm BST

PDB ID : 1GRF
Title : SUBSTRATE BINDING AND CATALYSIS BY GLUTATHIONE REDUCTASE AS DERIVED FROM REFINED ENZYME: SUBSTRATE CRYSTAL STRUCTURES AT 2 ANGSTROMS RESOLUTION
Authors : Karplus, P.A.; Schulz, G.E.
Deposited on : 1992-12-15
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACM	A	481	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4092 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTATHIONE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	461	3499	2212	603	660	24	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is ACETAMIDE (three-letter code: ACM) (formula: C₂H₅NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			4	2	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	531	Total 531	O 531	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GLUTATHIONE REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	B 1 1 2	Depositor
Cell constants a, b, c, α , β , γ	119.80 Å 84.50 Å 63.20 Å 90.00° 90.00° 58.70°	Depositor
Resolution (Å)	10.00 – 2.00 72.20 – 1.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.00) 96.4 (72.20-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.164 , (Not available) 0.154 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	20.0	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 100.1	EDS
L-test for twinning ¹	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4092	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, FAD, ACM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	18/3566 (0.5%)	1.26	40/4824 (0.8%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	124	GLU	CD-OE1	7.10	1.33	1.25
1	A	358	GLU	CD-OE2	6.90	1.33	1.25
1	A	185	GLU	CD-OE1	6.75	1.33	1.25
1	A	394	GLU	CD-OE1	6.59	1.32	1.25
1	A	427	GLU	CD-OE1	6.54	1.32	1.25
1	A	91	GLU	CD-OE2	6.52	1.32	1.25
1	A	77	GLU	CD-OE1	6.28	1.32	1.25
1	A	317	GLU	CD-OE1	6.24	1.32	1.25
1	A	236	GLU	CD-OE1	6.07	1.32	1.25
1	A	384	GLU	CD-OE1	6.06	1.32	1.25
1	A	262	GLU	CD-OE1	5.98	1.32	1.25
1	A	253	GLU	CD-OE2	5.88	1.32	1.25
1	A	473	GLU	CD-OE2	5.74	1.31	1.25
1	A	101	GLU	CD-OE2	5.68	1.31	1.25
1	A	428	GLU	CD-OE1	5.65	1.31	1.25
1	A	386	GLU	CD-OE1	5.58	1.31	1.25
1	A	239	GLU	CD-OE1	5.31	1.31	1.25
1	A	41	GLU	CD-OE2	5.02	1.31	1.25

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	91	GLU	C-N-CA	-10.11	101.06	122.30
1	A	159	MET	CB-CA-C	-8.34	93.72	110.40
1	A	461	ASP	CB-CG-OD2	-8.30	110.83	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	291	ARG	NE-CZ-NH1	-8.02	116.29	120.30
1	A	283	ASP	CB-CG-OD1	-7.67	111.39	118.30
1	A	104	ASP	CB-CG-OD1	7.62	125.16	118.30
1	A	104	ASP	CB-CG-OD2	-7.50	111.55	118.30
1	A	309	ASP	CB-CG-OD2	-7.31	111.72	118.30
1	A	359	ASP	CB-CG-OD2	-7.25	111.77	118.30
1	A	308	ASP	CB-CG-OD2	-7.16	111.86	118.30
1	A	352	ARG	NE-CZ-NH2	7.02	123.81	120.30
1	A	297	ASP	CB-CG-OD2	-6.97	112.03	118.30
1	A	84	ASP	CB-CG-OD1	-6.94	112.06	118.30
1	A	308	ASP	CB-CG-OD1	6.88	124.49	118.30
1	A	22	ASP	CB-CG-OD1	-6.64	112.32	118.30
1	A	97	ARG	NE-CZ-NH2	6.50	123.55	120.30
1	A	359	ASP	CB-CG-OD1	6.39	124.05	118.30
1	A	461	ASP	CB-CG-OD1	6.30	123.97	118.30
1	A	147	TYR	CB-CG-CD2	-6.28	117.23	121.00
1	A	178	ASP	CB-CG-OD1	6.27	123.94	118.30
1	A	178	ASP	CB-CG-OD2	-6.22	112.70	118.30
1	A	283	ASP	CB-CG-OD2	6.15	123.83	118.30
1	A	441	ASP	CB-CG-OD1	6.12	123.80	118.30
1	A	385	ASP	CB-CG-OD2	5.88	123.59	118.30
1	A	84	ASP	CB-CG-OD2	5.75	123.48	118.30
1	A	135	ASP	CB-CG-OD1	-5.70	113.17	118.30
1	A	363	ASP	CB-CG-OD1	5.67	123.41	118.30
1	A	363	ASP	CB-CG-OD2	-5.65	113.22	118.30
1	A	385	ASP	CB-CG-OD1	-5.53	113.33	118.30
1	A	291	ARG	NE-CZ-NH2	5.49	123.04	120.30
1	A	459	ASP	CB-CG-OD2	-5.46	113.39	118.30
1	A	227	ASP	CB-CG-OD1	5.40	123.16	118.30
1	A	22	ASP	CB-CG-OD2	5.34	123.11	118.30
1	A	297	ASP	CB-CG-OD1	5.29	123.06	118.30
1	A	91	GLU	CA-C-N	5.24	126.68	116.20
1	A	220	ASP	CB-CG-OD1	-5.23	113.59	118.30
1	A	281	ASP	CB-CG-OD1	-5.22	113.60	118.30
1	A	220	ASP	CB-CG-OD2	5.15	122.94	118.30
1	A	309	ASP	CB-CG-OD1	5.12	122.91	118.30
1	A	97	ARG	N-CA-CB	5.04	119.67	110.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3499	0	3538	59	1
2	A	5	0	0	0	0
3	A	53	0	31	0	0
4	A	4	0	3	1	0
5	A	531	0	0	16	2
All	All	4092	0	3572	59	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:293:PRO:HG3	1:A:312:HIS:CD2	2.16	0.80
1:A:296:LYS:HB2	5:A:584:HOH:O	1.83	0.78
1:A:185:GLU:HG3	5:A:807:HOH:O	1.87	0.73
1:A:93:LYS:HG2	1:A:93:LYS:O	1.91	0.70
1:A:77:GLU:OE2	1:A:374:HIS:ND1	2.25	0.69
1:A:224:ARG:HD2	5:A:985:HOH:O	1.94	0.68
1:A:127:ARG:HD2	5:A:724:HOH:O	1.97	0.64
1:A:266:VAL:HG22	1:A:276:MET:HG2	1.81	0.62
1:A:373:SER:HB2	5:A:992:HOH:O	2.00	0.61
1:A:110:LEU:O	1:A:114:TYR:HD1	1.85	0.59
1:A:114:TYR:OH	4:A:481:ACM:N	2.32	0.58
1:A:125:ILE:HG22	1:A:127:ARG:HD3	1.84	0.57
1:A:315:VAL:HA	1:A:320:ASN:O	2.05	0.56
1:A:164:HIS:HB2	5:A:925:HOH:O	2.06	0.56
1:A:218:ARG:HG3	5:A:595:HOH:O	2.07	0.55
1:A:292:VAL:HG23	1:A:293:PRO:HD2	1.89	0.54
1:A:352:ARG:HB2	1:A:360:SER:HB3	1.90	0.54
1:A:403:PHE:CE1	1:A:473:GLU:HG3	2.43	0.53
1:A:421:MET:SD	1:A:457:LYS:HD3	2.47	0.53
1:A:202:MET:CE	1:A:202:MET:HA	2.39	0.52
1:A:115:GLN:O	1:A:119:THR:HG23	2.09	0.52
1:A:331:ASP:HA	1:A:337:LEU:HD22	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:324:LYS:HE2	5:A:752:HOH:O	2.10	0.52
1:A:159:MET:HB3	1:A:160:PRO:HD2	1.91	0.51
1:A:159:MET:HB3	1:A:160:PRO:CD	2.41	0.50
1:A:45:ARG:NH2	1:A:124:GLU:OE1	2.44	0.50
1:A:411:THR:O	1:A:414:LYS:HE2	2.11	0.50
1:A:339:THR:HB	1:A:340:PRO:HD3	1.93	0.50
1:A:192:ILE:HD13	1:A:203:ALA:HA	1.95	0.49
1:A:426:LYS:HE3	1:A:426:LYS:HB3	1.60	0.48
1:A:371:VAL:HB	1:A:377:ILE:HB	1.95	0.47
1:A:236:GLU:HG2	5:A:929:HOH:O	2.15	0.47
1:A:125:ILE:CG2	1:A:127:ARG:HD3	2.45	0.46
1:A:250:GLN:NE2	5:A:919:HOH:O	2.50	0.45
1:A:198:ILE:O	1:A:202:MET:HG2	2.17	0.45
1:A:45:ARG:HH22	1:A:124:GLU:HB2	1.81	0.45
1:A:388:ILE:HG23	1:A:393:ILE:HG12	1.99	0.44
1:A:307:THR:HA	1:A:312:HIS:O	2.17	0.44
1:A:78:PHE:HB3	5:A:973:HOH:O	2.16	0.44
1:A:100:LYS:O	1:A:100:LYS:HG2	2.17	0.43
1:A:474:LEU:HA	1:A:474:LEU:HD23	1.89	0.43
1:A:65:PRO:HB2	1:A:181:PHE:CE1	2.53	0.43
1:A:91:GLU:O	1:A:91:GLU:HG3	2.16	0.43
1:A:465:ALA:HB1	1:A:472:GLU:HB2	2.00	0.43
1:A:454:GLY:HA2	5:A:884:HOH:O	2.19	0.42
1:A:49:VAL:HG11	1:A:130:ALA:HB2	2.01	0.42
1:A:138:PRO:HA	5:A:582:HOH:O	2.19	0.42
1:A:252:LYS:HG3	5:A:975:HOH:O	2.20	0.42
1:A:467:HIS:HA	1:A:468:PRO:HA	1.87	0.41
1:A:357:LYS:NZ	5:A:791:HOH:O	2.53	0.41
1:A:138:PRO:HB3	1:A:326:ILE:HD11	2.03	0.41
1:A:333:CYS:HB2	5:A:525:HOH:O	2.19	0.41
1:A:400:SER:HA	1:A:419:MET:O	2.20	0.41
1:A:418:VAL:O	1:A:435:MET:HA	2.21	0.41
1:A:118:LEU:HD23	1:A:118:LEU:HA	1.88	0.41
1:A:135:ASP:HA	1:A:136:PRO:HD3	1.94	0.41
1:A:257:THR:HG21	1:A:262:GLU:HB2	2.03	0.41
1:A:202:MET:HE3	1:A:202:MET:HA	2.02	0.40
1:A:218:ARG:HE	1:A:218:ARG:HB3	1.73	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1011:HOH:O	5:A:1012:HOH:O[4_575]	0.45	1.75
5:A:580:HOH:O	5:A:580:HOH:O[2_665]	0.62	1.58
1:A:90:CYS:SG	1:A:90:CYS:SG[2_665]	1.56	0.64

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	459/478 (96%)	442 (96%)	16 (4%)	1 (0%)	47 44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	136	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	382/393 (97%)	361 (94%)	21 (6%)	21 17

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	HIS
1	A	89	SER
1	A	90	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	91	GLU
1	A	93	LYS
1	A	127	ARG
1	A	134	SER
1	A	167	GLN
1	A	185	GLU
1	A	212	LYS
1	A	252	LYS
1	A	253	GLU
1	A	275	VAL
1	A	276	MET
1	A	294	ASN
1	A	296	LYS
1	A	317	GLU
1	A	397	LYS
1	A	426	LYS
1	A	468	PRO
1	A	478	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	HIS
1	A	117	ASN
1	A	129	HIS
1	A	182	GLN
1	A	301	ASN
1	A	366	ASN
1	A	425	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FAD	A	479	-	51,58,58	1.68	8 (15%)	60,89,89	2.19	15 (25%)
4	ACM	A	481	1	3,3,3	3.55	2 (66%)	3,3,3	2.44	1 (33%)
2	PO4	A	480	-	4,4,4	1.33	0	6,6,6	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	479	-	-	3/30/50/50	0/6/6/6

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	479	FAD	C4X-C10	7.50	1.46	1.38
4	A	481	ACM	O-C1	-4.98	1.13	1.24
3	A	479	FAD	C4-C4X	3.97	1.48	1.41
4	A	481	ACM	C1-N	3.17	1.44	1.32
3	A	479	FAD	O4B-C1B	3.13	1.45	1.41
3	A	479	FAD	C4-N3	2.85	1.38	1.33
3	A	479	FAD	C5'-C4'	2.69	1.55	1.51
3	A	479	FAD	C10-N1	-2.60	1.29	1.33
3	A	479	FAD	C2B-C1B	-2.24	1.50	1.53
3	A	479	FAD	P-O2P	-2.17	1.45	1.55

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	479	FAD	C4-N3-C2	9.83	123.44	115.14
3	A	479	FAD	C4X-C4-N3	-5.85	115.42	123.43
3	A	479	FAD	C10-C4X-N5	4.96	124.69	121.26
3	A	479	FAD	C4-C4X-C10	-4.56	116.93	119.95
4	A	481	ACM	O-C1-C2	3.54	127.86	120.63
3	A	479	FAD	C4X-C10-N10	-3.43	116.78	120.30
3	A	479	FAD	C4A-C5A-N7A	2.97	112.49	109.40
3	A	479	FAD	O4B-C1B-C2B	-2.93	102.65	106.93
3	A	479	FAD	C5'-C4'-C3'	-2.90	106.60	112.20
3	A	479	FAD	O2A-PA-O1A	2.55	124.87	112.24
3	A	479	FAD	C1'-N10-C9A	2.46	120.23	118.29
3	A	479	FAD	O2P-P-O1P	2.35	123.87	112.24
3	A	479	FAD	C2A-N1A-C6A	2.32	122.72	118.75
3	A	479	FAD	O2B-C2B-C3B	-2.27	104.47	111.82
3	A	479	FAD	C5A-C6A-N1A	-2.17	115.42	120.35
3	A	479	FAD	O3B-C3B-C4B	2.17	117.33	111.05

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	479	FAD	O4B-C4B-C5B-O5B
3	A	479	FAD	C3B-C4B-C5B-O5B
3	A	479	FAD	P-O3P-PA-O2A


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	481	ACM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	461/478 (96%)	-0.92	0 100 100	8, 19, 43, 67	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACM	A	481	4/4	0.95	0.14	23,23,35,36	0
2	PO4	A	480	5/5	0.96	0.18	31,32,37,42	5
3	FAD	A	479	53/53	0.98	0.07	5,12,21,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.