



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2020 – 02:11 AM BST

PDB ID : 5GTW
Title : The N253R mutant structures of trehalose synthase from *Deinococcus radiodurans* display two different active-site conformations
Authors : Chow, S.Y.; Wei, Y.J.; Liaw, S.H.
Deposited on : 2016-08-23
Resolution : 2.93 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2969 (2.98-2.90)
Clashscore	141614	3218 (2.98-2.90)
Ramachandran outliers	138981	3122 (2.98-2.90)
Sidechain outliers	138945	3124 (2.98-2.90)
RSRZ outliers	127900	2902 (2.98-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	571	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>15%</div> <div>• •</div> </div> </div>
1	B	571	<div> <div>2%</div> <div> <div></div> <div>78%</div> <div>17%</div> <div>• •</div> </div> </div>
1	C	571	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>• •</div> </div> </div>
1	D	571	<div> <div>4%</div> <div> <div></div> <div>78%</div> <div>16%</div> <div>• •</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 17815 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Trehalose synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	548	Total	C	N	O	S	0	0	0
			4407	2820	753	818	16			
1	B	548	Total	C	N	O	S	0	0	0
			4407	2820	753	818	16			
1	C	548	Total	C	N	O	S	0	0	0
			4407	2820	753	818	16			
1	D	548	Total	C	N	O	S	0	0	0
			4407	2820	753	818	16			

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP I3NX86
A	0	VAL	-	expression tag	UNP I3NX86
A	1	PRO	-	expression tag	UNP I3NX86
A	97	TRP	ARG	engineered mutation	UNP I3NX86
A	253	ARG	ASN	engineered mutation	UNP I3NX86
A	313	ILE	THR	engineered mutation	UNP I3NX86
A	380	VAL	ILE	engineered mutation	UNP I3NX86
A	553	SER	-	expression tag	UNP I3NX86
A	554	ARG	-	expression tag	UNP I3NX86
A	555	VAL	-	expression tag	UNP I3NX86
A	556	ASP	-	expression tag	UNP I3NX86
A	557	LYS	-	expression tag	UNP I3NX86
A	558	LEU	-	expression tag	UNP I3NX86
A	559	ALA	-	expression tag	UNP I3NX86
A	560	ALA	-	expression tag	UNP I3NX86
A	561	ALA	-	expression tag	UNP I3NX86
A	562	LEU	-	expression tag	UNP I3NX86
A	563	GLU	-	expression tag	UNP I3NX86
A	564	HIS	-	expression tag	UNP I3NX86
A	565	HIS	-	expression tag	UNP I3NX86
A	566	HIS	-	expression tag	UNP I3NX86

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	567	HIS	-	expression tag	UNP I3NX86
A	568	HIS	-	expression tag	UNP I3NX86
A	569	HIS	-	expression tag	UNP I3NX86
B	-1	MET	-	expression tag	UNP I3NX86
B	0	VAL	-	expression tag	UNP I3NX86
B	1	PRO	-	expression tag	UNP I3NX86
B	97	TRP	ARG	engineered mutation	UNP I3NX86
B	253	ARG	ASN	engineered mutation	UNP I3NX86
B	313	ILE	THR	engineered mutation	UNP I3NX86
B	380	VAL	ILE	engineered mutation	UNP I3NX86
B	553	SER	-	expression tag	UNP I3NX86
B	554	ARG	-	expression tag	UNP I3NX86
B	555	VAL	-	expression tag	UNP I3NX86
B	556	ASP	-	expression tag	UNP I3NX86
B	557	LYS	-	expression tag	UNP I3NX86
B	558	LEU	-	expression tag	UNP I3NX86
B	559	ALA	-	expression tag	UNP I3NX86
B	560	ALA	-	expression tag	UNP I3NX86
B	561	ALA	-	expression tag	UNP I3NX86
B	562	LEU	-	expression tag	UNP I3NX86
B	563	GLU	-	expression tag	UNP I3NX86
B	564	HIS	-	expression tag	UNP I3NX86
B	565	HIS	-	expression tag	UNP I3NX86
B	566	HIS	-	expression tag	UNP I3NX86
B	567	HIS	-	expression tag	UNP I3NX86
B	568	HIS	-	expression tag	UNP I3NX86
B	569	HIS	-	expression tag	UNP I3NX86
C	-1	MET	-	expression tag	UNP I3NX86
C	0	VAL	-	expression tag	UNP I3NX86
C	1	PRO	-	expression tag	UNP I3NX86
C	97	TRP	ARG	engineered mutation	UNP I3NX86
C	253	ARG	ASN	engineered mutation	UNP I3NX86
C	313	ILE	THR	engineered mutation	UNP I3NX86
C	380	VAL	ILE	engineered mutation	UNP I3NX86
C	553	SER	-	expression tag	UNP I3NX86
C	554	ARG	-	expression tag	UNP I3NX86
C	555	VAL	-	expression tag	UNP I3NX86
C	556	ASP	-	expression tag	UNP I3NX86
C	557	LYS	-	expression tag	UNP I3NX86
C	558	LEU	-	expression tag	UNP I3NX86
C	559	ALA	-	expression tag	UNP I3NX86
C	560	ALA	-	expression tag	UNP I3NX86

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	561	ALA	-	expression tag	UNP I3NX86
C	562	LEU	-	expression tag	UNP I3NX86
C	563	GLU	-	expression tag	UNP I3NX86
C	564	HIS	-	expression tag	UNP I3NX86
C	565	HIS	-	expression tag	UNP I3NX86
C	566	HIS	-	expression tag	UNP I3NX86
C	567	HIS	-	expression tag	UNP I3NX86
C	568	HIS	-	expression tag	UNP I3NX86
C	569	HIS	-	expression tag	UNP I3NX86
D	-1	MET	-	expression tag	UNP I3NX86
D	0	VAL	-	expression tag	UNP I3NX86
D	1	PRO	-	expression tag	UNP I3NX86
D	97	TRP	ARG	engineered mutation	UNP I3NX86
D	253	ARG	ASN	engineered mutation	UNP I3NX86
D	313	ILE	THR	engineered mutation	UNP I3NX86
D	380	VAL	ILE	engineered mutation	UNP I3NX86
D	553	SER	-	expression tag	UNP I3NX86
D	554	ARG	-	expression tag	UNP I3NX86
D	555	VAL	-	expression tag	UNP I3NX86
D	556	ASP	-	expression tag	UNP I3NX86
D	557	LYS	-	expression tag	UNP I3NX86
D	558	LEU	-	expression tag	UNP I3NX86
D	559	ALA	-	expression tag	UNP I3NX86
D	560	ALA	-	expression tag	UNP I3NX86
D	561	ALA	-	expression tag	UNP I3NX86
D	562	LEU	-	expression tag	UNP I3NX86
D	563	GLU	-	expression tag	UNP I3NX86
D	564	HIS	-	expression tag	UNP I3NX86
D	565	HIS	-	expression tag	UNP I3NX86
D	566	HIS	-	expression tag	UNP I3NX86
D	567	HIS	-	expression tag	UNP I3NX86
D	568	HIS	-	expression tag	UNP I3NX86
D	569	HIS	-	expression tag	UNP I3NX86

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0
2	D	1	Total Ca 1 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total	Ca	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	1	3		
4	B	1	Total	C	N	O	0	0
			8	4	1	3		
4	C	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	43	Total 43	O 43	0	0
5	B	33	Total 33	O 33	0	0
5	C	29	Total 29	O 29	0	0
5	D	50	Total 50	O 50	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Trehalose synthase




• Molecule 1: Trehalose synthase





• Molecule 1: Trehalose synthase





● Molecule 1: Trehalose synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	98.38Å 134.06Å 197.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.93 19.93 – 2.93	Depositor EDS
% Data completeness (in resolution range)	95.2 (20.00-2.93) 95.6 (19.93-2.93)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.81 (at 2.93Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.182 , 0.267 0.186 , 0.265	Depositor DCC
R_{free} test set	2696 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	52.8	Xtriage
Anisotropy	0.627	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 65.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	17815	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, MG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/4540	0.83	2/6185 (0.0%)
1	B	0.65	0/4540	0.79	0/6185
1	C	0.67	0/4540	0.80	1/6185 (0.0%)
1	D	0.67	0/4540	0.83	1/6185 (0.0%)
All	All	0.67	0/18160	0.81	4/24740 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	89	ARG	NE-CZ-NH1	6.91	123.75	120.30
1	C	72	ARG	NE-CZ-NH1	6.57	123.58	120.30
1	A	352	ARG	NE-CZ-NH1	5.20	122.90	120.30
1	A	331	ARG	NE-CZ-NH2	-5.01	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4407	0	4208	43	0
1	B	4407	0	4208	59	0
1	C	4407	0	4208	40	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	4407	0	4208	49	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	8	0	12	0	0
4	B	8	0	12	2	0
4	C	8	0	12	1	0
5	A	43	0	0	2	0
5	B	33	0	0	1	0
5	C	29	0	0	0	0
5	D	50	0	0	0	0
All	All	17815	0	16868	189	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 189 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:THR:HG1	1:A:171:HIS:HE2	1.25	0.81
1:B:182:TYR:OH	1:B:191:LEU:HD13	1.81	0.78
1:C:366:GLU:O	1:C:370:THR:HG23	1.86	0.75
1:C:182:TYR:OH	1:C:191:LEU:HD22	1.88	0.74
1:B:236:ARG:NH1	1:B:270:GLU:O	2.24	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	546/571 (96%)	514 (94%)	32 (6%)	0	100	100
1	B	546/571 (96%)	509 (93%)	34 (6%)	3 (0%)	29	60
1	C	546/571 (96%)	498 (91%)	45 (8%)	3 (0%)	29	60
1	D	546/571 (96%)	510 (93%)	33 (6%)	3 (0%)	29	60
All	All	2184/2284 (96%)	2031 (93%)	144 (7%)	9 (0%)	34	64

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	304	LYS
1	D	214	LEU
1	D	252	ALA
1	B	70	ASP
1	C	77	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	465/484 (96%)	440 (95%)	25 (5%)	22	51
1	B	465/484 (96%)	441 (95%)	24 (5%)	23	53
1	C	465/484 (96%)	445 (96%)	20 (4%)	29	60
1	D	465/484 (96%)	424 (91%)	41 (9%)	10	28
All	All	1860/1936 (96%)	1750 (94%)	110 (6%)	19	47

5 of 110 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	11	SER
1	C	381	LEU
1	D	397	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	52	TRP
1	C	147	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 8 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TRS	B	603	-	7,7,7	0.34	0	9,9,9	0.74	0
4	TRS	C	603	-	7,7,7	0.34	0	9,9,9	1.00	1 (11%)
4	TRS	A	603	-	7,7,7	0.55	0	9,9,9	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TRS	B	603	-	-	3/9/9/9	-
4	TRS	C	603	-	-	6/9/9/9	-
4	TRS	A	603	-	-	6/9/9/9	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	603	TRS	C3-C-N	2.26	114.72	107.98

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	603	TRS	C2-C-C3-O3
4	C	603	TRS	C2-C-C1-O1
4	C	603	TRS	C3-C-C1-O1
4	C	603	TRS	N-C-C1-O1
4	C	603	TRS	C1-C-C3-O3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	603	TRS	2	0
4	C	603	TRS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	548/571 (95%)	-0.29	14 (2%) 56 56	21, 51, 80, 115	3 (0%)
1	B	548/571 (95%)	-0.27	10 (1%) 68 69	24, 52, 80, 98	3 (0%)
1	C	548/571 (95%)	-0.19	18 (3%) 46 45	23, 55, 82, 126	3 (0%)
1	D	548/571 (95%)	-0.18	22 (4%) 38 37	25, 52, 90, 130	3 (0%)
All	All	2192/2284 (95%)	-0.23	64 (2%) 51 51	21, 53, 82, 130	12 (0%)

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	146	ASP	6.9
1	C	146	ASP	4.0
1	D	153	ASP	4.0
1	D	152	THR	3.8
1	D	156	VAL	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	TRS	A	603	8/8	0.86	0.28	63,75,83,83	0
4	TRS	B	603	8/8	0.92	0.19	64,79,83,88	0
4	TRS	C	603	8/8	0.94	0.21	57,62,72,80	0
3	MG	B	602	1/1	0.95	0.47	63,63,63,63	0
2	CA	B	601	1/1	0.95	0.21	78,78,78,78	0
3	MG	C	602	1/1	0.96	0.07	40,40,40,40	0
3	MG	D	601	1/1	0.97	0.09	35,35,35,35	0
3	MG	A	602	1/1	0.97	0.06	29,29,29,29	0
2	CA	C	601	1/1	0.98	0.10	59,59,59,59	0
2	CA	A	601	1/1	0.99	0.19	57,57,57,57	0
2	CA	D	600	1/1	0.99	0.18	56,56,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.