



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 10:59 am BST

PDB ID : 3GUU  
Title : X-ray structure of Candida Antarctica lipase A  
Authors : Brandt, A.-M.; Li, X.-G.; Nymalm-Rejstrom, Y.; Airenne, T.; Kanerva, L.T.;  
Salminen, T.A.  
Deposited on : 2009-03-30  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	462	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 81%, yellow 11%, orange 7%, grey 3%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>81%</span> <span>11%</span> <span>7%</span> </div> </div>
1	B	462	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 85%, yellow 8%, orange 7%, grey 3%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>85%</span> <span>8%</span> <span>7%</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	443	-	-	X	-

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7077 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lipase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	431	Total	C	N	O	S	0	1	0
			3272	2107	536	623	6			
1	B	431	Total	C	N	O	S	0	0	0
			3263	2102	535	620	6			

- Molecule 2 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula:  $C_{10}H_{22}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			16	10	6		
2	B	1	Total	C	O	0	0
			16	10	6		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	249	Total 249	O 249	0	0
5	B	250	Total 250	O 250	0	0



- Molecule 1: Lipase A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.10 Å 92.10 Å 300.40 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.92 – 2.10 19.92 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.92-2.10) 99.0 (19.92-2.10)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.66 (at 2.09 Å)	Xtriage
Refinement program	REFMAC 5.3.0040	Depositor
R, $R_{free}$	0.189 , 0.234 0.196 , 0.234	Depositor DCC
$R_{free}$ test set	3784 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.7	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 54.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	7077	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.96	2/3363 (0.1%)	0.84	7/4605 (0.2%)
1	B	1.01	2/3354 (0.1%)	0.83	5/4593 (0.1%)
All	All	0.99	4/6717 (0.1%)	0.84	12/9198 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	332	ILE	C-O	-5.54	1.12	1.23
1	B	320	SER	C-N	-5.25	1.22	1.34
1	B	22	TYR	CD1-CE1	5.17	1.47	1.39
1	A	246	PRO	C-N	-5.02	1.22	1.34

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	326	ARG	NE-CZ-NH2	-9.85	115.38	120.30
1	A	367	LEU	CA-CB-CG	-9.83	92.70	115.30
1	B	367	LEU	CA-CB-CG	-9.63	93.14	115.30
1	B	60	ARG	NE-CZ-NH2	-9.36	115.62	120.30
1	B	60	ARG	NE-CZ-NH1	9.04	124.82	120.30
1	A	326	ARG	NE-CZ-NH1	8.21	124.41	120.30
1	A	247	ASP	O-C-N	-6.42	112.42	122.70
1	B	367	LEU	CB-CG-CD1	6.37	121.83	111.00
1	A	95	ASP	CB-CG-OD1	6.29	123.97	118.30
1	B	60	ARG	CD-NE-CZ	5.76	131.66	123.60
1	A	247	ASP	CB-CG-OD2	5.23	123.01	118.30
1	A	367	LEU	CB-CG-CD1	5.08	119.64	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3272	0	3209	38	0
1	B	3263	0	3204	28	0
2	A	16	0	22	1	0
2	B	16	0	22	3	0
3	A	6	0	8	8	0
4	B	5	0	0	0	0
5	A	249	0	0	4	0
5	B	250	0	0	6	0
All	All	7077	0	6465	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:127:ILE:HG13	5:B:631:HOH:O	1.33	1.25
1:A:247:ASP:HB2	3:A:443:GOL:O1	1.67	0.92
1:A:304:ILE:HD11	3:A:443:GOL:H11	1.51	0.91
1:A:245:HIS:HD1	3:A:443:GOL:HO3	1.20	0.85
1:B:238:VAL:HG22	2:B:442:1PE:H222	1.59	0.83
1:A:198:GLU:HG3	5:A:507:HOH:O	1.78	0.83
1:B:127:ILE:CD1	5:B:631:HOH:O	2.24	0.80
1:A:245:HIS:ND1	3:A:443:GOL:O3	2.16	0.71
1:B:330:HIS:HD2	5:B:447:HOH:O	1.73	0.70
1:A:247:ASP:HB2	3:A:443:GOL:C1	2.22	0.69
1:B:127:ILE:HG23	1:B:137:VAL:HG11	1.75	0.69
1:B:367:LEU:HD22	1:B:429:SER:CB	2.23	0.68
1:A:365:GLU:OE1	5:A:453:HOH:O	2.13	0.67
1:B:238:VAL:HG22	2:B:442:1PE:C22	2.25	0.66
1:A:83:SER:HA	1:A:84:PRO:C	2.20	0.62
1:B:33:LYS:HA	1:B:60:ARG:HD2	1.83	0.60
1:B:65:GLN:OE1	5:B:673:HOH:O	2.15	0.59
1:B:127:ILE:CG1	5:B:631:HOH:O	2.05	0.57
1:B:281:TYR:N	1:B:282:PRO:HD3	2.20	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:116:LYS:HE2	5:A:602:HOH:O	2.04	0.56
1:B:259:LYS:HE2	1:B:289:LEU:HD23	1.88	0.56
1:A:155:GLU:OE1	1:A:188:HIS:ND1	2.39	0.55
1:A:210:SER:HB2	1:A:381:ILE:HD11	1.88	0.55
1:A:304:ILE:HD11	3:A:443:GOL:C1	2.32	0.54
1:A:367:LEU:HD13	1:A:433:LYS:HB2	1.89	0.54
1:A:159:ILE:HD11	1:A:189:ALA:HB1	1.88	0.54
1:B:367:LEU:HD13	1:B:433:LYS:HB2	1.89	0.53
1:B:222:PHE:HB2	2:B:442:1PE:H131	1.91	0.53
1:A:222:PHE:HB2	2:A:442:1PE:H231	1.92	0.52
1:A:242:SER:HB3	1:A:249:GLU:OE1	2.12	0.50
1:B:159:ILE:HD11	1:B:189:ALA:HB1	1.94	0.49
1:B:300:PRO:O	1:B:304:ILE:HG12	2.14	0.48
1:A:149:PHE:O	1:A:150:ILE:HG22	2.14	0.47
1:A:235:LEU:HD21	1:A:263:THR:HG22	1.96	0.47
1:A:245:HIS:HB3	3:A:443:GOL:H2	1.97	0.47
1:B:281:TYR:N	1:B:282:PRO:CD	2.77	0.47
1:A:292:ASP:OD2	1:A:298:GLU:OE1	2.33	0.47
1:A:428:GLN:NE2	1:A:432:GLY:C	2.68	0.46
1:A:181:GLU:HA	1:A:210:SER:O	2.15	0.46
1:A:122:ASP:OD2	1:A:370:GLU:OE2	2.34	0.46
1:A:217:SER:HB3	1:A:220:ASP:HB2	1.97	0.45
1:A:115:ASN:HD22	1:A:115:ASN:HA	1.65	0.45
1:A:219:LYS:HB2	1:A:305:LEU:HG	1.98	0.45
1:A:290:VAL:HG21	1:A:295:LEU:HD22	1.99	0.45
1:A:326:ARG:HD3	5:A:445:HOH:O	2.16	0.44
1:B:259:LYS:HE2	1:B:289:LEU:CD2	2.48	0.44
1:B:295:LEU:HG	1:B:301:ILE:HD12	1.99	0.44
1:A:248:MET:H	3:A:443:GOL:H12	1.83	0.43
1:B:212:GLY:HA2	1:B:329:TRP:O	2.18	0.43
1:A:13:LEU:HB3	1:A:14:PRO:HD2	2.01	0.43
1:B:60:ARG:HD3	1:B:61:THR:O	2.19	0.43
1:B:181:GLU:HA	1:B:210:SER:O	2.18	0.42
1:B:95:ASP:HB2	1:B:274:LEU:HD12	2.00	0.42
1:A:274:LEU:C	1:A:274:LEU:HD23	2.39	0.42
1:A:54:SER:HB3	1:A:76:TRP:CD1	2.55	0.42
1:A:367:LEU:HD22	1:A:429:SER:CB	2.50	0.42
1:A:428:GLN:NE2	1:A:432:GLY:O	2.51	0.42
1:A:127:ILE:HG23	1:A:137:VAL:HG11	2.03	0.41
1:B:292:ASP:HB3	1:B:295:LEU:HD13	2.02	0.41
1:A:326:ARG:NH2	1:A:345:TYR:OH	2.54	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:432:GLY:C	5:B:684:HOH:O	2.59	0.41
1:B:367:LEU:HD22	1:B:429:SER:OG	2.21	0.41
1:B:54:SER:HB3	1:B:76:TRP:CD1	2.55	0.41
1:B:167:LYS:HE2	1:B:167:LYS:HB3	1.64	0.41
1:A:376:PRO:HA	1:A:406:THR:HG21	2.04	0.40
1:A:177:LYS:HG2	1:A:207:VAL:HG11	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	430/462 (93%)	413 (96%)	16 (4%)	1 (0%)	47	49
1	B	429/462 (93%)	416 (97%)	11 (3%)	2 (0%)	29	26
All	All	859/924 (93%)	829 (96%)	27 (3%)	3 (0%)	41	41

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	150	ILE
1	B	150	ILE
1	B	281	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	347/368 (94%)	340 (98%)	7 (2%)	55	60
1	B	346/368 (94%)	342 (99%)	4 (1%)	71	77
All	All	693/736 (94%)	682 (98%)	11 (2%)	62	69

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	ARG
1	A	115	ASN
1	A	210	SER
1	A	259	LYS
1	A	296	LEU
1	A	305	LEU
1	A	411	GLN
1	B	38	GLN
1	B	60	ARG
1	B	264	LEU
1	B	383	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	115	ASN
1	A	330	HIS
1	A	428	GLN
1	B	330	HIS
1	B	383	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	1PE	A	442	-	15,15,15	0.49	0	14,14,14	0.62	0
4	SO4	B	443	-	4,4,4	0.16	0	6,6,6	0.90	0
2	1PE	B	442	-	15,15,15	0.61	0	14,14,14	0.52	0
3	GOL	A	443	-	5,5,5	0.70	0	5,5,5	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1PE	A	442	-	-	7/13/13/13	-
2	1PE	B	442	-	-	6/13/13/13	-
3	GOL	A	443	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	442	1PE	OH7-C16-C26-OH6
2	B	442	1PE	OH4-C13-C23-OH3
3	A	443	GOL	O1-C1-C2-C3
3	A	443	GOL	C1-C2-C3-O3
2	A	442	1PE	OH2-C12-C22-OH3
2	B	442	1PE	OH7-C16-C26-OH6
3	A	443	GOL	O2-C2-C3-O3
2	A	442	1PE	C12-C22-OH3-C23

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	A	442	1PE	C13-C23-OH3-C22
2	A	442	1PE	OH5-C14-C24-OH4
2	B	442	1PE	C13-C23-OH3-C22
2	B	442	1PE	C23-C13-OH4-C24
2	B	442	1PE	OH5-C14-C24-OH4
2	A	442	1PE	OH4-C13-C23-OH3
2	A	442	1PE	OH6-C15-C25-OH5
2	B	442	1PE	OH6-C15-C25-OH5

There are no ring outliers.

3 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	442	1PE	1	0
2	B	442	1PE	3	0
3	A	443	GOL	8	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	431/462 (93%)	-0.49	4 (0%) 84 86	5, 12, 24, 40	0
1	B	431/462 (93%)	-0.54	3 (0%) 87 89	6, 12, 22, 39	0
All	All	862/924 (93%)	-0.51	7 (0%) 86 88	5, 12, 23, 40	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	291	ASN	3.2
1	B	293	THR	3.1
1	A	293	THR	2.6
1	A	247	ASP	2.5
1	B	171	ASN	2.3
1	A	198	GLU	2.1
1	B	416	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	443	6/6	0.81	0.19	29,29,33,34	0
2	1PE	B	442	16/16	0.92	0.13	14,22,36,38	0
2	1PE	A	442	16/16	0.93	0.11	17,23,29,30	0
4	SO4	B	443	5/5	0.96	0.17	32,32,36,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.