



wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 03:03 am BST

PDB ID : 3GWP
Title : Crystal structure of carbon-sulfur lyase involved in aluminum resistance (YP_878183.1) from CLOSTRIDIUM NOVYI NT at 2.90 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2009-04-01
Resolution : 2.90 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	427	
1	B	427	
1	C	427	
1	D	427	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 12910 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called carbon-sulfur lyase involved in aluminum resistance.



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S	Se			
1	A	424	Total 3223	C 2042	N 529	O 631	P 1	S 9	Se 11	0	0	0
1	B	427	Total 3272	C 2079	N 535	O 637	P 1	S 9	Se 11	0	2	0
1	C	418	Total 3149	C 1999	N 523	O 607	P 1	S 9	Se 10	0	1	0
1	D	427	Total 3266	C 2074	N 539	O 632	P 1	S 9	Se 11	0	2	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	leader sequence	UNP A0Q0N1
B	0	GLY	-	leader sequence	UNP A0Q0N1
C	0	GLY	-	leader sequence	UNP A0Q0N1
D	0	GLY	-	leader sequence	UNP A0Q0N1



- Molecule 1: carbon-sulfur lyase involved in aluminum resistance



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.74Å 108.95Å 207.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.77 – 2.90 29.77 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.77-2.90) 100.0 (29.77-2.90)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	0.24	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.90Å)	Xtrriage
Refinement program	REFMAC 5.2.0019, PHENIX	Depositor
R, R_{free}	0.223 , 0.271 0.228 , 0.272	Depositor DCC
R_{free} test set	2022 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtrriage
Anisotropy	0.394	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 9.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	12910	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/3241	0.55	0/4364
1	B	0.54	0/3301	0.54	0/4445
1	C	0.53	0/3169	0.54	0/4277
1	D	0.54	0/3295	0.54	0/4442
All	All	0.53	0/13006	0.54	0/17528

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3223	0	3101	67	0
1	B	3272	0	3167	56	0
1	C	3149	0	3002	48	0
1	D	3266	0	3149	58	0
All	All	12910	0	12419	218	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 218 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:18:VAL:HG23	1:C:425:VAL:HG21	1.52	0.92
1:D:256:LEU:HD13	1:D:266:THR:HG21	1.52	0.91
1:B:368:MSE:HE2	1:B:375:VAL:HG21	1.57	0.87
1:C:103:LEU:HD23	1:C:238:ALA:HB3	1.61	0.82
1:B:275:ILE:HD13	1:D:98:ASN:HD21	1.46	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	419/427 (98%)	398 (95%)	19 (4%)	2 (0%)	29	61
1	B	426/427 (100%)	405 (95%)	19 (4%)	2 (0%)	29	61
1	C	412/427 (96%)	389 (94%)	22 (5%)	1 (0%)	47	78
1	D	426/427 (100%)	405 (95%)	19 (4%)	2 (0%)	29	61
All	All	1683/1708 (98%)	1597 (95%)	79 (5%)	7 (0%)	34	66

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	126	ASP
1	B	244	ASN
1	D	244	ASN
1	C	244	ASN
1	A	139	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/348 (96%)	331 (99%)	2 (1%)	86	96
1	B	340/348 (98%)	338 (99%)	2 (1%)	86	96
1	C	318/348 (91%)	317 (100%)	1 (0%)	92	98
1	D	336/348 (97%)	336 (100%)	0	100	100
All	All	1327/1392 (95%)	1322 (100%)	5 (0%)	91	97

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	GLU
1	A	137	ASN
1	B	17	ARG
1	B	367	ASP
1	C	137	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	ASN
1	C	30	GLN
1	C	137	ASN
1	D	98	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	243	1	23,24,25	1.76	4 (17%)	25,32,34	1.68	4 (16%)
1	LLP	D	243	1	23,24,25	1.78	4 (17%)	25,32,34	1.72	3 (12%)
1	LLP	B	243	1	23,24,25	1.90	4 (17%)	25,32,34	1.69	3 (12%)
1	LLP	C	243	1	23,24,25	1.82	5 (21%)	25,32,34	1.66	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	243	1	-	7/16/17/19	0/1/1/1
1	LLP	D	243	1	-	3/16/17/19	0/1/1/1
1	LLP	B	243	1	-	3/16/17/19	0/1/1/1
1	LLP	C	243	1	-	5/16/17/19	0/1/1/1

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	243	LLP	O3-C3	-6.18	1.22	1.37
1	D	243	LLP	O3-C3	-5.95	1.23	1.37
1	C	243	LLP	O3-C3	-5.82	1.23	1.37
1	A	243	LLP	O3-C3	-5.68	1.23	1.37
1	B	243	LLP	C4-C4'	3.47	1.53	1.46

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	243	LLP	OP4-C5'-C5	5.84	120.47	109.35
1	C	243	LLP	OP4-C5'-C5	5.54	119.90	109.35
1	B	243	LLP	OP4-C5'-C5	5.44	119.71	109.35
1	A	243	LLP	OP4-C5'-C5	5.42	119.67	109.35
1	B	243	LLP	C4-C4'-NZ	-3.92	106.33	124.31

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	243	LLP	C5'-OP4-P-OP2
1	A	243	LLP	C5'-OP4-P-OP3
1	A	243	LLP	O-C-CA-CB
1	A	243	LLP	C4-C4'-NZ-CE
1	D	243	LLP	C3-C4-C4'-NZ

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	243	LLP	1	0
1	B	243	LLP	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	412/427 (96%)	-0.19	4 (0%) 82 82	20, 24, 31, 39	0
1	B	415/427 (97%)	-0.25	6 (1%) 75 75	20, 24, 32, 43	0
1	C	407/427 (95%)	-0.09	5 (1%) 79 79	20, 24, 30, 37	0
1	D	415/427 (97%)	-0.16	6 (1%) 75 75	19, 24, 31, 40	0
All	All	1649/1708 (96%)	-0.17	21 (1%) 77 77	19, 24, 31, 43	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	135	THR	4.4
1	D	398	GLU	3.4
1	B	369	PRO	3.2
1	A	0	GLY	3.0
1	D	339	ASN	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	A	243	24/25	0.90	0.21	25,32,33,33	0
1	LLP	B	243	24/25	0.90	0.20	25,32,34,34	0
1	LLP	D	243	24/25	0.95	0.15	25,32,33,33	0
1	LLP	C	243	24/25	0.95	0.16	25,32,32,33	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.