



wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 04:29 am BST

PDB ID : 3GYX
Title : Crystal structure of adenylylsulfate reductase from *Desulfovibrio gigas*
Authors : Chiang, Y.-L.; Hsieh, Y.-C.; Liu, E.-H.; Liu, M.-Y.; Chen, C.-J.
Deposited on : 2009-04-06
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	662	<div> <div>76%</div> <div>21%</div> <div>.</div> </div>
1	C	662	<div> <div>73%</div> <div>23%</div> <div>.</div> </div>
1	E	662	<div> <div>%</div> <div>75%</div> <div>21%</div> <div>.</div> </div>
1	G	662	<div> <div>79%</div> <div>20%</div> <div>.</div> </div>
1	I	662	<div> <div>%</div> <div>78%</div> <div>19%</div> <div>.</div> </div>
1	K	662	<div> <div>74%</div> <div>23%</div> <div>.</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	166	<div><div></div><div>3%</div><div>82%</div><div>17%</div><div></div></div>
2	D	166	<div><div></div><div>%</div><div>85%</div><div>13%</div><div></div></div>
2	F	166	<div><div></div><div>4%</div><div>80%</div><div>17%</div><div></div></div>
2	H	166	<div><div></div><div>5%</div><div>83%</div><div>17%</div><div></div></div>
2	J	166	<div><div></div><div>2%</div><div>80%</div><div>17%</div><div></div></div>
2	L	166	<div><div></div><div>2%</div><div>79%</div><div>19%</div><div></div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 39537 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Adenylylsulfate Reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	662	Total	C	N	O	S	0	0	0
			5232	3320	914	957	41			
1	C	662	Total	C	N	O	S	0	0	0
			5232	3320	914	957	41			
1	E	662	Total	C	N	O	S	0	0	0
			5232	3320	914	957	41			
1	G	662	Total	C	N	O	S	0	0	0
			5232	3320	914	957	41			
1	I	662	Total	C	N	O	S	0	0	0
			5232	3320	914	957	41			
1	K	662	Total	C	N	O	S	0	0	0
			5232	3320	914	957	41			

- Molecule 2 is a protein called Adenylylsulfate Reductase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	166	Total	C	N	O	S	0	0	0
			1277	806	209	246	16			
2	D	166	Total	C	N	O	S	0	0	0
			1277	806	209	246	16			
2	F	166	Total	C	N	O	S	0	0	0
			1277	806	209	246	16			
2	H	166	Total	C	N	O	S	0	0	0
			1277	806	209	246	16			
2	J	166	Total	C	N	O	S	0	0	0
			1277	806	209	246	16			
2	L	166	Total	C	N	O	S	0	0	0
			1277	806	209	246	16			

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
3	C	1	Total 53	C 27	N 9	O 15	P 2	0	0
3	E	1	Total 53	C 27	N 9	O 15	P 2	0	0
3	G	1	Total 53	C 27	N 9	O 15	P 2	0	0
3	I	1	Total 53	C 27	N 9	O 15	P 2	0	0
3	K	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 4 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe_4S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	Fe	S	0	0
			8	4	4		
4	B	1	Total	Fe	S	0	0
			8	4	4		
4	D	1	Total	Fe	S	0	0
			8	4	4		
4	D	1	Total	Fe	S	0	0
			8	4	4		
4	F	1	Total	Fe	S	0	0
			8	4	4		
4	F	1	Total	Fe	S	0	0
			8	4	4		
4	H	1	Total	Fe	S	0	0
			8	4	4		
4	H	1	Total	Fe	S	0	0
			8	4	4		
4	J	1	Total	Fe	S	0	0
			8	4	4		
4	J	1	Total	Fe	S	0	0
			8	4	4		
4	L	1	Total	Fe	S	0	0
			8	4	4		
4	L	1	Total	Fe	S	0	0
			8	4	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	11	Total 11	O 11	0	0
5	B	1	Total 1	O 1	0	0
5	C	14	Total 14	O 14	0	0
5	D	3	Total 3	O 3	0	0
5	E	12	Total 12	O 12	0	0
5	F	3	Total 3	O 3	0	0
5	G	6	Total 6	O 6	0	0
5	H	5	Total 5	O 5	0	0
5	I	2	Total 2	O 2	0	0
5	J	3	Total 3	O 3	0	0
5	K	7	Total 7	O 7	0	0
5	L	2	Total 2	O 2	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Adenylylsulfate Reductase





• Molecule 1: Adenylylsulfate Reductase








• Molecule 1: Adenylylsulfate Reductase




• Molecule 1: Adenylylsulfate Reductase




• Molecule 1: Adenylylsulfate Reductase





● Molecule 1: Adenylylsulfate Reductase

Chain K: 74% 23%



● Molecule 2: Adenylylsulfate Reductase


Chain B: 3% 82% 17%





● Molecule 2: Adenylylsulfate Reductase

Chain D: 85% 13%








• Molecule 2: Adenylylsulfate Reductase




• Molecule 2: Adenylylsulfate Reductase



• Molecule 2: Adenylylsulfate Reductase



• Molecule 2: Adenylylsulfate Reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	199.63Å 199.63Å 317.42Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 3.20 29.85 – 3.10	Depositor EDS
% Data completeness (in resolution range)	91.7 (30.00-3.20) 97.7 (29.85-3.10)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.66 (at 3.11Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.192 , 0.245 0.199 , 0.246	Depositor DCC
R_{free} test set	6533 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	53.7	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 26.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.015 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	39537	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/5364	0.53	1/7254 (0.0%)
1	C	0.34	0/5364	0.51	0/7254
1	E	0.35	0/5364	0.51	0/7254
1	G	0.34	0/5364	0.52	0/7254
1	I	0.34	0/5364	0.53	0/7254
1	K	0.34	0/5364	0.52	0/7254
2	B	0.36	0/1309	0.54	0/1773
2	D	0.36	0/1309	0.57	0/1773
2	F	0.37	0/1309	0.54	0/1773
2	H	0.36	0/1309	0.53	0/1773
2	J	0.36	0/1309	0.53	0/1773
2	L	0.36	0/1309	0.52	0/1773
All	All	0.35	0/40038	0.52	1/54162 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	295	LEU	CA-CB-CG	5.86	128.78	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5232	0	5114	86	0
1	C	5232	0	5114	118	0
1	E	5232	0	5114	97	0
1	G	5232	0	5114	74	0
1	I	5232	0	5114	87	0
1	K	5232	0	5114	103	0
2	B	1277	0	1228	15	0
2	D	1277	0	1228	20	0
2	F	1277	0	1228	18	0
2	H	1277	0	1228	17	0
2	J	1277	0	1228	19	0
2	L	1277	0	1228	16	0
3	A	53	0	31	3	0
3	C	53	0	31	1	0
3	E	53	0	31	1	0
3	G	53	0	31	1	0
3	I	53	0	31	2	0
3	K	53	0	31	2	0
4	B	16	0	0	0	0
4	D	16	0	0	1	0
4	F	16	0	0	0	0
4	H	16	0	0	0	0
4	J	16	0	0	0	0
4	L	16	0	0	0	0
5	A	11	0	0	0	0
5	B	1	0	0	0	0
5	C	14	0	0	0	0
5	D	3	0	0	0	0
5	E	12	0	0	0	0
5	F	3	0	0	0	0
5	G	6	0	0	1	0
5	H	5	0	0	0	0
5	I	2	0	0	0	0
5	J	3	0	0	0	0
5	K	7	0	0	0	0
5	L	2	0	0	0	0
All	All	39537	0	38238	611	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 611 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:205:ARG:HH21	1:K:205:ARG:HG3	1.26	0.98
2:D:65:ARG:HH21	2:D:65:ARG:HG2	1.26	0.97
1:I:205:ARG:HG3	1:I:205:ARG:HH21	1.28	0.96
1:G:102:ARG:NH1	1:G:426:ALA:O	1.99	0.95
2:F:65:ARG:HH21	2:F:65:ARG:HG2	1.32	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	660/662 (100%)	600 (91%)	45 (7%)	15 (2%)	6	34
1	C	660/662 (100%)	612 (93%)	42 (6%)	6 (1%)	17	56
1	E	660/662 (100%)	604 (92%)	45 (7%)	11 (2%)	9	42
1	G	660/662 (100%)	618 (94%)	32 (5%)	10 (2%)	10	44
1	I	660/662 (100%)	610 (92%)	36 (6%)	14 (2%)	7	37
1	K	660/662 (100%)	612 (93%)	32 (5%)	16 (2%)	6	34
2	B	164/166 (99%)	147 (90%)	12 (7%)	5 (3%)	4	28
2	D	164/166 (99%)	145 (88%)	17 (10%)	2 (1%)	13	49
2	F	164/166 (99%)	145 (88%)	15 (9%)	4 (2%)	6	34
2	H	164/166 (99%)	142 (87%)	19 (12%)	3 (2%)	8	41
2	J	164/166 (99%)	154 (94%)	7 (4%)	3 (2%)	8	41
2	L	164/166 (99%)	147 (90%)	14 (8%)	3 (2%)	8	41
All	All	4944/4968 (100%)	4536 (92%)	316 (6%)	92 (2%)	8	39

5 of 92 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	132	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	329	HIS
1	A	441	ARG
2	B	17	GLU
2	B	161	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	540/540 (100%)	499 (92%)	41 (8%)	13	45
1	C	540/540 (100%)	494 (92%)	46 (8%)	10	38
1	E	540/540 (100%)	490 (91%)	50 (9%)	9	33
1	G	540/540 (100%)	494 (92%)	46 (8%)	10	38
1	I	540/540 (100%)	493 (91%)	47 (9%)	10	37
1	K	540/540 (100%)	499 (92%)	41 (8%)	13	45
2	B	139/139 (100%)	132 (95%)	7 (5%)	24	60
2	D	139/139 (100%)	130 (94%)	9 (6%)	17	51
2	F	139/139 (100%)	126 (91%)	13 (9%)	8	33
2	H	139/139 (100%)	133 (96%)	6 (4%)	29	64
2	J	139/139 (100%)	127 (91%)	12 (9%)	10	38
2	L	139/139 (100%)	121 (87%)	18 (13%)	4	19
All	All	4074/4074 (100%)	3738 (92%)	336 (8%)	11	41

5 of 336 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	631	LEU
1	G	254	VAL
1	K	449	ASN
2	F	80	LEU
1	G	49	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 94 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	449	ASN
1	G	217	ASN
1	K	256	ASN
1	E	476	HIS
1	G	23	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SF4	D	2000	2	0,12,12	0.00	-	-		
4	SF4	H	2000	2	0,12,12	0.00	-	-		
4	SF4	B	2000	2	0,12,12	0.00	-	-		
4	SF4	D	1000	2	0,12,12	0.00	-	-		
4	SF4	F	1000	2	0,12,12	0.00	-	-		
4	SF4	F	2000	2	0,12,12	0.00	-	-		
4	SF4	B	1000	2	0,12,12	0.00	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SF4	L	1000	2	0,12,12	0.00	-	-		
4	SF4	L	2000	2	0,12,12	0.00	-	-		
4	SF4	H	1000	2	0,12,12	0.00	-	-		
3	FAD	A	1000	-	51,58,58	1.43	6 (11%)	60,89,89	1.59	8 (13%)
4	SF4	J	1000	2	0,12,12	0.00	-	-		
3	FAD	C	1000	-	51,58,58	1.40	6 (11%)	60,89,89	1.62	9 (15%)
3	FAD	E	1000	-	51,58,58	1.43	6 (11%)	60,89,89	1.64	10 (16%)
3	FAD	G	1000	-	51,58,58	1.44	6 (11%)	60,89,89	1.63	8 (13%)
3	FAD	I	1000	-	51,58,58	1.40	6 (11%)	60,89,89	1.59	7 (11%)
3	FAD	K	1000	-	51,58,58	1.41	6 (11%)	60,89,89	1.59	8 (13%)
4	SF4	J	2000	2	0,12,12	0.00	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	D	2000	2	-	-	0/6/5/5
4	SF4	H	2000	2	-	-	0/6/5/5
4	SF4	B	2000	2	-	-	0/6/5/5
4	SF4	D	1000	2	-	-	0/6/5/5
4	SF4	F	1000	2	-	-	0/6/5/5
4	SF4	F	2000	2	-	-	0/6/5/5
4	SF4	B	1000	2	-	-	0/6/5/5
4	SF4	L	1000	2	-	-	0/6/5/5
4	SF4	L	2000	2	-	-	0/6/5/5
4	SF4	H	1000	2	-	-	0/6/5/5
3	FAD	A	1000	-	-	5/30/50/50	0/6/6/6
4	SF4	J	1000	2	-	-	0/6/5/5
3	FAD	C	1000	-	-	2/30/50/50	0/6/6/6
3	FAD	E	1000	-	-	5/30/50/50	0/6/6/6
3	FAD	G	1000	-	-	7/30/50/50	0/6/6/6
3	FAD	I	1000	-	-	10/30/50/50	0/6/6/6
3	FAD	K	1000	-	-	8/30/50/50	0/6/6/6
4	SF4	J	2000	2	-	-	0/6/5/5

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1000	FAD	C10-N1	4.65	1.39	1.33
3	E	1000	FAD	C10-N1	4.60	1.39	1.33
3	K	1000	FAD	C10-N1	4.58	1.39	1.33
3	C	1000	FAD	C10-N1	4.53	1.39	1.33
3	G	1000	FAD	C10-N1	4.51	1.39	1.33

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	1000	FAD	C4-N3-C2	6.13	120.32	115.14
3	G	1000	FAD	C4-N3-C2	6.04	120.24	115.14
3	A	1000	FAD	C4-N3-C2	5.92	120.14	115.14
3	I	1000	FAD	N3A-C2A-N1A	-5.91	119.44	128.68
3	C	1000	FAD	C4-N3-C2	5.88	120.11	115.14

There are no chirality outliers.

5 of 37 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1000	FAD	C5'-O5'-P-O1P
3	A	1000	FAD	C5'-O5'-P-O2P
3	E	1000	FAD	C3'-C4'-C5'-O5'
3	E	1000	FAD	O4'-C4'-C5'-O5'
3	G	1000	FAD	O4'-C4'-C5'-O5'


There are no ring outliers.


7 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	2000	SF4	1	0
3	A	1000	FAD	3	0
3	C	1000	FAD	1	0
3	E	1000	FAD	1	0
3	G	1000	FAD	1	0
3	I	1000	FAD	2	0
3	K	1000	FAD	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.


Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	662/662 (100%)	-0.47	1 (0%) 95 94	32, 40, 56, 69	8 (1%)
1	C	662/662 (100%)	-0.49	2 (0%) 94 92	32, 41, 60, 73	8 (1%)
1	E	662/662 (100%)	-0.35	4 (0%) 89 83	34, 52, 69, 76	8 (1%)
1	G	662/662 (100%)	-0.45	1 (0%) 95 94	34, 44, 56, 63	8 (1%)
1	I	662/662 (100%)	-0.45	8 (1%) 79 67	28, 40, 59, 76	8 (1%)
1	K	662/662 (100%)	-0.46	3 (0%) 91 86	32, 41, 57, 66	9 (1%)
2	B	166/166 (100%)	-0.25	5 (3%) 50 34	34, 46, 73, 76	1 (0%)
2	D	166/166 (100%)	-0.27	2 (1%) 79 67	31, 43, 66, 72	1 (0%)
2	F	166/166 (100%)	-0.01	6 (3%) 42 27	43, 59, 69, 71	1 (0%)
2	H	166/166 (100%)	-0.12	8 (4%) 30 18	33, 44, 72, 78	1 (0%)
2	J	166/166 (100%)	-0.24	3 (1%) 68 55	34, 46, 59, 65	1 (0%)
2	L	166/166 (100%)	-0.20	3 (1%) 68 55	40, 50, 66, 73	1 (0%)
All	All	4968/4968 (100%)	-0.39	46 (0%) 84 75	28, 43, 64, 78	55 (1%)

The worst 5 of 46 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	263	MET	5.3
1	C	263	MET	5.2
1	A	263	MET	4.8
2	B	17	GLU	4.2
2	B	152	GLU	3.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	FAD	A	1000	53/53	0.96	0.14	31,33,38,38	0
3	FAD	I	1000	53/53	0.96	0.15	26,31,37,37	0
3	FAD	K	1000	53/53	0.96	0.15	33,35,38,38	0
3	FAD	G	1000	53/53	0.97	0.14	34,36,38,38	0
3	FAD	C	1000	53/53	0.97	0.15	32,34,41,42	0
3	FAD	E	1000	53/53	0.97	0.14	38,39,44,44	0
4	SF4	F	2000	8/8	0.98	0.06	57,59,60,60	0
4	SF4	J	2000	8/8	0.98	0.07	44,45,46,47	0
4	SF4	L	2000	8/8	0.99	0.08	52,53,54,55	0
4	SF4	H	1000	8/8	0.99	0.14	30,31,32,32	0
4	SF4	H	2000	8/8	0.99	0.07	42,44,44,45	0
4	SF4	J	1000	8/8	0.99	0.12	34,35,35,35	0
4	SF4	B	2000	8/8	0.99	0.09	49,50,50,52	0
4	SF4	D	1000	8/8	0.99	0.14	30,31,31,32	0
4	SF4	F	1000	8/8	0.99	0.13	43,44,45,45	0
4	SF4	D	2000	8/8	0.99	0.07	39,41,42,42	0
4	SF4	B	1000	8/8	0.99	0.12	33,34,34,35	0
4	SF4	L	1000	8/8	0.99	0.13	41,41,42,42	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD I 1000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD K 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD G 1000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD C 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD E 1000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.