



wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 02:54 pm BST

PDB ID : 1GZ4
Title : molecular mechanism of the regulation of human mitochondrial NAD(P)⁺-dependent malic enzyme by ATP and fumarate
Authors : Yang, Z.; Lanks, C.W.; Tong, L.
Deposited on : 2002-05-15
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	551	
1	B	551	
1	C	551	
1	D	551	

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 18289 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-DEPENDENT MALIC ENZYME.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	551	Total	C	N	O	S	Se	0	0	0
			4350	2784	741	802	9	14			
1	B	551	Total	C	N	O	S	Se	0	0	0
			4350	2784	741	802	9	14			
1	C	551	Total	C	N	O	S	Se	0	0	0
			4350	2784	741	802	9	14			
1	D	551	Total	C	N	O	S	Se	0	0	0
			4351	2784	741	803	9	14			


- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is TARTRONATE (three-letter code: TTN) (formula: $C_3H_2O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	3	5		
3	B	1	Total	C	O	0	0
			8	3	5		
3	C	1	Total	C	O	0	0
			8	3	5		
3	D	1	Total	C	O	0	0
			8	3	5		

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Mn 1 1	0	0
4	A	1	Total Mn 1 1	0	0
4	D	1	Total Mn 1 1	0	0
4	C	1	Total Mn 1 1	0	0

- Molecule 5 is FUMARIC ACID (three-letter code: FUM) (formula: C₄H₄O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 8 4 4	0	0
5	B	1	Total C O 8 4 4	0	0
5	C	1	Total C O 8 4 4	0	0
5	D	1	Total C O 8 4 4	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	150	Total O 150 150	0	0
6	B	142	Total O 142 142	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	130	Total	O	0	0
			130	130		
6	D	150	Total	O	0	0
			150	150		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: NAD-DEPENDENT MALIC ENZYME




• Molecule 1: NAD-DEPENDENT MALIC ENZYME




• Molecule 1: NAD-DEPENDENT MALIC ENZYME

Chain C:  73% 25%



• Molecule 1: NAD-DEPENDENT MALIC ENZYME

Chain D:  73% 24%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	224.94Å 117.45Å 111.61Å 90.00° 109.33° 90.00°	Depositor
Resolution (Å)	20.00 – 2.20	Depositor
% Data completeness (in resolution range)	95.0 (20.00-2.20)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.209 , 0.245	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	18289	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUM, TTN, ATP, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/4430	0.58	0/5976
1	B	0.35	0/4430	0.60	0/5976
1	C	0.35	0/4430	0.60	1/5976 (0.0%)
1	D	0.35	0/4431	0.59	0/5976
All	All	0.35	0/17721	0.59	1/23904 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	91	ARG	NE-CZ-NH2	-6.08	117.26	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4350	0	4383	107	0
1	B	4350	0	4383	137	0
1	C	4350	0	4383	122	0
1	D	4351	0	4383	123	0
2	A	62	0	24	0	0
2	B	62	0	24	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	62	0	24	1	0
2	D	62	0	24	0	0
3	A	8	0	1	1	0
3	B	8	0	1	1	0
3	C	8	0	1	1	0
3	D	8	0	1	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	8	0	2	1	0
5	B	8	0	2	0	0
5	C	8	0	2	0	0
5	D	8	0	2	0	0
6	A	150	0	0	6	0
6	B	142	0	0	3	0
6	C	130	0	0	3	0
6	D	150	0	0	4	0
All	All	18289	0	17640	479	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 479 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:29:MSE:HE2	1:B:50:LEU:HD22	1.28	1.15
1:B:315:ALA:HB3	1:B:392:VAL:HG21	1.25	1.10
1:A:381:VAL:HG13	1:A:407:MSE:HE3	1.38	1.06
1:C:286:VAL:HG21	1:C:467:ASN:HA	1.39	1.04
1:D:343:MSE:HE2	1:D:365:PHE:HB2	1.41	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	549/551 (100%)	531 (97%)	17 (3%)	1 (0%)	47	55
1	B	549/551 (100%)	530 (96%)	17 (3%)	2 (0%)	34	37
1	C	549/551 (100%)	528 (96%)	20 (4%)	1 (0%)	47	55
1	D	549/551 (100%)	527 (96%)	21 (4%)	1 (0%)	47	55
All	All	2196/2204 (100%)	2116 (96%)	75 (3%)	5 (0%)	47	55

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	397	ARG
1	B	397	ARG
1	D	397	ARG
1	C	392	VAL
1	B	392	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	467/453 (103%)	445 (95%)	22 (5%)	26	33
1	B	467/453 (103%)	448 (96%)	19 (4%)	30	39
1	C	467/453 (103%)	445 (95%)	22 (5%)	26	33
1	D	467/453 (103%)	438 (94%)	29 (6%)	18	21
All	All	1868/1812 (103%)	1776 (95%)	92 (5%)	25	31

5 of 92 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	66	LEU
1	C	300	LYS
1	D	371	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	70	ARG
1	C	133	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	482	ASN
1	C	154	HIS
1	D	261	ASN
1	B	518	ASN
1	B	520	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 4 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	FUM	B	605	-	1,7,7	1.70	0	2,8,8	0.55	0
5	FUM	D	605	-	1,7,7	1.70	0	2,8,8	0.47	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ATP	B	602	-	26,33,33	1.60	6 (23%)	31,52,52	1.57	3 (9%)
3	TTN	B	603	4	1,7,7	0.11	0	0,9,9	0.00	-
3	TTN	A	603	4	1,7,7	0.04	0	0,9,9	0.00	-
3	TTN	D	603	4	1,7,7	0.17	0	0,9,9	0.00	-
2	ATP	A	601	-	26,33,33	1.55	6 (23%)	31,52,52	1.52	3 (9%)
5	FUM	C	605	-	1,7,7	2.00	1 (100%)	2,8,8	1.06	0
2	ATP	B	601	-	26,33,33	1.62	5 (19%)	31,52,52	1.50	3 (9%)
3	TTN	C	603	4	1,7,7	0.21	0	0,9,9	0.00	-
5	FUM	A	605	-	1,7,7	1.67	0	2,8,8	0.50	0
2	ATP	D	601	-	26,33,33	1.59	6 (23%)	31,52,52	1.52	3 (9%)
2	ATP	C	602	-	26,33,33	1.62	6 (23%)	31,52,52	1.54	3 (9%)
2	ATP	A	602	-	26,33,33	1.71	6 (23%)	31,52,52	1.55	3 (9%)
2	ATP	D	602	-	26,33,33	1.63	6 (23%)	31,52,52	1.54	3 (9%)
2	ATP	C	601	-	26,33,33	1.58	6 (23%)	31,52,52	1.52	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FUM	B	605	-	-	0/0/5/5	-
5	FUM	D	605	-	-	0/0/5/5	-
2	ATP	B	602	-	-	2/18/38/38	0/3/3/3
3	TTN	B	603	4	-	0/0/8/8	-
3	TTN	A	603	4	-	0/0/8/8	-
3	TTN	D	603	4	-	0/0/8/8	-
2	ATP	A	601	-	-	3/18/38/38	0/3/3/3
5	FUM	C	605	-	-	0/0/5/5	-
2	ATP	B	601	-	-	3/18/38/38	0/3/3/3
3	TTN	C	603	4	-	0/0/8/8	-
5	FUM	A	605	-	-	0/0/5/5	-
2	ATP	D	601	-	-	2/18/38/38	0/3/3/3
2	ATP	C	602	-	-	2/18/38/38	0/3/3/3
2	ATP	A	602	-	-	2/18/38/38	0/3/3/3
2	ATP	D	602	-	-	2/18/38/38	0/3/3/3
2	ATP	C	601	-	-	3/18/38/38	0/3/3/3

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	602	ATP	O4'-C1'	4.35	1.47	1.41
2	D	601	ATP	O4'-C1'	4.07	1.46	1.41
2	B	601	ATP	O4'-C1'	3.89	1.46	1.41
2	C	602	ATP	O4'-C1'	3.85	1.46	1.41
2	D	602	ATP	O4'-C1'	3.61	1.46	1.41

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	601	ATP	N3-C2-N1	-5.18	120.58	128.68
2	A	601	ATP	N3-C2-N1	-5.17	120.60	128.68
2	A	602	ATP	N3-C2-N1	-5.16	120.61	128.68
2	B	602	ATP	N3-C2-N1	-5.16	120.61	128.68
2	D	601	ATP	N3-C2-N1	-5.15	120.64	128.68

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	602	ATP	PG-O3B-PB-O1B
2	A	602	ATP	PG-O3B-PB-O1B
2	B	602	ATP	PG-O3B-PB-O1B
2	B	602	ATP	PG-O3B-PB-O2B
2	B	601	ATP	PG-O3B-PB-O2B


There are no ring outliers.


7 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	603	TTN	1	0
3	A	603	TTN	1	0
3	D	603	TTN	1	0
2	B	601	ATP	1	0
3	C	603	TTN	1	0
5	A	605	FUM	1	0
2	C	601	ATP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.


Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.