



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 17, 2020 – 07:47 pm BST

PDB ID : 1H18  
Title : Pyruvate Formate-Lyase (E.coli) in complex with Pyruvate  
Authors : Becker, A.; Kabsch, W.  
Deposited on : 2002-07-04  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	759	<div> <div>6%</div> <div>81%</div> <div>18%</div> <div>.</div> </div>
1	B	759	<div> <div>7%</div> <div>76%</div> <div>23%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PYR	B	1001	-	-	X	-
5	PG4	B	9013	-	-	-	X

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 13667 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FORMATE ACETYLTRANSFERASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	759	Total	C	N	O	S	0	39	0
			6172	3906	1050	1174	42			
1	B	759	Total	C	N	O	S	0	39	0
			6172	3906	1050	1174	42			

- Molecule 2 is PYRUVIC ACID (three-letter code: PYR) (formula: C<sub>3</sub>H<sub>4</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	4	Total	Na	0	0
			4	4		
3	A	3	Total	Na	0	0
			3	3		

- Molecule 4 is L-TREITOL (three-letter code: DTL) (formula:  $C_4H_{10}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	4	4		
4	A	1	Total	C	O	0	0
			8	4	4		
4	B	1	Total	C	O	0	0
			8	4	4		
4	B	1	Total	C	O	0	0
			8	4	4		

- Molecule 5 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula:  $C_8H_{18}O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			13	8	5		
5	A	1	Total	C	O	0	0
			13	8	5		
5	A	1	Total	C	O	0	0
			13	8	5		
5	B	1	Total	C	O	0	0
			13	8	5		
5	B	1	Total	C	O	0	0
			13	8	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	633	Total	O	0	0
			633	633		
6	B	574	Total	O	0	0
			574	574		


### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: FORMATE ACETYLTRANSFERASE 1



#### • Molecule 1: FORMATE ACETYLTRANSFERASE 1







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	158.91Å 158.91Å 159.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.30 48.16 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.2 (15.00-2.30) 95.3 (48.16-2.30)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.55 (at 2.29Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.183 , 0.234 0.182 , 0.230	Depositor DCC
$R_{free}$ test set	1734 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.5	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 50.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for -h,l,k 0.004 for -l,-k,-h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	13667	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PYR, PG4, DTL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	2.04	4/6477 (0.1%)	0.86	6/8742 (0.1%)
1	B	0.46	0/6477	0.65	0/8742
All	All	1.48	4/12954 (0.0%)	0.76	6/17484 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	146[A]	GLN	CD-NE2	112.60	4.14	1.32
1	A	146[B]	GLN	CD-NE2	112.60	4.14	1.32
1	A	146[A]	GLN	CB-CG	12.89	1.87	1.52
1	A	146[B]	GLN	CB-CG	12.89	1.87	1.52

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	146[A]	GLN	CG-CD-NE2	-30.02	44.66	116.70
1	A	146[B]	GLN	CG-CD-NE2	-30.02	44.66	116.70
1	A	146[A]	GLN	OE1-CD-NE2	-17.91	80.70	121.90
1	A	146[B]	GLN	OE1-CD-NE2	-17.91	80.70	121.90
1	A	146[A]	GLN	CA-CB-CG	-9.83	91.77	113.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	146[A]	GLN	Sidechain
1	A	181	TYR	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6172	0	6153	122	0
1	B	6172	0	6153	140	0
2	A	6	0	3	2	0
2	B	6	0	3	4	0
3	A	3	0	0	0	0
3	B	4	0	0	0	0
4	A	16	0	20	2	0
4	B	16	0	20	2	0
5	A	39	0	54	2	0
5	B	26	0	36	1	0
6	A	633	0	0	8	0
6	B	574	0	0	14	0
All	All	13667	0	12442	262	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 262 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146[A]:GLN:CB	1:A:146[A]:GLN:CG	1.87	1.49
1:A:132:MET:SD	6:A:2183:HOH:O	1.97	1.22
1:A:110[B]:ILE:HD12	1:A:270:ASN:HB3	1.49	0.94
1:A:146[A]:GLN:CA	1:A:146[A]:GLN:CG	2.46	0.93
1:B:606:ILE:HG22	1:B:607:THR:H	1.32	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	799/759 (105%)	763 (96%)	34 (4%)	2 (0%)	41	50
1	B	799/759 (105%)	759 (95%)	36 (4%)	4 (0%)	29	35
All	All	1598/1518 (105%)	1522 (95%)	70 (4%)	6 (0%)	34	42

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	733	SER
1	B	606	ILE
1	A	606	ILE
1	B	554	ALA
1	B	733	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	680/638 (107%)	666 (98%)	14 (2%)	53	70
1	B	680/638 (107%)	671 (99%)	9 (1%)	69	82
All	All	1360/1276 (107%)	1337 (98%)	23 (2%)	62	76

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	468[B]	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	607	THR
1	B	435	ARG
1	A	490	TYR
1	A	733	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	706	ASN
1	B	92	GLN
1	B	358	ASN
1	A	410	ASN
1	B	250	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 7 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	DTL	B	9009	-	7,7,7	0.62	0	8,8,8	0.60	0
2	PYR	B	1001	-	2,5,5	0.81	0	2,6,6	1.49	0
5	PG4	A	9011	-	12,12,12	0.51	0	11,11,11	0.54	0
5	PG4	B	9011	-	12,12,12	0.60	0	11,11,11	0.47	0
4	DTL	A	9009	-	7,7,7	0.40	0	8,8,8	0.56	0
5	PG4	A	9012	-	12,12,12	0.61	0	11,11,11	0.55	0
5	PG4	B	9013	-	12,12,12	0.70	0	11,11,11	0.44	0
5	PG4	A	9013	-	12,12,12	0.66	0	11,11,11	0.44	0
2	PYR	A	1001	-	2,5,5	1.01	0	2,6,6	1.38	0
4	DTL	B	9010	-	7,7,7	0.61	0	8,8,8	0.21	0
4	DTL	A	9010	-	7,7,7	0.41	0	8,8,8	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTL	B	9009	-	-	0/8/8/8	-
2	PYR	B	1001	-	-	0/0/4/4	-
5	PG4	A	9011	-	-	6/10/10/10	-
5	PG4	B	9011	-	-	6/10/10/10	-
4	DTL	A	9009	-	-	2/8/8/8	-
5	PG4	A	9012	-	-	7/10/10/10	-
5	PG4	B	9013	-	-	8/10/10/10	-
5	PG4	A	9013	-	-	3/10/10/10	-
2	PYR	A	1001	-	-	0/0/4/4	-
4	DTL	B	9010	-	-	1/8/8/8	-
4	DTL	A	9010	-	-	1/8/8/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	9010	DTL	C1-C2-C3-C4
4	A	9010	DTL	C1-C2-C3-C4
5	A	9011	PG4	O2-C3-C4-O3
5	A	9011	PG4	O3-C5-C6-O4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
5	B	9011	PG4	O2-C3-C4-O3

There are no ring outliers.

6 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1001	PYR	4	0
5	B	9013	PG4	1	0
5	A	9013	PG4	2	0
2	A	1001	PYR	2	0
4	B	9010	DTL	2	0
4	A	9010	DTL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	759/759 (100%)	-0.01	46 (6%)	21 27	32, 48, 67, 115	26 (3%)
1	B	759/759 (100%)	0.13	56 (7%)	14 19	33, 51, 81, 113	27 (3%)
All	All	1518/1518 (100%)	0.06	102 (6%)	17 23	32, 50, 77, 115	53 (3%)

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	SER	12.4
1	B	1	SER	11.7
1	B	2	GLU	7.2
1	A	2	GLU	5.0
1	B	419	CYS	4.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	PG4	B	9013	13/13	0.59	0.53	104,105,106,106	0
3	NA	A	9004	1/1	0.64	0.17	95,95,95,95	0
4	DTL	B	9010	8/8	0.71	0.35	83,88,90,92	0
5	PG4	B	9011	13/13	0.72	0.27	87,89,92,93	0
5	PG4	A	9013	13/13	0.76	0.25	89,91,102,102	0
3	NA	B	9006	1/1	0.80	0.32	77,77,77,77	0
4	DTL	A	9010	8/8	0.81	0.30	81,83,84,85	0
5	PG4	A	9012	13/13	0.81	0.25	74,76,93,95	0
5	PG4	A	9011	13/13	0.81	0.21	88,91,95,95	0
4	DTL	B	9009	8/8	0.83	0.22	59,67,70,72	0
3	NA	B	9003	1/1	0.85	0.52	77,77,77,77	0
4	DTL	A	9009	8/8	0.88	0.17	71,75,77,77	0
2	PYR	B	1001	6/6	0.94	0.34	53,55,55,55	0
3	NA	A	9001	1/1	0.95	0.05	69,69,69,69	0
3	NA	B	9001	1/1	0.95	0.07	70,70,70,70	0
3	NA	A	9002	1/1	0.97	0.47	92,92,92,92	0
3	NA	B	9007	1/1	0.98	0.16	97,97,97,97	0
2	PYR	A	1001	6/6	0.99	0.28	40,42,44,45	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.