



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2021 – 05:14 AM EDT

PDB ID : 3H3N
Title : Glycerol Kinase H232R with Glycerol
Authors : Yeh, J.I.; Kettering, R.D.
Deposited on : 2009-04-16
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	O	506	<div> <div>6%</div> <div>81%</div> <div>17%</div> <div>..</div> </div>
1	X	506	<div> <div>7%</div> <div>84%</div> <div>13%</div> <div>..</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycerol kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	501	Total	C	N	O	S	0	0	0
			3885	2461	649	761	14			
1	O	501	Total	C	N	O	S	0	0	0
			3890	2465	650	761	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	232	ARG	HIS	engineered mutation	UNP O34153
O	232	ARG	HIS	engineered mutation	UNP O34153

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	X	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	O	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	X	1	Total	C	O	0	0
			4	2	2		
3	X	1	Total	C	O	0	0
			4	2	2		
3	X	1	Total	C	O	0	0
			4	2	2		
3	O	1	Total	C	O	0	0
			4	2	2		
3	O	1	Total	C	O	0	0
			4	2	2		
3	O	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	X	1	Total	O	P	0	0
			5	4	1		
4	O	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	X	164	Total	O	0	0
			164	164		
5	O	148	Total	O	0	0
			148	148		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycerol kinase



- Molecule 1: Glycerol kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	96.81Å 200.46Å 56.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.73 9.98 – 1.73	Depositor EDS
% Data completeness (in resolution range)	98.6 (10.00-1.73) 99.8 (9.98-1.73)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 1.73Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.207 , 0.216 0.211 , 0.221	Depositor DCC
R_{free} test set	5762 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtriage
Anisotropy	0.066	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 51.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8133	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, GOL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	O	0.37	2/3970 (0.1%)	0.58	1/5380 (0.0%)
1	X	0.42	6/3965 (0.2%)	0.62	3/5374 (0.1%)
All	All	0.40	8/7935 (0.1%)	0.60	4/10754 (0.0%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	O	85	GLU	CD-OE1	-7.94	1.17	1.25
1	O	85	GLU	C-O	-7.49	1.09	1.23
1	X	461	LEU	C-O	-7.44	1.09	1.23
1	X	303	LEU	C-O	-6.51	1.10	1.23
1	X	501	ALA	C-O	-5.98	1.11	1.23
1	X	303	LEU	CG-CD2	-5.60	1.31	1.51
1	X	501	ALA	CA-CB	-5.53	1.40	1.52
1	X	461	LEU	CG-CD2	-5.12	1.32	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	303	LEU	CA-CB-CG	7.35	132.21	115.30
1	X	461	LEU	CB-CG-CD2	5.51	120.37	111.00
1	X	282	LEU	N-CA-CB	-5.37	99.66	110.40
1	O	240	ILE	N-CA-C	-5.14	97.13	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	O	3890	0	3798	60	0
1	X	3885	0	3786	61	0
2	O	6	0	8	0	0
2	X	6	0	8	0	0
3	O	12	0	18	1	0
3	X	12	0	18	0	0
4	O	5	0	0	0	0
4	X	5	0	0	0	0
5	O	148	0	0	1	0
5	X	164	0	0	2	0
All	All	8133	0	7636	121	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (121) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:178:GLN:HG3	1:X:229:ARG:HH21	1.26	0.98
1:O:221:ASN:HD22	1:O:294:GLY:H	1.15	0.95
1:X:436:ASN:H	1:X:436:ASN:HD22	1.17	0.90
1:X:12:GLN:HE22	1:X:83:GLN:HE21	1.10	0.90
1:X:221:ASN:HD22	1:X:294:GLY:H	1.21	0.88
1:O:12:GLN:HE22	1:O:83:GLN:HE21	1.17	0.88
1:X:178:GLN:HG3	1:X:229:ARG:NH2	1.94	0.82
1:X:178:GLN:CG	1:X:229:ARG:HH21	1.95	0.80
1:O:221:ASN:ND2	1:O:294:GLY:H	1.81	0.79
1:X:85:GLU:OE2	1:X:189:ARG:HD3	1.84	0.76
1:O:112:ILE:HD13	1:O:143:ARG:HG3	1.66	0.75
1:O:133:ILE:HA	1:O:191:MET:HE1	1.69	0.75
1:X:221:ASN:ND2	1:X:294:GLY:H	1.89	0.70
1:X:229:ARG:NH1	1:X:231:TYR:OH	2.24	0.69
1:X:281:GLN:O	1:X:282:LEU:HB3	1.91	0.69
1:X:28:LYS:HD2	1:X:31:SER:HB2	1.73	0.68
1:X:12:GLN:HE21	1:X:166:ILE:HG21	1.61	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:341:ASN:HD21	1:O:383:GLN:HE22	1.41	0.66
1:X:436:ASN:HD22	1:X:436:ASN:N	1.85	0.65
1:O:143:ARG:O	1:O:143:ARG:HD3	1.96	0.65
1:X:436:ASN:HB2	1:X:438:GLU:OE1	1.96	0.65
1:O:68:SER:HB2	1:O:70:ILE:HD13	1.80	0.64
1:X:176:ASP:HB3	1:X:229:ARG:NH1	2.13	0.63
1:O:12:GLN:HE21	1:O:166:ILE:HG21	1.63	0.63
1:X:12:GLN:NE2	1:X:83:GLN:HE21	1.92	0.63
1:X:181:VAL:HG22	1:X:182:THR:N	2.17	0.60
1:O:379:ARG:O	1:O:383:GLN:HG3	2.02	0.59
1:O:329:GLN:HE21	1:O:333:LEU:HD11	1.66	0.59
1:O:402:ILE:HG22	1:O:403:ASP:N	2.17	0.59
1:X:263:LYS:HD2	1:X:263:LYS:C	2.24	0.57
1:X:283:SER:O	1:X:399:ASP:HB3	2.03	0.57
1:X:397:LYS:HG3	1:X:402:ILE:O	2.04	0.57
1:X:438:GLU:HG2	1:X:438:GLU:O	2.03	0.57
1:O:436:ASN:HB3	1:O:438:GLU:OE2	2.05	0.57
1:O:457:ASP:OD2	1:O:459:ASP:HB2	2.04	0.57
1:O:438:GLU:H	1:O:438:GLU:CD	2.08	0.57
1:O:390:LYS:HD2	1:O:426:ILE:HG22	1.87	0.56
1:O:433:ARG:O	1:O:468:GLY:HA3	2.05	0.56
1:X:322:ARG:HD2	5:X:580:HOH:O	2.06	0.56
1:O:327:SER:HB2	1:O:328:PRO:HD3	1.88	0.55
1:O:181:VAL:HG22	1:O:182:THR:N	2.21	0.55
1:X:189:ARG:HH22	1:X:288:LEU:CD1	2.20	0.55
1:O:12:GLN:NE2	1:O:83:GLN:HE21	1.97	0.54
1:O:314:ILE:HD11	1:O:382:LEU:HD23	1.90	0.54
1:X:189:ARG:HH22	1:X:288:LEU:HD13	1.73	0.53
1:O:27:LYS:HZ3	1:O:27:LYS:HB3	1.74	0.52
1:X:436:ASN:N	1:X:436:ASN:ND2	2.57	0.52
1:O:81:THR:OG1	1:O:246:ASP:HA	2.09	0.52
1:O:298:LYS:HE2	1:O:300:TYR:OH	2.09	0.52
1:X:81:THR:OG1	1:X:246:ASP:HA	2.10	0.52
1:O:180:HIS:CE1	1:O:216:PRO:HB3	2.45	0.52
1:O:484:TYR:CE2	1:O:488:LYS:HD2	2.45	0.51
1:O:263:LYS:HD2	1:O:263:LYS:C	2.30	0.51
1:X:229:ARG:HH11	1:X:229:ARG:HG3	1.74	0.51
1:O:125:ILE:HD12	1:O:133:ILE:HG12	1.92	0.51
1:X:379:ARG:O	1:X:383:GLN:HG3	2.11	0.50
1:X:229:ARG:NH1	1:X:229:ARG:HG3	2.27	0.50
1:X:438:GLU:OE1	1:X:438:GLU:N	2.42	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:279:GLU:C	1:X:281:GLN:H	2.15	0.50
1:O:175:THR:O	1:O:178:GLN:HG2	2.12	0.50
1:O:84:ARG:O	1:O:85:GLU:HB2	2.11	0.49
1:X:81:THR:HG21	1:X:440:THR:HG22	1.95	0.49
1:O:124:MET:SD	1:O:128:LYS:HD2	2.53	0.49
1:X:280:PRO:HD3	1:X:301:TYR:CD2	2.48	0.49
1:X:180:HIS:HD2	5:X:574:HOH:O	1.96	0.48
1:X:331:GLU:HG3	1:X:416:ASN:ND2	2.28	0.48
1:O:118:VAL:HG13	1:O:119:ASP:OD1	2.14	0.48
1:X:241:ALA:HB1	1:X:451:ALA:HB3	1.96	0.47
1:O:241:ALA:HB1	1:O:451:ALA:HB3	1.95	0.47
1:O:27:LYS:HB3	1:O:27:LYS:NZ	2.29	0.47
1:O:110:SER:OG	1:O:111:PRO:HD3	2.14	0.46
1:X:172:TRP:CE2	1:X:177:GLY:HA2	2.50	0.46
1:O:492:ALA:O	1:O:496:THR:HG23	2.16	0.46
1:O:254:GLN:NE2	1:O:410:ASP:OD2	2.50	0.45
1:X:433:ARG:O	1:X:468:GLY:HA3	2.16	0.45
1:X:282:LEU:N	1:X:282:LEU:HD23	2.32	0.45
1:X:266:TYR:HB3	1:X:413:ALA:HB3	1.99	0.45
1:O:386:ALA:HA	1:O:423:GLN:HE22	1.82	0.45
1:O:390:LYS:HD3	1:O:484:TYR:CD1	2.52	0.44
1:X:181:VAL:HG22	1:X:182:THR:H	1.81	0.44
1:X:225:TYR:CE1	1:X:243:MET:HG3	2.53	0.44
1:X:280:PRO:O	1:X:282:LEU:HD23	2.17	0.44
1:X:42:LYS:HE3	1:X:42:LYS:HB2	1.84	0.44
1:O:132:VAL:O	1:O:191:MET:HE1	2.18	0.44
1:X:87:THR:HG23	1:X:163:PHE:CE1	2.53	0.43
1:X:184:TYR:HB3	1:X:291:ILE:HG21	1.99	0.43
1:X:189:ARG:NH2	1:X:288:LEU:HD13	2.33	0.43
1:X:150:GLU:N	1:X:150:GLU:OE1	2.51	0.43
1:X:255:MET:SD	1:X:462:LYS:HA	2.58	0.43
1:O:282:LEU:N	1:O:282:LEU:HD22	2.33	0.43
1:X:92:LYS:HG3	1:X:93:THR:N	2.34	0.42
1:O:87:THR:HG23	1:O:163:PHE:CE1	2.54	0.42
1:X:181:VAL:CG2	1:X:182:THR:N	2.82	0.42
1:O:403:ASP:OD2	1:O:404:ILE:N	2.53	0.42
1:O:125:ILE:HD12	1:O:133:ILE:CD1	2.50	0.42
1:O:265:THR:HG23	1:O:410:ASP:OD1	2.20	0.42
1:X:324:ILE:HG22	1:X:333:LEU:CD1	2.50	0.42
1:O:184:TYR:HB3	1:O:291:ILE:HG21	2.01	0.42
1:O:435:ALA:HB3	1:O:466:GLU:HG2	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:54:TRP:CD1	1:O:173:LYS:HE2	2.55	0.41
1:O:115:GLN:HA	1:O:118:VAL:HG12	2.01	0.41
1:O:266:TYR:HB3	1:O:413:ALA:HB3	2.02	0.41
1:X:30:GLY:HA3	1:X:68:SER:HB3	2.01	0.41
1:O:402:ILE:CG2	1:O:403:ASP:N	2.81	0.41
1:O:81:THR:HG21	1:O:440:THR:HG22	2.03	0.41
1:X:282:LEU:N	1:X:282:LEU:CD2	2.84	0.41
1:O:110:SER:N	1:O:111:PRO:CD	2.84	0.41
1:O:181:VAL:HG22	1:O:182:THR:H	1.85	0.41
1:O:217:GLU:HG3	5:O:555:HOH:O	2.21	0.41
1:O:435:ALA:HB3	1:O:466:GLU:CG	2.51	0.41
1:X:178:GLN:CD	1:X:229:ARG:HH21	2.24	0.41
1:X:18:ARG:HG3	1:X:32:SER:O	2.21	0.40
1:O:262:ILE:HD11	1:O:272:ILE:CG2	2.51	0.40
1:X:110:SER:OG	1:X:111:PRO:HD3	2.21	0.40
1:X:110:SER:N	1:X:111:PRO:CD	2.84	0.40
1:O:183:ASP:OD2	1:O:185:SER:OG	2.27	0.40
1:O:427:LEU:HB3	1:O:429:ILE:HG12	2.04	0.40
1:X:180:HIS:CE1	1:X:216:PRO:HB3	2.56	0.40
1:X:238:VAL:HA	1:X:239:PRO:HD3	1.88	0.40
1:X:475:MET:HA	1:X:476:PRO:HD3	1.97	0.40
1:O:382:LEU:HD23	3:O:3126:EDO:H12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	O	499/506 (99%)	492 (99%)	7 (1%)	0	100	100
1	X	499/506 (99%)	488 (98%)	10 (2%)	1 (0%)	47	29
All	All	998/1012 (99%)	980 (98%)	17 (2%)	1 (0%)	51	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	281	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	O	406/410 (99%)	399 (98%)	7 (2%)	60	41
1	X	405/410 (99%)	398 (98%)	7 (2%)	60	41
All	All	811/820 (99%)	797 (98%)	14 (2%)	60	41

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	12	GLN
1	X	150	GLU
1	X	165	THR
1	X	189	ARG
1	X	263	LYS
1	X	436	ASN
1	X	438	GLU
1	O	12	GLN
1	O	127	GLU
1	O	165	THR
1	O	263	LYS
1	O	303	LEU
1	O	438	GLU
1	O	461	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	12	GLN
1	X	121	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	180	HIS
1	X	202	GLN
1	X	221	ASN
1	X	341	ASN
1	X	423	GLN
1	X	436	ASN
1	X	469	GLN
1	X	489	GLN
1	O	12	GLN
1	O	58	GLN
1	O	126	HIS
1	O	180	HIS
1	O	202	GLN
1	O	221	ASN
1	O	285	ASN
1	O	315	GLN
1	O	329	GLN
1	O	383	GLN
1	O	423	GLN
1	O	432	GLN
1	O	489	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	X	1000	-	5,5,5	0.42	0	5,5,5	0.26	0
3	EDO	X	3122	-	3,3,3	0.44	0	2,2,2	0.29	0
2	GOL	O	1001	-	5,5,5	0.33	0	5,5,5	0.29	0
4	PO4	O	507	-	4,4,4	1.63	0	6,6,6	0.44	0
3	EDO	X	3125	-	3,3,3	0.45	0	2,2,2	0.32	0
3	EDO	O	3124	-	3,3,3	0.45	0	2,2,2	0.36	0
3	EDO	O	3127	-	3,3,3	0.45	0	2,2,2	0.34	0
3	EDO	O	3126	-	3,3,3	0.45	0	2,2,2	0.31	0
3	EDO	X	3123	-	3,3,3	0.46	0	2,2,2	0.34	0
4	PO4	X	507	-	4,4,4	1.66	0	6,6,6	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	X	1000	-	-	0/4/4/4	-
3	EDO	X	3122	-	-	0/1/1/1	-
2	GOL	O	1001	-	-	0/4/4/4	-
3	EDO	X	3125	-	-	0/1/1/1	-
3	EDO	O	3124	-	-	0/1/1/1	-
3	EDO	O	3127	-	-	0/1/1/1	-
3	EDO	O	3126	-	-	0/1/1/1	-
3	EDO	X	3123	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	O	3126	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	O	501/506 (99%)	0.40	28 (5%) 24 29	23, 35, 54, 73	0
1	X	501/506 (99%)	0.38	36 (7%) 15 19	21, 34, 55, 79	0
All	All	1002/1012 (99%)	0.39	64 (6%) 19 23	21, 34, 55, 79	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	281	GLN	8.2
1	X	282	LEU	6.0
1	X	340	ASP	5.9
1	O	401	GLY	5.9
1	X	280	PRO	5.7
1	O	402	ILE	5.2
1	O	467	GLU	4.8
1	X	467	GLU	4.8
1	X	199	GLU	4.3
1	O	340	ASP	4.2
1	O	282	LEU	4.0
1	O	123	GLU	4.0
1	X	206	ASP	3.9
1	O	466	GLU	3.9
1	X	403	ASP	3.6
1	O	127	GLU	3.5
1	X	150	GLU	3.5
1	X	3	GLU	3.4
1	X	2	ALA	3.4
1	X	279	GLU	3.3
1	O	206	ASP	3.1
1	O	459	ASP	3.1
1	O	500	LYS	3.1
1	X	119	ASP	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	401	GLY	3.0
1	O	403	ASP	3.0
1	O	436	ASN	2.9
1	X	338	LYS	2.8
1	O	279	GLU	2.8
1	O	118	VAL	2.8
1	X	172	TRP	2.8
1	O	156	ALA	2.7
1	X	402	ILE	2.7
1	O	150	GLU	2.6
1	X	325	GLU	2.6
1	O	437	LEU	2.6
1	O	332	GLU	2.6
1	X	126	HIS	2.5
1	O	465	ALA	2.5
1	X	477	ALA	2.4
1	X	474	GLU	2.4
1	X	437	LEU	2.4
1	X	93	THR	2.4
1	X	438	GLU	2.4
1	X	466	GLU	2.4
1	X	125	ILE	2.3
1	O	230	SER	2.3
1	O	325	GLU	2.3
1	O	339	GLY	2.3
1	X	332	GLU	2.3
1	O	285	ASN	2.3
1	X	410	ASP	2.2
1	X	227	HIS	2.2
1	X	415	LYS	2.2
1	X	500	LYS	2.2
1	O	468	GLY	2.2
1	O	120	GLY	2.2
1	X	472	THR	2.2
1	X	120	GLY	2.1
1	X	337	ALA	2.1
1	X	229	ARG	2.1
1	O	258	GLU	2.1
1	X	459	ASP	2.0
1	O	469	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	O	3126	4/4	0.87	0.12	57,58,61,61	0
3	EDO	X	3123	4/4	0.88	0.14	37,46,56,61	0
4	PO4	O	507	5/5	0.88	0.18	51,54,61,63	0
3	EDO	O	3127	4/4	0.91	0.10	46,47,55,55	0
3	EDO	O	3124	4/4	0.91	0.11	36,41,45,48	0
3	EDO	X	3125	4/4	0.93	0.13	61,61,62,64	0
4	PO4	X	507	5/5	0.94	0.21	50,52,63,63	0
2	GOL	X	1000	6/6	0.95	0.08	23,27,30,32	0
3	EDO	X	3122	4/4	0.96	0.16	29,32,35,36	0
2	GOL	O	1001	6/6	0.97	0.10	27,27,27,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.