



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 09:29 AM BST

PDB ID : 4H55  
Title : Crystal structure of Canavalia brasiliensis seed lectin (ConBr) in complex with beta-d-ribofuranose  
Authors : Salviano, E.; Rocha, B.A.M.; Cavada, B.S.; Delatorre, P.; Santi-Gadelha, T.; Gadelha, C.A.A.; Silva-Filho, J.C.; Nobrega, R.B.; Farias, D.L.  
Deposited on : 2012-09-18  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	<div> <div>6%</div> <div>65%</div> <div>32%</div> <div>••</div> </div>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 1874 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Concanavalin-Br.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	233	Total	C	N	O	S	0	0	0
			1775	1122	299	352	2			


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		

- Molecule 4 is beta-D-ribofuranose (three-letter code: BDR) (formula: C<sub>5</sub>H<sub>10</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	5	5		

- Molecule 5 is D-ALPHA-AMINOBUTYRIC ACID (three-letter code: DBB) (formula:  $C_4H_9NO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			7	4	1	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	80	Total	O	0	0
			80	80		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Concanavalin-Br



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.24Å 70.43Å 97.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.39 – 2.15 24.39 – 2.07	Depositor EDS
% Data completeness (in resolution range)	87.8 (24.39-2.15) 90.3 (24.39-2.07)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.45 (at 2.06Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.183 , 0.227 0.197 , 0.237	Depositor DCC
$R_{free}$ test set	610 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.8	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 44.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.40$ , $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.035 for -1/2*h+1/2*k+1/2*l,1/2*h-1/2*k +1/2*l,h+k 0.055 for -1/2*h+1/2*k-1/2*l,1/2*h-1/2*k- 1/2*l,-h-k 0.025 for k,h,-l 0.055 for -1/2*h-1/2*k+1/2*l,-1/2*h-1/2*k- 1/2*l,h-k 0.064 for -1/2*h-1/2*k-1/2*l,-1/2*h-1/2*k+ 1/2*l,-h+k	Xtriage
Reported twinning fraction	0.939 for H, K, L 0.061 for K, H, -L	Depositor
Outliers	0 of 13226 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	1874	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DBB, CA, MN, BDR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.00	1/1815 (0.1%)	0.99	2/2469 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	192	GLU	CG-CD	5.65	1.60	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	82	ASP	CB-CG-OD2	-5.41	113.43	118.30
1	A	33	ARG	NE-CZ-NH2	-5.33	117.63	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1775	0	1735	54	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	10	0	7	4	0
5	A	7	0	8	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	80	0	0	1	0
All	All	1874	0	1750	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:HIS:H	5:A:304:DBB:H	1.10	0.93
1:A:205:HIS:HD2	1:A:206:PRO:O	1.66	0.78
1:A:202:PRO:HD2	1:A:203:ASP:H	1.53	0.74
1:A:202:PRO:CD	1:A:203:ASP:H	2.10	0.65
1:A:98:GLY:HA3	4:A:303:BDR:O4	1.97	0.64
1:A:174:LEU:HD12	1:A:174:LEU:N	2.13	0.64
1:A:143:GLN:HE22	1:A:172:ARG:HH21	1.46	0.64
1:A:208:ASP:OD2	1:A:227:GLY:HA2	1.99	0.62
1:A:132:GLN:NE2	6:A:452:HOH:O	2.34	0.60
1:A:143:GLN:NE2	1:A:172:ARG:HE	1.99	0.60
1:A:92:GLY:HA2	1:A:109:TRP:CH2	2.36	0.59
1:A:181:ILE:HD12	1:A:182:TRP:N	2.18	0.59
1:A:156:LEU:O	1:A:171:GLY:HA3	2.05	0.57
1:A:4:ILE:HD13	1:A:215:SER:HB3	1.88	0.54
1:A:100:TYR:HB3	1:A:205:HIS:O	2.08	0.54
1:A:12:TYR:HD2	1:A:205:HIS:CD2	2.26	0.53
1:A:51:HIS:O	1:A:63:ALA:HA	2.08	0.53
1:A:133:PHE:O	1:A:152:GLY:HA2	2.10	0.52
1:A:102:GLU:C	1:A:165:PRO:HB3	2.29	0.52
1:A:201:SER:OG	1:A:203:ASP:HB2	2.10	0.52
1:A:180:HIS:N	5:A:304:DBB:H	1.93	0.51
1:A:151:GLU:OE2	1:A:155:ARG:NH2	2.44	0.51
1:A:87:GLU:HG2	1:A:180:HIS:CD2	2.46	0.50
1:A:4:ILE:HB	1:A:233:PHE:CE2	2.47	0.50
1:A:115:LEU:O	1:A:123:THR:HA	2.13	0.48
1:A:202:PRO:CD	1:A:203:ASP:N	2.77	0.48
1:A:205:HIS:CD2	1:A:206:PRO:O	2.56	0.48
1:A:181:ILE:CD1	1:A:182:TRP:CD1	2.97	0.48
1:A:8:GLU:O	1:A:25:ILE:HA	2.14	0.47
1:A:126:LEU:HD22	1:A:179:VAL:HG22	1.97	0.47
1:A:48:GLY:O	1:A:196:THR:HA	2.15	0.47
1:A:102:GLU:HG2	1:A:207:ALA:O	2.16	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:HIS:O	5:A:304:DBB:O	2.33	0.46
1:A:107:LEU:HD12	1:A:107:LEU:N	2.32	0.45
1:A:181:ILE:CD1	1:A:182:TRP:HD1	2.30	0.45
1:A:67:TYR:HB3	1:A:68:PRO:HD2	1.99	0.45
1:A:111:PHE:O	1:A:127:HIS:HA	2.17	0.45
1:A:27:ILE:HG23	1:A:61:LEU:HD23	1.99	0.45
1:A:32:VAL:O	1:A:32:VAL:HG22	2.17	0.44
1:A:23:PRO:HB2	1:A:40:TRP:O	2.18	0.44
1:A:164:SER:HA	1:A:165:PRO:HD2	1.69	0.43
1:A:87:GLU:HG2	1:A:180:HIS:NE2	2.33	0.43
1:A:175:PHE:CE2	1:A:177:ALA:HB3	2.54	0.42
1:A:221:ILE:HA	1:A:222:PRO:HD3	1.92	0.42
1:A:12:TYR:HA	1:A:13:PRO:HD3	1.75	0.42
1:A:9:LEU:O	1:A:209:GLY:HA3	2.20	0.41
1:A:112:THR:O	1:A:191:PHE:HA	2.20	0.41
1:A:5:VAL:HG22	1:A:29:ILE:HD12	2.02	0.41
1:A:105:THR:O	1:A:197:PHE:HA	2.21	0.41
1:A:208:ASP:OD2	4:A:303:BDR:C1	2.68	0.41
1:A:175:PHE:HE2	1:A:177:ALA:HB3	1.86	0.41
1:A:174:LEU:CD1	1:A:174:LEU:N	2.84	0.41
1:A:100:TYR:O	4:A:303:BDR:O1	2.39	0.41
1:A:99:LEU:HB2	4:A:303:BDR:H51	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	229/237 (97%)	216 (94%)	12 (5%)	1 (0%)	34 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	202	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/202 (98%)	188 (95%)	10 (5%)	24	20

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	LYS
1	A	118	ASN
1	A	150	THR
1	A	155	ARG
1	A	159	VAL
1	A	161	SER
1	A	185	SER
1	A	203	ASP
1	A	217	ILE
1	A	223	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	143	GLN
1	A	205	HIS
1	A	237	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	DBB	A	304	-	3,6,6	0.69	0	1,7,7	0.50	0
4	BDR	A	303	-	10,10,10	1.53	1 (10%)	13,14,14	2.37	4 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DBB	A	304	-	-	2/2/6/6	-
4	BDR	A	303	-	-	0/2/18/18	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	303	BDR	C1-C2	-4.20	1.47	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	303	BDR	O4-C1-C2	-5.50	97.69	104.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	303	BDR	C1-C2-C3	-3.55	97.85	102.30
4	A	303	BDR	O3-C3-C4	-3.32	101.44	111.05
4	A	303	BDR	O5-C5-C4	-2.73	101.93	111.29

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	304	DBB	N-CA-CB-CG
5	A	304	DBB	C-CA-CB-CG

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	304	DBB	3	0
4	A	303	BDR	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	233/237 (98%)	0.24	14 (6%) 21 29	7, 21, 34, 44	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	ALA	4.7
1	A	159	VAL	4.6
1	A	162	ASN	3.6
1	A	118	ASN	3.2
1	A	163	GLY	3.0
1	A	230	LEU	2.8
1	A	236	ALA	2.4
1	A	223	SER	2.2
1	A	9	LEU	2.1
1	A	237	ASN	2.1
1	A	211	ALA	2.0
1	A	6	ALA	2.0
1	A	218	ASP	2.0
1	A	17	ILE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	DBB	A	304	7/7	0.88	0.19	23,25,26,29	1
4	BDR	A	303	10/10	0.89	0.25	34,35,39,41	0
2	CA	A	301	1/1	0.99	0.05	28,28,28,28	0
3	MN	A	302	1/1	1.00	0.05	21,21,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.