



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 12:48 pm BST

PDB ID : 4H58  
Title : BRAF in complex with compound 3  
Authors : Vasbinder, M.; Aquila, B.; Augustin, M.; Chueng, T.; Cook, D.; Drew, L.; Fauber, B.; Glossop, S.; Godin, R.; Grondine, M.; Hennessy, E.; Johannes, J.; Lee, S.; Lyne, P.; Moertl, M.; Omer, C.; Palakurthi, S.; Pontz, T.; Read, J.; Sha, L.; Shen, M.; Steinbacher, S.; Wang, H.; Wu, A.; Ye, M.; Bagal, B.  
Deposited on : 2012-09-18  
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>2%</div> <div> <div></div> <div>70%</div> <div>23%</div> <div>• •</div> </div> </div>
1	B	275	<div> <div>%</div> <div> <div></div> <div>68%</div> <div>23%</div> <div>• 5%</div> </div> </div>
1	C	275	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>16%</div> <div>••</div> </div> </div>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6432 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase B-raf.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	1	0
			2108	1351	366	378	13			
1	B	262	Total	C	N	O	S	0	1	0
			2102	1350	364	375	13			
1	C	272	Total	C	N	O	S	0	0	0
			2177	1396	381	387	13			

- Molecule 2 is N-(4-[(2-methoxyethyl)amino]methyl}phenyl)-6-(pyridin-4-yl)quinazolin-2-amine (three-letter code: 10Z) (formula: C<sub>23</sub>H<sub>23</sub>N<sub>5</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			29	23	5	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	O	0	0
			4	4		
4	B	9	Total	O	0	0
			9	9		
4	C	2	Total	O	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein kinase B-raf



- Molecule 1: Serine/threonine-protein kinase B-raf



- Molecule 1: Serine/threonine-protein kinase B-raf



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	204.74Å 204.74Å 152.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.25 – 3.10 47.23 – 3.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.25-3.10) 95.3 (47.23-3.10)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 3.12Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.222 , 0.272 0.225 , 0.272	Depositor DCC
$R_{free}$ test set	1412 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.8	Xtriage
Anisotropy	0.139	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 74.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6432	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 10Z, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.49	0/2153	0.65	0/2906
1	B	0.54	0/2148	0.67	1/2900 (0.0%)
1	C	0.47	0/2226	0.61	0/3005
All	All	0.50	0/6527	0.65	1/8811 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	703	ARG	NE-CZ-NH1	5.68	123.14	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	626	MET	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2108	0	2145	34	0
1	B	2102	0	2137	40	0
1	C	2177	0	2209	25	0
2	A	29	0	23	0	0
3	C	1	0	0	0	0
4	A	4	0	0	1	0
4	B	9	0	0	2	0
4	C	2	0	0	0	0
All	All	6432	0	6514	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:485:ASN:HD22	1:B:493:GLN:HE22	1.24	0.86
1:A:616:ILE:HG21	1:A:665:ILE:HG12	1.63	0.80
1:B:573:HIS:O	1:B:574:ARG:HB2	1.90	0.71
1:B:551:LYS:HZ1	1:B:587:LEU:HA	1.57	0.68
1:B:574:ARG:NE	1:B:598:THR:HG21	2.08	0.68
1:B:575:ASP:O	1:B:580:ASN:ND2	2.27	0.68
1:B:490:THR:OG1	1:B:492:GLN:NE2	2.28	0.65
1:B:551:LYS:NZ	1:B:587:LEU:HA	2.12	0.65
1:A:487:THR:O	1:A:489:PRO:HD3	1.98	0.64
1:A:649:MET:HG3	1:A:684:CYS:SG	2.42	0.60
1:A:658:ILE:HD12	1:A:673:LEU:HD11	1.83	0.60
1:A:508:ARG:HD3	1:C:515:PHE:O	2.03	0.58
1:B:448:ASP:HA	4:B:806:HOH:O	2.04	0.57
1:A:575:ASP:HB2	1:A:596:LEU:HD23	1.88	0.56
1:B:573:HIS:NE2	1:B:593:ASP:O	2.39	0.55
1:B:596:LEU:HB2	4:B:807:HOH:O	2.06	0.55
1:B:660:ASN:HD22	1:B:663:GLN:HB2	1.71	0.55
1:B:594:PHE:HB3	1:B:597:ALA:HB3	1.89	0.54
1:B:708:GLN:HE21	1:B:708:GLN:C	2.11	0.54
1:C:462:ILE:HG13	1:C:470:VAL:HG12	1.89	0.54
1:A:557:ARG:HG3	1:A:713:ILE:HG22	1.90	0.54
1:A:550:ILE:HD12	1:A:720:LEU:HD23	1.90	0.53
1:A:567:HIS:HD2	1:A:571:ILE:O	1.92	0.53

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:625:ARG:O	1:A:626:MET:HB2	2.09	0.52
1:A:452:PRO:HB2	1:A:455:GLN:HE21	1.73	0.52
1:B:540:LEU:HD23	1:B:647:GLU:HB3	1.91	0.52
1:A:630:ASN:N	1:A:631:PRO:CD	2.72	0.52
1:B:465:GLY:O	1:B:466:SER:C	2.48	0.51
1:B:547:PHE:HB2	1:B:552:LEU:HD21	1.93	0.51
1:C:658:ILE:CD1	1:C:673:LEU:HD11	2.40	0.51
1:B:664:ILE:O	1:B:665:ILE:C	2.49	0.51
1:C:581:ILE:HG23	1:C:589:VAL:HG13	1.93	0.51
1:A:485:ASN:O	1:A:523:GLN:NE2	2.45	0.49
1:C:603:TRP:HD1	1:C:617:LEU:HD11	1.77	0.49
1:C:482:LYS:O	1:C:525:ALA:HA	2.12	0.49
1:C:603:TRP:HB3	1:C:661:ARG:HD2	1.95	0.49
1:B:598:THR:C	1:C:490:THR:HG22	2.33	0.49
1:C:501:VAL:HG23	1:C:526:ILE:HD11	1.94	0.49
1:B:620:ALA:HA	1:B:639:TYR:CD2	2.48	0.49
1:B:484:LEU:HD22	1:B:497:PHE:HB2	1.94	0.49
1:A:507:THR:HB	1:A:512:ILE:HG21	1.93	0.49
1:C:532:GLU:O	1:C:585:GLU:N	2.45	0.49
1:B:584:HIS:CD2	1:B:585:GLU:HG3	2.48	0.48
1:C:504:LEU:HB3	1:C:515:PHE:HB2	1.94	0.48
1:A:619:MET:HE3	1:A:623:VAL:HG12	1.96	0.48
1:B:695:CYS:O	1:B:703:ARG:HD2	2.13	0.48
1:A:596:LEU:O	1:A:597:ALA:HB2	2.14	0.48
1:C:511:ASN:HA	1:C:589:VAL:O	2.14	0.47
1:B:630:ASN:N	1:B:631:PRO:CD	2.78	0.47
1:B:554:ASP:O	1:B:558:GLN:HG3	2.15	0.47
1:A:630:ASN:CG	1:A:630:ASN:O	2.52	0.46
1:A:616:ILE:HD11	1:A:661:ARG:HD2	1.96	0.46
1:B:511:ASN:O	1:B:512:ILE:HD13	2.16	0.46
1:A:547:PHE:HB2	1:A:552:LEU:HD21	1.97	0.46
1:C:660:ASN:O	1:C:664:ILE:HG13	2.15	0.46
1:A:662:ASP:HA	1:A:665:ILE:HD12	1.98	0.46
1:B:646:TYR:CE1	1:B:650:THR:HG21	2.51	0.46
1:B:625:ARG:NE	1:B:669:GLY:O	2.49	0.45
1:C:481:VAL:HG13	1:C:527:VAL:HG13	1.98	0.45
1:A:625:ARG:O	1:A:626:MET:CB	2.65	0.45
1:A:572:ILE:HG22	1:A:574:ARG:HG3	1.98	0.45
1:B:567:HIS:O	1:B:570:SER:N	2.47	0.45
1:A:638:VAL:HG21	1:A:704:PRO:O	2.17	0.44
1:A:516:MET:HE3	1:C:509:HIS:HA	1.98	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:634:PHE:O	1:B:638:VAL:HG23	2.17	0.44
1:C:658:ILE:HD12	1:C:673:LEU:HD11	2.00	0.43
1:C:664:ILE:O	1:C:668:VAL:HG23	2.19	0.43
1:B:523:GLN:HA	1:B:523:GLN:HE21	1.83	0.43
1:A:664:ILE:O	1:A:668:VAL:HG23	2.19	0.43
1:B:529:GLN:HB2	1:B:529:GLN:HE21	1.68	0.43
1:B:548:GLU:O	1:B:551:LYS:N	2.52	0.43
1:C:465:GLY:HA3	1:C:470:VAL:HG23	2.00	0.43
1:B:574:ARG:HD2	1:B:596:LEU:O	2.19	0.43
1:A:706:PHE:N	1:A:707:PRO:HD2	2.34	0.43
1:C:499:ASN:O	1:C:503:VAL:HG23	2.19	0.43
1:A:574:ARG:HG2	1:A:632:TYR:CD1	2.54	0.42
1:A:570:SER:O	1:A:599:VAL:HG13	2.19	0.42
1:A:549:MET:HB2	1:A:683:ASN:HD21	1.84	0.42
1:B:539:HIS:O	1:B:545:THR:N	2.53	0.42
1:B:514:LEU:HB3	1:B:529:GLN:HG2	2.01	0.42
1:C:620:ALA:HA	1:C:639:TYR:CE1	2.55	0.42
1:C:617:LEU:HD12	1:C:617:LEU:H	1.84	0.42
1:B:614:GLY:H	1:B:661:ARG:HH22	1.69	0.41
1:C:542:ILE:CD1	1:C:599:VAL:HG21	2.49	0.41
1:C:542:ILE:HD11	1:C:599:VAL:HG21	2.02	0.41
1:C:662:ASP:HA	1:C:665:ILE:HD12	2.01	0.41
1:A:574:ARG:HD3	1:A:598:THR:HG22	2.03	0.41
1:B:574:ARG:HE	1:B:598:THR:HG21	1.83	0.41
1:A:657:ASN:N	1:A:657:ASN:OD1	2.54	0.41
1:B:586:ASP:O	1:B:587:LEU:HD12	2.21	0.41
1:A:499:ASN:O	1:A:503:VAL:HG23	2.21	0.41
1:A:640:ALA:O	1:A:644:VAL:HG23	2.21	0.40
1:B:614:GLY:N	1:B:661:ARG:HH22	2.19	0.40
1:B:687:ALA:HB3	1:B:716:LEU:HD22	2.03	0.40
1:A:616:ILE:HD12	4:A:902:HOH:O	2.22	0.40
1:C:470:VAL:HA	1:C:481:VAL:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/275 (94%)	229 (88%)	23 (9%)	8 (3%)	4	23
1	B	259/275 (94%)	229 (88%)	25 (10%)	5 (2%)	8	33
1	C	268/275 (98%)	240 (90%)	26 (10%)	2 (1%)	22	57
All	All	787/825 (95%)	698 (89%)	74 (9%)	15 (2%)	8	33

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	626	MET
1	A	628	ASP
1	B	466	SER
1	A	489	PRO
1	A	586	ASP
1	A	491	PRO
1	A	597	ALA
1	B	549	MET
1	B	628	ASP
1	C	611	GLN
1	A	577	LYS
1	B	665	ILE
1	C	575	ASP
1	A	595	GLY
1	B	630	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/242 (96%)	214 (92%)	18 (8%)	12	40
1	B	231/242 (96%)	210 (91%)	21 (9%)	9	33
1	C	239/242 (99%)	227 (95%)	12 (5%)	24	57

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	702/726 (97%)	651 (93%)	51 (7%)	14	43

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	467	PHE
1	A	486	VAL
1	A	519	SER
1	A	532	GLU
1	A	534	SER
1	A	570	SER
1	A	583	LEU
1	A	596	LEU
1	A	615	SER
1	A	616	ILE
1	A	630	ASN
1	A	649	MET
1	A	673	LEU
1	A	681	ARG
1	A	682	SER
1	A	686	LYS
1	A	698	LYS
1	A	710	LEU
1	B	469	THR
1	B	487	THR
1	B	490	THR
1	B	492	GLN
1	B	523	GLN
1	B	524	LEU
1	B	529	GLN
1	B	532	GLU
1	B	542	ILE
1	B	572	ILE
1	B	588	THR
1	B	593	ASP
1	B	596	LEU
1	B	598	THR
1	B	627	GLN
1	B	630	ASN
1	B	649	MET
1	B	655	TYR
1	B	681	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	698	LYS
1	B	708	GLN
1	C	464	SER
1	C	469	THR
1	C	471	TYR
1	C	527	VAL
1	C	535	SER
1	C	544	GLU
1	C	583	LEU
1	C	673	LEU
1	C	681	ARG
1	C	682	SER
1	C	715	LEU
1	C	720	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	455	GLN
1	A	485	ASN
1	A	523	GLN
1	A	627	GLN
1	B	455	GLN
1	B	492	GLN
1	B	493	GLN
1	B	511	ASN
1	B	529	GLN
1	B	657	ASN
1	B	660	ASN
1	B	663	GLN
1	B	708	GLN
1	C	455	GLN
1	C	485	ASN
1	C	492	GLN
1	C	567	HIS
1	C	652	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	10Z	A	801	-	32,32,32	1.39	3 (9%)	41,42,42	1.96	8 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	10Z	A	801	-	-	2/14/14/14	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	10Z	C9-N7	3.53	1.39	1.32
2	A	801	10Z	C8-N7	3.20	1.39	1.34
2	A	801	10Z	C8-N17	3.18	1.43	1.36

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	10Z	N7-C8-N10	-9.21	117.62	126.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	10Z	C9-N7-C8	4.01	121.82	115.88
2	A	801	10Z	C8-N10-C12	3.08	120.62	115.60
2	A	801	10Z	C1-C3-N5	-2.28	119.65	123.62
2	A	801	10Z	C23-N17-C8	2.12	135.38	129.23
2	A	801	10Z	C16-C11-C12	2.07	120.98	118.27
2	A	801	10Z	C21-C23-N17	-2.06	113.75	120.64
2	A	801	10Z	C22-C23-N17	2.02	127.40	120.64

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	10Z	N25-C26-C27-O28
2	A	801	10Z	C27-C26-N25-C24

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	263/275 (95%)	-0.09	6 (2%)	60 39	51, 71, 109, 117	18 (6%)
1	B	262/275 (95%)	-0.09	3 (1%)	80 64	31, 59, 81, 95	13 (4%)
1	C	272/275 (98%)	0.11	4 (1%)	73 54	52, 77, 107, 111	23 (8%)
All	All	797/825 (96%)	-0.02	13 (1%)	72 51	31, 69, 104, 117	54 (6%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	519	SER	4.1
1	A	467	PHE	4.1
1	A	487	THR	3.4
1	B	627	GLN	3.4
1	C	463	GLY	2.7
1	C	471	TYR	2.5
1	C	597	ALA	2.5
1	B	628	ASP	2.4
1	B	721	PRO	2.4
1	A	447	ASP	2.1
1	A	488	ALA	2.0
1	A	484	LEU	2.0
1	A	489	PRO	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CL	C	801	1/1	0.84	0.17	72,72,72,72	0
2	10Z	A	801	29/29	0.90	0.26	87,88,97,98	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.