



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 03:01 am BST

PDB ID : 6H5A  
Title : Crystal structure of Mycobacterium tuberculosis phosphatidylinositol phosphate synthase (PgsA1) in complex with manganese and citrate  
Authors : Grave, K.; Hogbom, M.  
Deposited on : 2018-07-24  
Resolution : 1.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>7%</div> <div>7%</div> </div> </div>
1	B	223	<div> <div>6%</div> <div> <div></div> <div>86%</div> <div>8%</div> <div>6%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SO4	B	306	-	-	X	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6963 atoms, of which 3461 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CDP-diacylglycerol--inositol 3-phosphatidyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	207	Total	C	H	N	O	S	0	5	0
			3212	1028	1632	274	267	11			
1	B	209	Total	C	H	N	O	S	0	0	0
			3200	1027	1620	271	271	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	218	GLU	-	expression tag	UNP P9WPG7
A	219	ASN	-	expression tag	UNP P9WPG7
A	220	LEU	-	expression tag	UNP P9WPG7
A	221	TYR	-	expression tag	UNP P9WPG7
A	222	PHE	-	expression tag	UNP P9WPG7
A	223	GLN	-	expression tag	UNP P9WPG7
B	218	GLU	-	expression tag	UNP P9WPG7
B	219	ASN	-	expression tag	UNP P9WPG7
B	220	LEU	-	expression tag	UNP P9WPG7
B	221	TYR	-	expression tag	UNP P9WPG7
B	222	PHE	-	expression tag	UNP P9WPG7
B	223	GLN	-	expression tag	UNP P9WPG7

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Mn	0	0
			3	3		
2	A	4	Total	Mn	0	0
			4	4		

- Molecule 3 is CITRATE ANION (three-letter code: FLC) (formula: C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	0	0	
			13	6	7			
3	A	1	Total	C	O	0	0	
			13	6	7			
3	B	1	Total	C	H	O	0	0
			18	6	5	7		

- Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: C<sub>21</sub>H<sub>40</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 25 21 4	0	0
4	A	1	Total C H O 65 21 40 4	0	0
4	B	1	Total C O 25 21 4	0	0

- Molecule 5 is EICOSANE (three-letter code: LFA) (formula:  $C_{20}H_{42}$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C H 33 11 22	0	0
5	A	1	Total C H 62 20 42	0	0
5	B	1	Total C H 62 20 42	0	0
5	B	1	Total C H 21 7 14	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C<sub>10</sub>H<sub>22</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	H	O	0	0
			38	10	22	6		
7	B	1	Total	C	H	O	0	0
			38	10	22	6		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	70	Total 70	O 70	0	0
8	B	56	Total 56	O 56	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CDP-diacylglycerol--inositol 3-phosphatidyltransferase



- Molecule 1: CDP-diacylglycerol--inositol 3-phosphatidyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.07Å 77.94Å 100.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.70 – 1.88 45.70 – 1.88	Depositor EDS
% Data completeness (in resolution range)	98.9 (45.70-1.88) 98.8 (45.70-1.88)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.28 (at 1.88Å)	Xtriage
Refinement program	PHENIX 1.12 _2829	Depositor
R, $R_{free}$	0.204 , 0.230 0.204 , 0.229	Depositor DCC
$R_{free}$ test set	2196 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.4	Xtriage
Anisotropy	0.253	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 46.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6963	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.03 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.2213e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OLC, LFA, MN, 1PE, SO4, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/1643	0.61	1/2238 (0.0%)
1	B	0.45	0/1618	0.56	0/2205
All	All	0.47	0/3261	0.59	1/4443 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	93	ASP	CB-CG-OD1	7.98	125.48	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1580	1632	1605	12	0
1	B	1580	1620	1620	11	0
2	A	4	0	0	0	0
2	B	3	0	0	0	0
3	A	26	0	8	0	0
3	B	13	5	4	0	0
4	A	50	40	80	5	0
4	B	25	0	40	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	31	64	63	0	0
5	B	27	56	52	0	0
6	B	5	0	0	3	0
7	B	32	44	43	0	0
8	A	70	0	0	1	0
8	B	56	0	0	1	0
All	All	3502	3461	3515	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:306:SO4:O2	8:B:401:HOH:O	1.97	0.82
4:B:305:OLC:H12	4:B:305:OLC:H7A	1.73	0.70
1:B:217:ARG:NE	6:B:306:SO4:O2	2.23	0.68
1:B:186:VAL:HG21	4:B:305:OLC:H8A	1.78	0.66
1:A:157:ILE:HD11	4:A:308:OLC:H22	1.82	0.60

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/223 (94%)	206 (98%)	4 (2%)	0	100	100
1	B	207/223 (93%)	203 (98%)	4 (2%)	0	100	100
All	All	417/446 (94%)	409 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/175 (94%)	165 (100%)	0	100	100
1	B	164/175 (94%)	162 (99%)	2 (1%)	71	67
All	All	329/350 (94%)	327 (99%)	2 (1%)	86	86

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	71	ASP
1	B	219	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

Of 20 ligands modelled in this entry, 7 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	OLC	A	308	-	24,24,24	0.73	1 (4%)	25,25,25	1.15	2 (8%)
7	1PE	B	308	-	15,15,15	0.59	0	14,14,14	0.45	0
5	LFA	B	310	-	6,6,19	0.18	0	5,5,18	0.41	0
5	LFA	B	307	-	19,19,19	0.17	0	18,18,18	1.04	0
7	1PE	B	309	-	15,15,15	0.68	0	14,14,14	0.43	0
5	LFA	A	309	-	10,10,19	0.15	0	9,9,18	0.79	0
3	FLC	A	305	2	3,12,12	2.37	2 (66%)	3,17,17	6.13	3 (100%)
4	OLC	B	305	-	24,24,24	0.73	1 (4%)	25,25,25	1.21	1 (4%)
3	FLC	B	304	2	3,12,12	1.69	1 (33%)	3,17,17	2.62	2 (66%)
5	LFA	A	310	-	19,19,19	0.19	0	18,18,18	0.73	0
6	SO4	B	306	-	4,4,4	0.26	0	6,6,6	0.25	0
3	FLC	A	306	2	3,12,12	2.22	1 (33%)	3,17,17	3.77	3 (100%)
4	OLC	A	307	-	24,24,24	0.67	1 (4%)	25,25,25	1.33	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLC	A	308	-	-	9/24/24/24	-
7	1PE	B	308	-	-	4/13/13/13	-
5	LFA	B	310	-	-	2/4/4/17	-
5	LFA	B	307	-	-	6/17/17/17	-
7	1PE	B	309	-	-	2/13/13/13	-
5	LFA	A	309	-	-	4/8/8/17	-
3	FLC	A	305	2	-	2/6/16/16	-
4	OLC	B	305	-	-	9/24/24/24	-
3	FLC	B	304	2	-	0/6/16/16	-
5	LFA	A	310	-	-	7/17/17/17	-
3	FLC	A	306	2	-	1/6/16/16	-
4	OLC	A	307	-	-	9/24/24/24	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	305	FLC	CA-CB	-3.43	1.50	1.54
3	A	306	FLC	OHB-CB	-3.15	1.38	1.43
4	B	305	OLC	O20-C1	3.07	1.42	1.33
4	A	308	OLC	O20-C1	2.94	1.41	1.33
4	A	307	OLC	O20-C1	2.85	1.41	1.33

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	305	FLC	CG-CB-CA	-8.99	85.27	109.33
3	A	306	FLC	CG-CB-CA	-5.15	95.54	109.33
3	A	305	FLC	CB-CA-CAC	-4.31	108.08	114.98
4	B	305	OLC	O20-C1-C2	4.01	124.50	111.91
4	A	307	OLC	O20-C1-C2	3.81	123.85	111.91

There are no chirality outliers.

5 of 55 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	305	FLC	CAC-CA-CB-OHB
4	B	305	OLC	O20-C21-C22-C24
4	A	307	OLC	C21-C22-C24-O25
4	A	308	OLC	C2-C1-O20-C21
4	A	308	OLC	O19-C1-O20-C21


There are no ring outliers.

4 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	308	OLC	3	0
4	B	305	OLC	6	0
6	B	306	SO4	3	0
4	A	307	OLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	207/223 (92%)	0.32	11 (5%) 26 28	18, 26, 43, 58	0
1	B	209/223 (93%)	0.45	13 (6%) 20 22	19, 28, 48, 58	1 (0%)
All	All	416/446 (93%)	0.39	24 (5%) 23 24	18, 27, 47, 58	1 (0%)

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	216	ASP	6.7
1	A	170	PHE	4.8
1	B	210	ALA	4.7
1	B	171	VAL	4.7
1	A	6	PHE	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	LFA	A	310	20/20	0.66	0.20	35,48,56,57	0
2	MN	B	303	1/1	0.70	0.13	47,47,47,47	1
4	OLC	A	307	25/25	0.77	0.20	25,36,48,51	0
4	OLC	A	308	25/25	0.80	0.20	33,51,81,84	0
4	OLC	B	305	25/25	0.81	0.20	25,37,48,52	0
5	LFA	B	310	7/20	0.82	0.12	35,43,46,50	0
7	1PE	B	309	16/16	0.84	0.24	33,47,62,67	0
5	LFA	B	307	20/20	0.86	0.22	25,37,49,56	0
5	LFA	A	309	11/20	0.86	0.19	34,49,64,67	0
3	FLC	A	305	13/13	0.87	0.13	23,25,37,37	0
2	MN	B	302	1/1	0.88	0.07	47,47,47,47	1
7	1PE	B	308	16/16	0.89	0.18	31,47,61,67	0
3	FLC	B	304	13/13	0.91	0.16	28,34,42,42	18
3	FLC	A	306	13/13	0.91	0.12	20,25,28,28	0
2	MN	A	303	1/1	0.93	0.11	25,25,25,25	0
2	MN	A	304	1/1	0.94	0.08	24,24,24,24	0
6	SO4	B	306	5/5	0.95	0.46	29,30,41,45	5
2	MN	A	302	1/1	0.98	0.11	27,27,27,27	0
2	MN	B	301	1/1	0.99	0.07	37,37,37,37	1
2	MN	A	301	1/1	1.00	0.11	22,22,22,22	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around LFA A 310:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around OLC A 307:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around OLC A 308:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around OLC B 305:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around LFA B 310:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around LFA B 307:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around LFA A 309:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.