



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 08:11 am BST

PDB ID : 3H7N  
Title : Structure of Nup120  
Authors : Seo, H.S.; Ma, Y.; Debler, E.W.; Blobel, G.; Hoelz, A.  
Deposited on : 2009-04-27  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	729	<div> <div>11%</div> <div> <div>31%</div> <div>53%</div> <div>12%</div> </div> </div>
1	B	729	<div> <div>9%</div> <div> <div>31%</div> <div>53%</div> <div>12%</div> </div> </div>
1	C	729	<div> <div>10%</div> <div> <div>31%</div> <div>54%</div> <div>12%</div> </div> </div>
1	D	729	<div> <div>9%</div> <div> <div>32%</div> <div>53%</div> <div>11%</div> </div> </div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 22992 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nucleoporin NUP120.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	706	Total	C	N	O	S	0	0	0
			5748	3718	913	1099	18			
1	B	706	Total	C	N	O	S	0	0	0
			5748	3718	913	1099	18			
1	C	706	Total	C	N	O	S	0	0	0
			5748	3718	913	1099	18			
1	D	706	Total	C	N	O	S	0	0	0
			5748	3718	913	1099	18			






• Molecule 1: Nucleoporin NUP120





• Molecule 1: Nucleoporin NUP120





● Molecule 1: Nucleoporin NUP120





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.99Å 115.72Å 156.08Å 90.06° 89.96° 90.02°	Depositor
Resolution (Å)	50.00 – 3.00 47.47 – 3.01	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-3.00) 85.3 (47.47-3.01)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.58 (at 3.01Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.255 , 0.274 0.254 , 0.272	Depositor DCC
$R_{free}$ test set	3365 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	93.7	Xtriage
Anisotropy	0.330	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 67.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.448 for h,-k,-l 0.449 for -h,k,-l 0.447 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	22992	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	123.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	2/5878 (0.0%)	0.71	3/7981 (0.0%)
1	B	0.46	0/5878	0.71	2/7981 (0.0%)
1	C	0.46	0/5878	0.71	2/7981 (0.0%)
1	D	0.46	0/5878	0.71	3/7981 (0.0%)
All	All	0.47	2/23512 (0.0%)	0.71	10/31924 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	307	SER	C-O	5.94	1.34	1.23
1	A	306	ASP	CB-CG	5.73	1.63	1.51

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	358	GLY	N-CA-C	-5.81	98.58	113.10
1	D	358	GLY	N-CA-C	-5.73	98.78	113.10
1	C	358	GLY	N-CA-C	-5.72	98.81	113.10
1	A	307	SER	CB-CA-C	5.71	120.95	110.10
1	A	358	GLY	N-CA-C	-5.66	98.95	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5748	0	5657	593	0
1	B	5748	0	5657	564	0
1	C	5748	0	5657	591	0
1	D	5748	0	5657	567	0
All	All	22992	0	22628	2281	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 50.

The worst 5 of 2281 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:224:ILE:HG12	1:B:273:VAL:HG11	1.23	1.20
1:D:224:ILE:HG12	1:D:273:VAL:HG11	1.23	1.15
1:A:224:ILE:HG12	1:A:273:VAL:HG11	1.24	1.14
1:C:224:ILE:HG12	1:C:273:VAL:HG11	1.22	1.14
1:D:398:VAL:HB	1:D:669:ARG:NH1	1.62	1.14

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	702/729 (96%)	463 (66%)	169 (24%)	70 (10%)	0	2
1	B	702/729 (96%)	469 (67%)	159 (23%)	74 (10%)	0	2
1	C	702/729 (96%)	467 (66%)	165 (24%)	70 (10%)	0	2
1	D	702/729 (96%)	469 (67%)	165 (24%)	68 (10%)	0	2
All	All	2808/2916 (96%)	1868 (66%)	658 (23%)	282 (10%)	0	2

5 of 282 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	100	ALA
1	A	104	GLN
1	A	113	VAL
1	A	214	ARG
1	A	241	CYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	661/682 (97%)	608 (92%)	53 (8%)	12	40
1	B	661/682 (97%)	607 (92%)	54 (8%)	11	39
1	C	661/682 (97%)	609 (92%)	52 (8%)	12	41
1	D	661/682 (97%)	610 (92%)	51 (8%)	13	42
All	All	2644/2728 (97%)	2434 (92%)	210 (8%)	12	41

5 of 210 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	565	THR
1	C	270	PHE
1	D	541	PRO
1	B	596	PHE
1	B	717	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 81 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	575	ASN
1	C	132	GLN
1	D	487	ASN
1	B	589	ASN
1	C	22	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	706/729 (96%)	0.50	79 (11%) <b>5</b> <b>1</b>	51, 121, 174, 188	0
1	B	706/729 (96%)	0.51	69 (9%) <b>7</b> <b>2</b>	54, 121, 174, 188	0
1	C	706/729 (96%)	0.54	74 (10%) <b>6</b> <b>2</b>	52, 122, 174, 191	0
1	D	706/729 (96%)	0.48	68 (9%) <b>8</b> <b>2</b>	50, 121, 174, 186	0
All	All	2824/2916 (96%)	0.51	290 (10%) <b>6</b> <b>2</b>	50, 122, 174, 191	0

The worst 5 of 290 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	600	ASP	9.5
1	C	174	SER	9.3
1	B	244	LYS	9.3
1	C	186	LEU	8.5
1	B	194	TYR	7.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.