



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 10:36 am BST

PDB ID : 2H8M
Title : N-Domain Of Grp94 In Complex With the 2-Iodo-NECA
Authors : Immormino, R.M.; Gewirth, D.T.
Deposited on : 2006-06-07
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	<div> <div>8%</div> <div> <div></div> <div>78%</div> <div>12%</div> <div>9%</div> </div> </div>
1	B	236	<div> <div>7%</div> <div> <div></div> <div>76%</div> <div>13%</div> <div>11%</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3959 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Endoplasmin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	215	Total	C	N	O	S	0	5	0
			1698	1082	282	326	8			
1	B	210	Total	C	N	O	S	0	5	0
			1672	1063	277	325	7			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	65	GLY	-	CLONING ARTIFACT	UNP P41148
A	66	SER	-	CLONING ARTIFACT	UNP P41148
A	67	HIS	-	CLONING ARTIFACT	UNP P41148
A	68	MET	-	CLONING ARTIFACT	UNP P41148
A	287	GLY	-	SEE REMARK 999	UNP P41148
A	288	GLY	-	SEE REMARK 999	UNP P41148
A	289	GLY	-	SEE REMARK 999	UNP P41148
A	290	GLY	-	SEE REMARK 999	UNP P41148
B	65	GLY	-	CLONING ARTIFACT	UNP P41148
B	66	SER	-	CLONING ARTIFACT	UNP P41148
B	67	HIS	-	CLONING ARTIFACT	UNP P41148
B	68	MET	-	CLONING ARTIFACT	UNP P41148
B	287	GLY	-	SEE REMARK 999	UNP P41148
B	288	GLY	-	SEE REMARK 999	UNP P41148
B	289	GLY	-	SEE REMARK 999	UNP P41148
B	290	GLY	-	SEE REMARK 999	UNP P41148

- Molecule 2 is (2S,3S,4R,5R)-5-(6-AMINO-2-iodo-9H-purin-9-yl)-N-ethyl-3,4-dihydroxytetrahydrofuran-2-carboxamide (three-letter code: NEI) (formula: C₁₂H₁₅IN₆O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	I	N	O	2	1
			23	12	1	6	4		
2	B	1	Total	C	I	N	O	2	1
			23	12	1	6	4		

- Molecule 3 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	8	5		
3	A	1	Total	C	O	0	0
			13	8	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	6	1
			13	8	5		
3	A	1	Total	C	O	0	0
			13	8	5		
3	A	1	Total	C	O	0	0
			13	8	5		
3	B	1	Total	C	O	0	0
			13	8	5		
3	B	1	Total	C	O	0	0
			13	8	5		
3	B	1	Total	C	O	0	1
			13	8	5		
3	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 4 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C₁₀H₂₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			16	10	6		
4	B	1	Total	C	O	0	0
			16	10	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	206	Total 206	O 206	0	0
5	B	188	Total 188	O 188	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Endoplasmic



• Molecule 1: Endoplasmic



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.32Å 84.50Å 96.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.51 – 1.80 45.51 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.51-1.80) 99.9 (45.51-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 1.79Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.213 , 0.241 0.208 , 0.236	Depositor DCC
R_{free} test set	5010 reflections (10.04%)	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtriage
Anisotropy	0.657	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 59.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3959	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.08 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.8884e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, NEI, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/1721	0.60	0/2324
1	B	0.31	0/1695	0.59	0/2287
All	All	0.31	0/3416	0.59	0/4611

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1698	0	1669	31	0
1	B	1672	0	1642	29	0
2	A	23	0	15	3	0
2	B	23	0	15	3	0
3	A	65	0	90	3	0
3	B	52	0	72	1	0
4	B	32	0	44	2	0
5	A	206	0	0	5	1
5	B	188	0	0	5	1
All	All	3959	0	3547	61	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 9.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:VAL:HG13	2:A:401[A]:NEI:I2	2.56	0.75
1:B:152:VAL:HG13	2:B:402[A]:NEI:I2	2.58	0.73
1:A:93:LEU:HD11	1:A:192:ILE:HD12	1.75	0.68
1:A:152:VAL:CG1	2:A:401[A]:NEI:I2	3.14	0.64
1:B:192:ILE:HG23	1:B:197:VAL:HG23	1.80	0.64
1:B:76:PHE:HA	4:B:602:1PE:H252	1.79	0.63
1:B:152:VAL:CG1	2:B:402[A]:NEI:I2	3.18	0.61
1:B:93:LEU:HD11	1:B:192:ILE:HD12	1.83	0.60
1:A:93:LEU:HD11	1:A:192:ILE:CD1	2.32	0.59
1:B:138:CYS:SG	1:B:264:ILE:HG13	2.44	0.57
1:A:117:LEU:CD1	3:A:507:PG4:H61	2.36	0.56
1:A:117:LEU:HD12	3:A:507:PG4:H61	1.87	0.55
1:A:85[A]:MET:SD	1:A:197:VAL:HG11	2.48	0.54
1:A:99[B]:ILE:HD12	1:A:99[B]:ILE:C	2.29	0.53
1:A:208:LYS:HE2	1:A:224:GLU:OE2	2.09	0.53
1:A:137[A]:LYS:HD2	5:A:586:HOH:O	2.08	0.53
1:B:90:ILE:HD12	4:B:601:1PE:H122	1.92	0.51
1:B:137[A]:LYS:HD2	5:B:678:HOH:O	2.10	0.51
1:B:272:SER:CB	1:B:279:ILE:HD12	2.43	0.48
1:A:93:LEU:HD13	1:A:99[B]:ILE:CD1	2.44	0.48
1:B:175:LEU:HD23	1:B:175:LEU:O	2.14	0.48
1:A:188:THR:O	1:A:192:ILE:HG13	2.14	0.47
1:B:272:SER:HB3	1:B:279:ILE:HD12	1.97	0.47
1:A:175:LEU:HD23	1:A:175:LEU:O	2.14	0.47
1:B:93:LEU:HD11	1:B:192:ILE:CD1	2.44	0.47
1:A:192:ILE:HG23	1:A:197:VAL:HG23	1.97	0.47
1:A:93:LEU:HD13	1:A:99[B]:ILE:HD11	1.97	0.47
1:B:264:ILE:O	1:B:268:VAL:HG23	2.15	0.46
1:B:254:GLU:HG3	5:B:719:HOH:O	2.15	0.46
1:A:272:SER:OG	1:A:279:ILE:HD12	2.15	0.45
1:A:122:ASP:HB3	1:A:125:ALA:HB2	1.99	0.45
1:A:272:SER:CB	1:A:279:ILE:HD12	2.46	0.45
1:B:93:LEU:HB2	1:B:204:LEU:HD11	1.98	0.45
1:B:140:LYS:HG2	1:B:259:LEU:HB3	1.98	0.45
1:A:99[B]:ILE:HD12	1:A:100:PHE:N	2.33	0.44
1:A:98:GLU:HG2	5:A:599:HOH:O	2.18	0.44
1:B:153:GLY:O	2:B:402[A]:NEI:I2	3.05	0.44
1:A:188:THR:N	5:A:633:HOH:O	2.50	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:282:TRP:CZ3	1:B:332:ASP:HA	2.54	0.43
1:A:134:VAL:HB	1:A:279:ILE:HG12	2.01	0.43
1:B:135:LYS:HB3	1:B:333:TRP:CH2	2.54	0.43
1:A:210:ILE:HB	1:A:248:THR:HB	1.99	0.43
1:B:275:ILE:HG22	1:B:277:PHE:H	1.83	0.43
1:B:75:LYS:HE2	5:B:722:HOH:O	2.18	0.43
1:B:90:ILE:HA	1:B:204:LEU:HD21	2.00	0.43
1:B:88:LEU:HD11	1:B:175:LEU:HG	2.01	0.43
1:B:238:GLY:HA2	5:B:688:HOH:O	2.18	0.42
1:A:85[B]:MET:CE	1:A:174:PHE:HD2	2.32	0.42
1:B:88:LEU:HD13	1:B:178:MET:HB2	2.02	0.42
1:A:153:GLY:O	2:A:401[A]:NEI:I2	3.07	0.42
1:A:154[B]:MET:CE	1:A:162:ASN:HB2	2.50	0.42
1:A:129:ASN:OD1	1:A:131:GLU:HG2	2.19	0.42
1:B:188:THR:N	5:B:768:HOH:O	2.53	0.41
3:A:507:PG4:H82	5:A:594:HOH:O	2.19	0.41
1:B:135:LYS:HB3	1:B:333:TRP:CZ3	2.54	0.41
1:A:253:GLU:HB3	5:A:699:HOH:O	2.21	0.41
1:A:272:SER:HB3	1:A:279:ILE:HD12	2.03	0.41
1:B:237:ARG:HH22	3:B:501:PG4:H22	1.85	0.41
1:B:188:THR:O	1:B:192:ILE:HG13	2.21	0.40
1:A:85[B]:MET:HE1	1:A:174:PHE:HD2	1.86	0.40
1:A:282:TRP:CZ3	1:A:332:ASP:HA	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:569:HOH:O	5:B:670:HOH:O[3_146]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/236 (90%)	203 (96%)	8 (4%)	1 (0%)	29	15
1	B	206/236 (87%)	198 (96%)	8 (4%)	0	100	100
All	All	418/472 (89%)	401 (96%)	16 (4%)	1 (0%)	47	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	129	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/207 (88%)	183 (100%)	0	100	100
1	B	183/207 (88%)	183 (100%)	0	100	100
All	All	366/414 (88%)	366 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PG4	B	508[B]	-	12,12,12	0.50	0	11,11,11	0.91	0
4	1PE	B	602	-	15,15,15	0.59	0	14,14,14	0.92	0
3	PG4	A	504	-	12,12,12	0.50	0	11,11,11	0.80	0
3	PG4	B	509	-	12,12,12	0.52	0	11,11,11	0.77	0
2	NEI	A	401[A]	-	19,25,25	0.78	1 (5%)	21,37,37	1.83	6 (28%)
3	PG4	B	501	-	12,12,12	0.46	0	11,11,11	0.83	0
3	PG4	A	507	-	12,12,12	0.53	0	11,11,11	0.73	0
2	NEI	B	402[A]	-	19,25,25	0.79	1 (5%)	21,37,37	1.78	6 (28%)
3	PG4	A	510	-	12,12,12	0.54	0	11,11,11	0.78	0
3	PG4	B	502	-	12,12,12	0.53	0	11,11,11	0.77	0
3	PG4	A	506[B]	-	12,12,12	0.46	0	11,11,11	1.11	2 (18%)
3	PG4	A	505	-	12,12,12	0.54	0	11,11,11	0.78	0
4	1PE	B	601	-	15,15,15	0.51	0	14,14,14	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PG4	B	508[B]	-	-	2/10/10/10	-
4	1PE	B	602	-	-	3/13/13/13	-
3	PG4	A	504	-	-	0/10/10/10	-
3	PG4	B	509	-	-	0/10/10/10	-
2	NEI	A	401[A]	-	-	3/7/27/27	0/3/3/3
3	PG4	B	501	-	-	3/10/10/10	-
3	PG4	A	507	-	-	3/10/10/10	-
2	NEI	B	402[A]	-	-	2/7/27/27	0/3/3/3
3	PG4	A	510	-	-	0/10/10/10	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PG4	B	502	-	-	1/10/10/10	-
3	PG4	A	506[B]	-	-	2/10/10/10	-
3	PG4	A	505	-	-	0/10/10/10	-
4	1PE	B	601	-	-	3/13/13/13	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	402[A]	NEI	O4'-C1'	2.19	1.44	1.41
2	A	401[A]	NEI	O4'-C1'	2.17	1.44	1.41

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401[A]	NEI	C2-N3-C4	4.07	119.98	115.03
2	B	402[A]	NEI	C2-N3-C4	3.87	119.73	115.03
2	A	401[A]	NEI	C2'-C3'-C4'	-3.56	97.49	101.64
2	B	402[A]	NEI	C2'-C3'-C4'	-3.26	97.84	101.64
2	A	401[A]	NEI	N3-C2-N1	-2.88	121.37	126.77
2	B	402[A]	NEI	N3-C2-N1	-2.79	121.53	126.77
2	B	402[A]	NEI	C4-C5-N7	-2.77	106.52	109.40
2	A	401[A]	NEI	C4-C5-N7	-2.74	106.54	109.40
2	B	402[A]	NEI	C5-C6-N1	-2.70	119.24	121.01
2	A	401[A]	NEI	C5-C6-N1	-2.62	119.29	121.01
2	B	402[A]	NEI	C1'-N9-C4	-2.39	122.44	126.64
3	A	506[B]	PG4	C7-O4-C6	-2.20	103.76	113.29
3	A	506[B]	PG4	C5-O3-C4	-2.11	104.13	113.29
2	A	401[A]	NEI	C1'-N9-C4	-2.10	122.95	126.64

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	402[A]	NEI	C52-C51-N5'-C5'
3	A	506[B]	PG4	O4-C7-C8-O5
3	B	508[B]	PG4	O3-C5-C6-O4
3	A	506[B]	PG4	O2-C3-C4-O3
2	A	401[A]	NEI	C52-C51-N5'-C5'
4	B	602	1PE	OH7-C16-C26-OH6
4	B	601	1PE	OH2-C12-C22-OH3
3	B	501	PG4	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	502	PG4	O1-C1-C2-O2
4	B	602	1PE	OH5-C14-C24-OH4
4	B	602	1PE	OH6-C15-C25-OH5
3	A	507	PG4	O3-C5-C6-O4
3	A	507	PG4	O4-C7-C8-O5
3	A	507	PG4	O2-C3-C4-O3
2	A	401[A]	NEI	O4'-C4'-C5'-N5'
4	B	601	1PE	OH7-C16-C26-OH6
3	B	508[B]	PG4	O1-C1-C2-O2
4	B	601	1PE	OH4-C13-C23-OH3
2	A	401[A]	NEI	O4'-C4'-C5'-O5'
2	B	402[A]	NEI	O4'-C4'-C5'-N5'
3	B	501	PG4	O4-C7-C8-O5
3	B	501	PG4	O3-C5-C6-O4

There are no ring outliers.

6 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	602	1PE	1	0
2	A	401[A]	NEI	3	0
3	B	501	PG4	1	0
3	A	507	PG4	3	0
2	B	402[A]	NEI	3	0
4	B	601	1PE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/236 (91%)	0.36	19 (8%) 10 7	12, 23, 76, 95	0
1	B	210/236 (88%)	0.24	16 (7%) 13 10	12, 25, 53, 88	1 (0%)
All	All	425/472 (90%)	0.30	35 (8%) 11 9	12, 24, 60, 95	1 (0%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	286	THR	7.4
1	A	170	GLY	7.3
1	B	286	THR	7.2
1	B	175	LEU	6.8
1	A	175	LEU	6.6
1	B	195	PHE	5.7
1	B	99[A]	ILE	5.5
1	A	179	THR	5.4
1	B	174	PHE	5.0
1	A	328	LYS	4.8
1	A	171	THR	4.8
1	B	179	THR	4.8
1	A	330	VAL	4.7
1	A	174	PHE	4.2
1	A	195	PHE	4.1
1	A	182	GLN	3.9
1	A	329	THR	3.8
1	A	180	GLU	3.8
1	A	181	ALA	3.5
1	B	330	VAL	3.4
1	B	178	MET	3.4
1	B	328	LYS	3.4
1	A	191	LEU	3.3
1	B	329	THR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	176	ASN	3.1
1	A	172	SER	3.0
1	A	166	ILE	2.6
1	B	188	THR	2.6
1	B	85[A]	MET	2.5
1	A	124	ASN	2.4
1	A	192	ILE	2.3
1	A	173	GLU	2.2
1	B	274	PHE	2.2
1	B	166	ILE	2.1
1	B	180	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PG4	A	507	13/13	0.65	0.27	65,65,66,66	0
3	PG4	A	510	13/13	0.68	0.25	47,53,55,55	0
4	1PE	B	602	16/16	0.72	0.26	44,46,49,52	0
3	PG4	B	502	13/13	0.73	0.24	64,65,66,66	0
3	PG4	A	505	13/13	0.75	0.19	52,54,58,60	0
3	PG4	B	509	13/13	0.76	0.21	54,55,58,58	0
2	NEI	A	401[A]	23/23	0.81	0.23	18,19,23,25	23
3	PG4	A	506[B]	13/13	0.85	0.29	18,25,34,34	13
4	1PE	B	601	16/16	0.85	0.20	38,43,52,53	0
3	PG4	B	501	13/13	0.86	0.22	34,39,56,58	0
3	PG4	A	504	13/13	0.90	0.13	21,26,34,36	0
3	PG4	B	508[B]	13/13	0.94	0.15	23,29,34,35	13

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NEI	B	402[A]	23/23	0.95	0.12	14,16,16,17	23

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NEI A 401 (A):

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NEI B 402 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.