



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 07:40 am BST

PDB ID : 5H94
Title : Crystal structure of Swine MHC CLASSI for 1.48 angstroms
Authors : Fan, S.; Zhang, N.; Wang, S.; Wu, Y.; Xia, C.
Deposited on : 2015-12-25
Resolution : 1.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4690 (1.50-1.46)
Clashscore	141614	4955 (1.50-1.46)
Ramachandran outliers	138981	4846 (1.50-1.46)
Sidechain outliers	138945	4844 (1.50-1.46)
RSRZ outliers	127900	4614 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>2%</div> <div> <div></div> <div>80%</div> <div>19%</div> <div>.</div> </div> </div>
1	D	275	<div> <div>4%</div> <div> <div></div> <div>85%</div> <div>12%</div> <div>.</div> </div> </div>
2	B	98	<div> <div></div> <div> <div></div> <div>83%</div> <div>15%</div> <div>..</div> </div> </div>
2	E	98	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>..</div> </div> </div>
3	C	9	<div> <div></div> <div> <div></div> <div>89%</div> <div>11%</div> </div> </div>
3	F	9	<div> <div></div> <div> <div></div> <div>67%</div> <div>33%</div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7048 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2238	1399	393	439	7			
1	D	275	Total	C	N	O	S	0	0	0
			2238	1399	393	439	7			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	97	Total	C	N	O	S	0	0	0
			800	510	140	147	3			
2	E	97	Total	C	N	O	S	0	0	0
			800	510	140	147	3			

- Molecule 3 is a protein called Nonapeptide from Influenza A virus HA protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	S	0	0	0
			72	45	12	14	1			
3	F	9	Total	C	N	O	S	0	0	0
			72	45	12	14	1			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	330	Total	O	0	0
			330	330		
4	B	148	Total	O	0	0
			148	148		
4	C	12	Total	O	0	0
			12	12		
4	D	228	Total	O	0	0
			228	228		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	103	Total	O	0	0
			103	103		
4	F	7	Total	O	0	0
			7	7		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I antigen




- Molecule 1: MHC class I antigen




- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin




- Molecule 3: Nonapeptide from Influenza A virus HA protein

Chain C:  89% 11%



- Molecule 3: Nonapeptide from Influenza A virus HA protein

Chain F:  67% 33%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	206.45Å 41.47Å 106.73Å 90.00° 117.46° 90.00°	Depositor
Resolution (Å)	50.00 – 1.48 35.32 – 1.48	Depositor EDS
% Data completeness (in resolution range)	99.6 (50.00-1.48) 99.6 (35.32-1.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 1.48Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.150 , 0.190 0.146 , 0.190	Depositor DCC
R_{free} test set	6668 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	11.1	Xtriage
Anisotropy	1.637	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7048	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.36	11/2302 (0.5%)	1.27	20/3131 (0.6%)
1	D	1.31	5/2302 (0.2%)	1.24	15/3131 (0.5%)
2	B	1.30	5/825 (0.6%)	1.06	1/1119 (0.1%)
2	E	1.25	1/825 (0.1%)	1.07	3/1119 (0.3%)
3	C	1.29	0/72	1.05	0/94
3	F	1.46	0/72	1.00	0/94
All	All	1.32	22/6398 (0.3%)	1.21	39/8688 (0.4%)

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	233	GLU	CG-CD	8.16	1.64	1.51
2	B	28	TYR	CG-CD1	7.76	1.49	1.39
1	A	75	TYR	CE1-CZ	7.14	1.47	1.38
1	D	167	GLU	CD-OE2	6.99	1.33	1.25
1	A	265	GLU	CD-OE1	6.84	1.33	1.25
1	D	160	TYR	CE1-CZ	6.75	1.47	1.38
1	D	159	SER	CB-OG	-6.64	1.33	1.42
2	B	28	TYR	CG-CD2	6.13	1.47	1.39
2	E	5	ARG	CZ-NH2	6.04	1.41	1.33
1	A	192	ARG	CD-NE	-5.79	1.36	1.46
1	A	174	GLU	CD-OE1	5.74	1.31	1.25
1	A	205	TRP	CE3-CZ3	5.69	1.48	1.38
1	D	275	TRP	CG-CD1	5.66	1.44	1.36
1	A	107	ASP	CB-CG	5.37	1.63	1.51
2	B	12	TYR	CD1-CE1	5.32	1.47	1.39
1	A	233	GLU	CD-OE2	-5.28	1.19	1.25
1	A	52	TRP	CG-CD1	5.25	1.44	1.36
2	B	51	GLU	CD-OE1	-5.17	1.20	1.25
1	A	258	TYR	CE2-CZ	5.16	1.45	1.38
1	A	18	ARG	CZ-NH1	5.15	1.39	1.33
1	A	230	GLU	CD-OE2	5.14	1.31	1.25
2	B	32	PHE	CE1-CZ	5.03	1.47	1.37

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	276	ASP	CB-CG-OD2	-12.52	107.04	118.30
1	A	192	ARG	NE-CZ-NH2	-12.24	114.18	120.30
1	A	15	ARG	NE-CZ-NH1	-9.94	115.33	120.30
1	A	192	ARG	NE-CZ-NH1	8.02	124.31	120.30
1	D	274	ARG	NE-CZ-NH2	7.88	124.24	120.30
1	A	10	TYR	CB-CG-CD2	-7.83	116.30	121.00
1	A	15	ARG	NE-CZ-NH2	7.58	124.09	120.30
1	D	158	ARG	NE-CZ-NH1	-7.49	116.55	120.30
2	E	5	ARG	NE-CZ-NH2	-7.29	116.65	120.30
1	D	274	ARG	NE-CZ-NH1	-7.20	116.70	120.30
1	A	198	LEU	CA-CB-CG	7.13	131.71	115.30
1	D	276	ASP	CB-CG-OD1	6.98	124.58	118.30
1	A	192	ARG	CG-CD-NE	-6.72	97.68	111.80
1	A	203	ARG	NE-CZ-NH1	-6.72	116.94	120.30
1	D	170	ARG	NE-CZ-NH2	-6.71	116.94	120.30
1	A	203	ARG	NE-CZ-NH2	6.67	123.64	120.30
1	D	192	ARG	NE-CZ-NH2	6.56	123.58	120.30
1	A	274	ARG	NE-CZ-NH1	6.51	123.56	120.30
1	D	203	ARG	NE-CZ-NH2	6.32	123.46	120.30
1	D	20	ASP	CB-CG-OD1	6.29	123.96	118.30
1	A	18	ARG	NE-CZ-NH2	-6.28	117.16	120.30
1	A	130	ASP	CB-CG-OD1	6.27	123.94	118.30
1	A	248	LEU	CB-CG-CD1	6.22	121.57	111.00
1	A	248	LEU	CB-CG-CD2	-6.08	100.66	111.00
1	A	18	ARG	NE-CZ-NH1	5.79	123.20	120.30
1	D	36	ARG	NE-CZ-NH2	-5.73	117.43	120.30
1	A	228	ASP	CB-CG-OD1	5.69	123.42	118.30
2	B	57	PHE	CB-CG-CD2	-5.52	116.94	120.80
1	D	160	TYR	CD1-CE1-CZ	-5.43	114.92	119.80
1	D	167	GLU	OE1-CD-OE2	-5.38	116.85	123.30
1	A	220	ARG	NE-CZ-NH2	5.37	122.98	120.30
1	A	34	PHE	CB-CG-CD1	5.35	124.54	120.80
2	E	47	LYS	CD-CE-NZ	5.34	123.98	111.70
1	D	242	PHE	CB-CG-CD1	-5.30	117.09	120.80
1	D	40	ASP	CB-CG-OD2	-5.29	113.54	118.30
1	A	228	ASP	CB-CG-OD2	-5.29	113.54	118.30
1	D	203	ARG	NE-CZ-NH1	-5.11	117.75	120.30
1	A	158	ARG	NE-CZ-NH2	5.05	122.82	120.30
2	E	40	ASP	CB-CG-OD2	5.03	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2238	0	2065	46	1
1	D	2238	0	2065	34	0
2	B	800	0	770	12	0
2	E	800	0	770	10	0
3	C	72	0	75	0	0
3	F	72	0	75	3	0
4	A	330	0	0	20	2
4	B	148	0	0	9	0
4	C	12	0	0	0	0
4	D	228	0	0	12	0
4	E	103	0	0	4	2
4	F	7	0	0	0	0
All	All	7048	0	5820	96	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:MET:SD	4:A:576:HOH:O	1.95	1.22
1:A:229:MET:CE	4:A:337:HOH:O	1.88	1.20
1:A:2:GLY:N	4:A:302:HOH:O	1.74	1.19
1:A:55:GLN:OE1	4:A:301:HOH:O	1.61	1.16
1:A:182:ARG:NH1	1:D:194:PRO:HG3	1.63	1.13
1:D:276:ASP:HB3	4:D:456:HOH:O	1.53	1.06
1:A:195:SER:HB3	4:A:534:HOH:O	1.56	1.04
1:A:229:MET:HE3	4:A:337:HOH:O	1.50	1.04
1:D:218:TRP:H	1:D:229:MET:HE1	1.27	0.99
1:A:218:TRP:H	1:A:229:MET:HE1	1.21	0.98
1:A:182:ARG:HH22	1:D:192:ARG:HH22	0.99	0.96
1:D:276:ASP:CB	4:D:456:HOH:O	2.06	0.96

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:89:ASP:CG	4:B:201:HOH:O	2.11	0.87
1:D:229:MET:CE	4:D:325:HOH:O	2.22	0.86
2:E:40:ASP:OD1	2:E:47:LYS:HE3	1.78	0.84
1:A:182:ARG:HH22	1:D:192:ARG:NH2	1.77	0.83
1:D:229:MET:HE1	4:D:325:HOH:O	1.76	0.81
2:B:68:HIS:HD2	4:B:318:HOH:O	1.63	0.79
1:A:69:LYS:NZ	4:A:304:HOH:O	2.10	0.78
1:A:182:ARG:HH12	1:D:194:PRO:HG3	1.47	0.77
1:D:112:HIS:HD2	4:D:487:HOH:O	1.68	0.77
1:A:182:ARG:NH2	1:D:192:ARG:HH22	1.81	0.77
2:E:18:GLU:OE2	4:E:201:HOH:O	2.03	0.76
1:D:193:HIS:HE1	2:E:99:ASP:OD2	1.72	0.73
1:A:182:ARG:HH11	1:D:194:PRO:HG3	1.48	0.73
2:E:68:HIS:HD2	4:E:292:HOH:O	1.71	0.73
1:D:275:TRP:O	4:D:301:HOH:O	2.06	0.72
1:A:94:HIS:HD2	1:A:120:ASP:OD2	1.73	0.72
2:E:18:GLU:OE1	2:E:21:LYS:NZ	2.23	0.72
2:B:5:ARG:NE	4:B:202:HOH:O	2.25	0.70
2:B:89:ASP:OD1	4:B:201:HOH:O	2.09	0.69
1:D:94:HIS:HD2	1:D:120:ASP:OD2	1.75	0.69
1:A:218:TRP:N	1:A:229:MET:HE1	2.04	0.68
1:D:226:SER:HA	1:D:229:MET:HE3	1.76	0.67
1:A:276:ASP:C	4:A:308:HOH:O	2.32	0.67
1:A:193:HIS:CE1	4:A:323:HOH:O	2.47	0.67
1:D:218:TRP:N	1:D:229:MET:HE1	2.07	0.67
2:B:5:ARG:CD	4:B:202:HOH:O	2.43	0.67
1:A:178:ASP:O	1:A:182:ARG:CZ	2.42	0.67
1:A:229:MET:HE1	4:A:337:HOH:O	1.74	0.66
1:A:94:HIS:HE1	4:A:388:HOH:O	1.78	0.66
2:B:59:LYS:HD3	4:B:266:HOH:O	1.97	0.64
1:D:220:ARG:HG2	1:D:221:GLU:HG2	1.79	0.64
1:A:196:SER:HB3	1:A:199:GLY:N	2.13	0.63
1:A:195:SER:CB	4:A:534:HOH:O	2.28	0.62
2:B:59:LYS:HD2	4:B:327:HOH:O	1.98	0.62
2:B:68:HIS:HE1	4:B:319:HOH:O	1.81	0.61
1:A:63:ARG:CZ	1:A:67:ILE:HD11	2.31	0.61
1:A:198:LEU:HD12	1:A:252:PRO:HG2	1.81	0.60
1:D:218:TRP:HB2	1:D:229:MET:HE2	1.83	0.60
2:E:68:HIS:HE1	4:E:290:HOH:O	1.83	0.59
1:A:43:ASN:ND2	4:A:304:HOH:O	2.35	0.59
2:B:18:GLU:CD	2:B:21:LYS:HD2	2.24	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:78:GLN:OE1	4:E:202:HOH:O	2.16	0.58
1:D:259:THR:HG22	1:D:274:ARG:HG3	1.86	0.58
1:A:195:SER:HA	4:A:534:HOH:O	2.03	0.57
1:D:107:ASP:HB3	1:D:109:LEU:H	1.68	0.57
1:D:182:ARG:NH2	1:D:184:GLU:OE1	2.38	0.57
1:A:178:ASP:O	1:A:182:ARG:NH1	2.37	0.57
1:D:187:LYS:HE3	4:D:368:HOH:O	2.03	0.57
1:D:94:HIS:HE1	4:D:321:HOH:O	1.89	0.56
1:A:225:GLN:NE2	4:A:311:HOH:O	2.40	0.55
1:A:195:SER:CA	4:A:534:HOH:O	2.52	0.55
1:D:75:TYR:O	1:D:79:LEU:HD13	2.06	0.55
1:D:118:ALA:HB2	2:E:61:TRP:CE2	2.42	0.55
2:B:5:ARG:CZ	4:B:202:HOH:O	2.54	0.54
3:F:5:GLN:HE21	3:F:6:PHE:N	2.06	0.54
1:A:197:ASP:OD1	1:A:198:LEU:N	2.41	0.53
1:A:196:SER:HB3	1:A:199:GLY:H	1.74	0.53
2:B:34:PRO:HB2	2:B:35:PRO:HD2	1.91	0.52
1:D:229:MET:HE3	4:D:325:HOH:O	1.93	0.52
1:A:118:ALA:HB2	2:B:61:TRP:CE2	2.45	0.52
1:A:76:ARG:NH1	4:A:315:HOH:O	2.42	0.51
1:D:220:ARG:HD2	1:D:258:TYR:CZ	2.46	0.50
1:D:276:ASP:HA	4:D:456:HOH:O	2.13	0.49
3:F:5:GLN:HE21	3:F:7:THR:H	1.60	0.48
1:A:269:GLU:HG3	1:A:270:PRO:HD2	1.96	0.48
1:A:191:THR:OG1	1:A:193:HIS:HE1	1.98	0.47
2:E:34:PRO:HB2	2:E:35:PRO:HD2	1.96	0.46
3:F:5:GLN:NE2	3:F:7:THR:H	2.12	0.46
1:A:231:LEU:HD11	1:A:244:LYS:HE3	1.97	0.46
1:D:256:GLN:HG2	4:D:301:HOH:O	2.16	0.45
1:A:193:HIS:HE1	4:A:323:HOH:O	1.94	0.45
1:A:54:GLN:HB2	1:A:54:GLN:HE21	1.50	0.44
1:D:218:TRP:HB2	1:D:229:MET:CE	2.47	0.44
1:D:276:ASP:CA	4:D:456:HOH:O	2.55	0.44
1:D:49:ARG:HA	1:D:49:ARG:HD3	1.82	0.44
1:A:59:GLU:HG2	4:A:520:HOH:O	2.18	0.44
1:D:153:TRP:O	1:D:156:GLN:HB2	2.19	0.43
4:A:476:HOH:O	2:E:18:GLU:HG2	2.18	0.43
1:A:146:ARG:HH11	1:A:146:ARG:HD2	1.63	0.42
1:A:52:TRP:O	1:A:175:MET:CE	2.67	0.42
1:A:198:LEU:HD12	1:A:252:PRO:CG	2.47	0.42
1:A:198:LEU:HD12	1:A:252:PRO:CD	2.50	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:GLN:NE2	1:A:119:TYR:OH	2.52	0.40
1:A:196:SER:O	1:A:197:ASP:C	2.60	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:602:HOH:O	4:E:298:HOH:O[2_557]	2.05	0.15
4:A:614:HOH:O	4:E:253:HOH:O[2_557]	2.13	0.07
1:A:268:GLN:CG	1:A:268:GLN:NE2[2_557]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/275 (99%)	269 (98%)	4 (2%)	0	100	100
1	D	273/275 (99%)	269 (98%)	4 (2%)	0	100	100
2	B	95/98 (97%)	93 (98%)	2 (2%)	0	100	100
2	E	95/98 (97%)	93 (98%)	2 (2%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	F	7/9 (78%)	7 (100%)	0	0	100	100
All	All	750/764 (98%)	738 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/235 (100%)	234 (100%)	1 (0%)	91	81
1	D	235/235 (100%)	230 (98%)	5 (2%)	53	22
2	B	90/91 (99%)	87 (97%)	3 (3%)	38	9
2	E	90/91 (99%)	88 (98%)	2 (2%)	52	20
3	C	8/8 (100%)	7 (88%)	1 (12%)	4	0
3	F	8/8 (100%)	8 (100%)	0	100	100
All	All	666/668 (100%)	654 (98%)	12 (2%)	59	29

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	248	LEU
2	B	59	LYS
2	B	71	PHE
2	B	97	ASP
3	C	4	THR
1	D	3	PRO
1	D	36	ARG
1	D	79	LEU
1	D	107	ASP
1	D	166	VAL
2	E	59	LYS
2	E	71	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	ASN
1	A	55	GLN
1	A	71	ASN
1	A	80	ASN
1	A	88	GLN
1	A	94	HIS
1	A	97	GLN
1	A	142	GLN
1	A	189	HIS
1	A	193	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	225	GLN
1	A	263	GLN
2	B	26	ASN
2	B	68	HIS
3	C	5	GLN
1	D	66	GLN
1	D	73	GLN
1	D	88	GLN
1	D	94	HIS
1	D	97	GLN
1	D	112	HIS
1	D	142	GLN
1	D	156	GLN
2	E	26	ASN
2	E	52	GLN
2	E	68	HIS
3	F	5	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	-0.17	5 (1%) 68 72	11, 16, 32, 78	0
1	D	275/275 (100%)	-0.04	10 (3%) 42 46	13, 20, 42, 61	0
2	B	97/98 (98%)	-0.48	0 100 100	12, 17, 30, 42	0
2	E	97/98 (98%)	-0.20	2 (2%) 63 67	14, 19, 32, 44	0
3	C	9/9 (100%)	-0.30	0 100 100	12, 13, 16, 21	0
3	F	9/9 (100%)	-0.28	0 100 100	13, 20, 30, 34	0
All	All	762/764 (99%)	-0.17	17 (2%) 62 66	11, 18, 36, 78	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	198	LEU	12.1
1	A	197	ASP	6.6
1	D	222	GLY	5.7
1	A	195	SER	5.4
1	D	226	SER	4.6
1	A	196	SER	3.9
1	D	153	TRP	3.4
1	D	227	GLN	3.4
1	D	223	GLN	2.9
2	E	4	ALA	2.6
1	A	227	GLN	2.5
1	D	276	ASP	2.3
2	E	89	ASP	2.3
1	D	221	GLU	2.2
1	D	137	ALA	2.1
1	D	87	ASN	2.1
1	D	18	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.