



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2021 – 12:35 AM EDT

PDB ID : 2HA5
Title : Crystal structure of mutant S203A of acetylcholinesterase complexed with acetylthiocholine
Authors : Bourne, Y.; Radic, Z.; Sulzenbacher, G.; Kim, E.; Taylor, P.; Marchot, P.
Deposited on : 2006-06-12
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	<div> <div>4%</div> <div>88%</div> <div>10%</div> <div>..</div> </div>
1	B	543	<div> <div>8%</div> <div>88%</div> <div>9%</div> <div>..</div> </div>
2	C	2	<div> <div>100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	1	-	-	-	X
2	FUC	C	2	-	-	-	X
4	ACT	A	1903	-	-	X	-
5	AT3	A	1901[A]	-	-	-	X
6	ETM	B	1955	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 9414 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholinesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	535	Total	C	N	O	S	1	7	0
			4211	2702	734	761	14			
1	B	533	Total	C	N	O	S	0	3	0
			4174	2681	722	757	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	203	ALA	SER	engineered mutation	UNP P21836
B	203	ALA	SER	engineered mutation	UNP P21836

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			24	14	1	9			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is ACETYLTHIOCHOLINE (three-letter code: AT3) (formula: $C_7H_{16}NOS$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	1
			10	7	1	1	1		
5	A	1	Total	C	N	O	S	0	0
			10	7	1	1	1		
5	B	1	Total	C	N	O	S	0	1
			10	7	1	1	1		
5	B	1	Total	C	N	O	S	0	0
			10	7	1	1	1		
5	B	1	Total	C	N	O	S	0	0
			10	7	1	1	1		

- Molecule 6 is 2-(TRIMETHYLAMMONIUM)ETHYL THIOL (three-letter code: ETM) (formula: C₅H₁₄NS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	S	0	1
			7	5	1	1		
6	A	1	Total	C	N	S	0	0
			7	5	1	1		
6	B	1	Total	C	N	S	0	1
			7	5	1	1		
6	B	1	Total	C	N	S	0	0
			7	5	1	1		

- Molecule 7 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: $C_{12}H_{26}O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			19	12	7		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	485	Total	O	0	0
			485	485		
8	B	401	Total	O	0	0
			401	401		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Acetylcholinesterase



• Molecule 1: Acetylcholinesterase



• Molecule 2: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.48Å 109.84Å 227.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.15 29.02 – 2.15	Depositor EDS
% Data completeness (in resolution range)	96.6 (30.00-2.15) 86.6 (29.02-2.15)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.25 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.177 , 0.210 0.187 , 0.222	Depositor DCC
R_{free} test set	2062 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	33.9	Xtriage
Anisotropy	0.772	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9414	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P6G, FUC, ACT, AT3, ETM, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	4/4365 (0.1%)	0.72	6/5961 (0.1%)
1	B	0.45	0/4309	0.62	0/5888
All	All	0.52	4/8674 (0.0%)	0.67	6/11849 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	369[A]	GLN	CB-CG	-13.22	1.16	1.52
1	A	369[B]	GLN	CB-CG	-13.22	1.16	1.52
1	A	369[A]	GLN	CA-CB	5.39	1.65	1.53
1	A	369[B]	GLN	CA-CB	5.39	1.65	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	369[A]	GLN	CA-CB-CG	15.94	148.47	113.40
1	A	369[B]	GLN	CA-CB-CG	15.94	148.47	113.40
1	A	369[A]	GLN	N-CA-CB	-5.50	100.69	110.60
1	A	369[B]	GLN	N-CA-CB	-5.50	100.69	110.60
1	A	369[A]	GLN	CB-CA-C	-5.04	100.32	110.40
1	A	369[B]	GLN	CB-CA-C	-5.04	100.32	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4211	0	4102	40	0
1	B	4174	0	4067	45	0
2	C	24	0	22	0	0
3	A	14	0	13	1	0
4	A	4	0	3	2	0
4	B	4	0	3	1	0
5	A	20	0	32	1	0
5	B	30	0	48	6	0
6	A	14	0	28	3	0
6	B	14	0	28	8	0
7	B	19	0	26	0	0
8	A	485	0	0	3	0
8	B	401	0	0	2	0
All	All	9414	0	8372	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:161:LEU:HD11	1:B:269:LEU:HD22	1.59	0.85
1:B:543:THR:HG23	8:B:3179:HOH:O	1.78	0.83
1:B:286:TRP:CZ2	6:B:1955:ETM:H53	2.15	0.81
1:A:337:TYR:OH	6:A:1909[B]:ETM:SD	2.43	0.76
1:B:341:TYR:CE2	6:B:1955:ETM:H42	2.24	0.72
1:B:286:TRP:CZ2	6:B:1955:ETM:C5	2.73	0.72
1:B:72:TYR:CE1	6:B:1955:ETM:H52	2.25	0.72
1:A:48:MET:HE1	1:A:165:ARG:O	1.90	0.71
1:A:380:LEU:HD23	1:A:385:TRP:HZ2	1.57	0.69
5:B:1951[A]:AT3:S24	5:B:1951[A]:AT3:C9	2.81	0.69
5:B:1951[A]:AT3:S24	5:B:1951[A]:AT3:H9C2	2.33	0.68
1:B:203:ALA:HB1	4:B:1953:ACT:C	2.25	0.67
1:A:337:TYR:HA	1:A:443[A]:MET:HE1	1.80	0.64
1:B:48:MET:HE3	1:B:48:MET:HA	1.80	0.63
1:A:203:ALA:HB1	4:A:1903:ACT:C	2.29	0.63
1:A:161:LEU:HD11	1:A:269:LEU:HD22	1.81	0.62
1:B:286:TRP:CE2	6:B:1955:ETM:H53	2.33	0.62
1:A:380:LEU:HD23	1:A:385:TRP:CZ2	2.36	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:MET:HA	1:B:48:MET:CE	2.31	0.61
1:B:495:SER:OG	1:B:496:LYS:N	2.33	0.60
1:B:465:TYR:OH	5:B:1952:AT3:H9C1	2.02	0.60
1:A:414:LEU:HG	1:A:418:LEU:HD22	1.84	0.59
1:B:341:TYR:CD2	6:B:1955:ETM:H42	2.37	0.59
1:B:48:MET:HE1	1:B:165:ARG:O	2.03	0.58
1:B:291:GLN:HE22	1:B:369:GLN:NE2	2.01	0.58
1:A:541:SER:O	1:A:542:ALA:HB3	2.05	0.57
1:A:316:ILE:O	1:A:421:GLN:NE2	2.25	0.56
1:B:291:GLN:HE22	1:B:369:GLN:HE21	1.52	0.56
1:B:134:ASP:OD1	1:B:136:ARG:HD2	2.06	0.55
1:A:48:MET:HE1	1:A:166:GLU:HA	1.88	0.55
1:A:291:GLN:NE2	1:A:369[B]:GLN:HE21	2.05	0.54
1:B:197:VAL:H	1:B:223:HIS:CD2	2.25	0.54
1:B:493:ARG:O	1:B:494:ASP:HB3	2.09	0.53
1:A:337:TYR:HA	1:A:443[A]:MET:CE	2.39	0.52
1:A:541:SER:O	1:A:542:ALA:CB	2.57	0.52
1:A:360:LEU:HD22	1:A:379:VAL:HG21	1.91	0.52
1:B:161:LEU:HD12	1:B:270:ILE:CG1	2.39	0.52
1:B:493:ARG:O	1:B:494:ASP:CB	2.59	0.51
1:B:10:VAL:HG22	1:B:107:ARG:NH1	2.26	0.50
1:A:76:LEU:HD22	1:A:341:TYR:CE2	2.46	0.50
1:A:324:LEU:HG	1:A:326:VAL:HG13	1.94	0.50
1:A:463:LEU:HA	3:A:701:NAG:H82	1.94	0.49
1:B:458:PRO:HA	1:B:465:TYR:CD2	2.48	0.49
1:B:291:GLN:NE2	1:B:369:GLN:HE21	2.11	0.49
1:A:291:GLN:NE2	1:A:369[B]:GLN:NE2	2.60	0.48
1:B:197:VAL:H	1:B:223:HIS:HD2	1.61	0.48
1:B:328:VAL:O	1:B:427:ALA:HA	2.13	0.48
1:B:291:GLN:NE2	1:B:369:GLN:NE2	2.62	0.48
1:A:166:GLU:HB2	1:A:270:ILE:HD13	1.95	0.47
1:A:291:GLN:NE2	8:A:2368:HOH:O	2.35	0.47
1:A:76:LEU:HD22	1:A:341:TYR:CD2	2.49	0.47
1:A:465:TYR:OH	5:A:1902:AT3:H9C1	2.15	0.47
1:B:115:LEU:HD23	1:B:198:THR:HB	1.97	0.47
1:B:312:PRO:O	1:B:316:ILE:HG23	2.15	0.46
1:A:197:VAL:H	1:A:223:HIS:HD2	1.63	0.46
1:B:286:TRP:CZ2	6:B:1955:ETM:H51	2.51	0.46
1:B:511:VAL:HB	1:B:518:LEU:HD22	1.98	0.46
1:B:454:ILE:HD13	1:B:476:LEU:HB3	1.98	0.45
1:A:141:VAL:HG21	1:A:459:LEU:CD2	2.45	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:453:PHE:HB3	1:A:476:LEU:HD12	1.98	0.45
1:B:200:PHE:CB	1:B:226:VAL:HB	2.47	0.45
1:B:209:VAL:CG1	1:B:225:ALA:HB1	2.47	0.45
1:B:539:LEU:O	1:B:543:THR:N	2.50	0.44
1:A:367:VAL:HG12	1:A:370:ALA:HB2	1.99	0.44
1:A:458:PRO:HA	1:A:465:TYR:CD2	2.53	0.44
1:A:300:VAL:HB	1:A:301:PRO:HD2	2.01	0.43
1:B:29:VAL:HG21	1:B:136:ARG:HB2	1.99	0.43
6:B:1959[B]:ETM:H33	6:B:1959[B]:ETM:H12	1.86	0.43
1:A:141:VAL:HG21	1:A:459:LEU:HD23	2.01	0.42
1:A:408:VAL:HG11	1:A:525:ARG:HG3	2.01	0.42
1:A:202:GLU:OE1	6:A:1909[B]:ETM:H41	2.20	0.42
1:B:200:PHE:HB2	1:B:226:VAL:HB	2.01	0.42
1:A:339:LEU:HD13	1:A:346:PHE:CE2	2.54	0.42
1:B:166:GLU:HB2	1:B:270:ILE:HD13	2.02	0.42
1:B:211:MET:HG2	1:B:308:LEU:HD21	2.00	0.42
1:A:297:PHE:CZ	4:A:1903:ACT:H2	2.55	0.41
1:B:112:THR:HG21	1:B:143:GLY:O	2.20	0.41
5:B:1951[A]:AT3:S24	5:B:1951[A]:AT3:H9C3	2.61	0.41
1:B:161:LEU:HD12	1:B:270:ILE:HD11	2.03	0.41
5:B:1952:AT3:H9C2	8:B:3066:HOH:O	2.19	0.41
1:A:265:ASN:ND2	8:A:2302:HOH:O	2.53	0.41
1:A:328:VAL:O	1:A:427:ALA:HA	2.20	0.41
1:A:337:TYR:CZ	6:A:1909[B]:ETM:SD	3.11	0.41
1:B:113:PRO:HG2	1:B:485:ARG:HG2	2.03	0.41
1:B:337:TYR:CE2	5:B:1951[A]:AT3:H6C1	2.56	0.41
1:A:257:CYS:C	8:A:2019:HOH:O	2.60	0.40
1:B:373:LEU:HD23	1:B:539:LEU:HD11	2.03	0.40
1:B:528:THR:O	1:B:531:PHE:HB3	2.21	0.40
1:A:197:VAL:H	1:A:223:HIS:CD2	2.39	0.40
1:A:202:GLU:HA	1:A:228:GLN:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	538/543 (99%)	519 (96%)	17 (3%)	2 (0%)	34	29
1	B	532/543 (98%)	513 (96%)	15 (3%)	4 (1%)	19	12
All	All	1070/1086 (98%)	1032 (96%)	32 (3%)	6 (1%)	25	18

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	494	ASP
1	B	495	SER
1	A	541	SER
1	B	496	LYS
1	A	342	GLY
1	B	342	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	445/442 (101%)	433 (97%)	12 (3%)	44	46
1	B	440/442 (100%)	434 (99%)	6 (1%)	67	72
All	All	885/884 (100%)	867 (98%)	18 (2%)	57	59

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ARG
1	A	45	ARG
1	A	200	PHE
1	A	216	LEU
1	A	291	GLN
1	A	295	PHE
1	A	327	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	369[A]	GLN
1	A	369[B]	GLN
1	A	418	LEU
1	A	437	LEU
1	A	493	ARG
1	B	136	ARG
1	B	200	PHE
1	B	291	GLN
1	B	295	PHE
1	B	376	GLU
1	B	413	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	HIS
1	A	291	GLN
1	A	322	GLN
1	A	325	GLN
1	B	223	HIS
1	B	291	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.65	0	17,19,21	1.23	2 (11%)
2	FUC	C	2	2	10,10,11	0.63	0	14,14,16	1.12	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	FUC	C	2	2	-	-	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	C1-O5-C5	3.15	116.46	112.19
2	C	1	NAG	O5-C1-C2	-2.31	107.64	111.29
2	C	2	FUC	C3-C4-C5	2.29	113.34	109.77
2	C	2	FUC	O5-C5-C4	2.04	113.19	109.52

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	P6G	B	2901	-	18,18,18	2.19	6 (33%)	17,17,17	1.28	1 (5%)
3	NAG	A	701	1	14,14,15	0.50	0	17,19,21	0.91	1 (5%)
6	ETM	A	1909[B]	-	6,6,6	1.02	0	7,8,8	1.03	1 (14%)
5	AT3	A	1902	-	9,9,9	0.79	0	12,12,12	0.45	0
6	ETM	B	1959[B]	-	6,6,6	1.06	0	7,8,8	0.68	0
4	ACT	B	1953	-	1,3,3	1.39	0	0,3,3	-	-
5	AT3	B	1952	-	9,9,9	0.86	0	12,12,12	0.62	0
5	AT3	B	1951[A]	-	9,9,9	1.18	0	12,12,12	0.50	0
5	AT3	B	1961	-	9,9,9	0.80	0	12,12,12	0.83	1 (8%)
4	ACT	A	1903	-	1,3,3	1.14	0	0,3,3	-	-
6	ETM	B	1955	-	6,6,6	0.71	0	7,8,8	0.96	0
5	AT3	A	1901[A]	-	9,9,9	1.17	0	12,12,12	0.75	0
6	ETM	A	1905	-	6,6,6	0.63	0	7,8,8	1.30	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	P6G	B	2901	-	-	8/16/16/16	-
3	NAG	A	701	1	-	0/6/23/26	0/1/1/1
6	ETM	A	1909[B]	-	-	1/4/4/4	-
5	AT3	A	1902	-	-	0/7/7/7	-
6	ETM	B	1959[B]	-	-	4/4/4/4	-
5	AT3	B	1952	-	-	3/7/7/7	-
5	AT3	B	1951[A]	-	-	7/7/7/7	-
5	AT3	B	1961	-	-	0/7/7/7	-
6	ETM	B	1955	-	-	1/4/4/4	-
5	AT3	A	1901[A]	-	-	5/7/7/7	-
6	ETM	A	1905	-	-	4/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	2901	P6G	O10-C9	4.33	1.60	1.42
7	B	2901	P6G	O13-C12	3.76	1.58	1.42
7	B	2901	P6G	O16-C15	3.55	1.57	1.42
7	B	2901	P6G	O19-C18	3.50	1.60	1.42
7	B	2901	P6G	O4-C3	3.46	1.57	1.42
7	B	2901	P6G	O7-C6	2.83	1.54	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	2901	P6G	O1-C2-C3	3.32	131.09	111.81
3	A	701	NAG	O5-C5-C6	2.34	110.88	107.20
6	A	1905	ETM	C3-N1-C2	2.30	119.34	109.92
5	B	1961	AT3	C3-S24-C5	2.27	113.61	101.68
6	A	1909[B]	ETM	C3-N1-C2	2.03	118.21	109.92

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	1951[A]	AT3	N1-C2-C3-S24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	B	1951[A]	AT3	C2-C3-S24-C5
5	B	1951[A]	AT3	O7-C5-S24-C3
5	B	1951[A]	AT3	C6-C5-S24-C3
5	B	1952	AT3	O7-C5-S24-C3
6	A	1905	ETM	SD-C1-C2-N1
6	B	1959[B]	ETM	SD-C1-C2-N1
7	B	2901	P6G	O16-C17-C18-O19
7	B	2901	P6G	O7-C8-C9-O10
5	B	1952	AT3	C6-C5-S24-C3
7	B	2901	P6G	O13-C14-C15-O16
6	B	1959[B]	ETM	C1-C2-N1-C3
6	B	1959[B]	ETM	C1-C2-N1-C5
5	A	1901[A]	AT3	N1-C2-C3-S24
5	B	1952	AT3	C2-C3-S24-C5
6	B	1955	ETM	SD-C1-C2-N1
7	B	2901	P6G	C2-C3-O4-C5
7	B	2901	P6G	C8-C9-O10-C11
7	B	2901	P6G	C11-C12-O13-C14
5	A	1901[A]	AT3	C3-C2-N1-C9
5	A	1901[A]	AT3	C3-C2-N1-C10
5	B	1951[A]	AT3	C3-C2-N1-C9
5	B	1951[A]	AT3	C3-C2-N1-C10
6	A	1909[B]	ETM	C1-C2-N1-C3
6	A	1905	ETM	C1-C2-N1-C3
6	B	1959[B]	ETM	C1-C2-N1-C4
7	B	2901	P6G	O4-C5-C6-O7
7	B	2901	P6G	C6-C5-O4-C3
5	A	1901[A]	AT3	C2-C3-S24-C5
5	A	1901[A]	AT3	C3-C2-N1-C8
5	B	1951[A]	AT3	C3-C2-N1-C8
6	A	1905	ETM	C1-C2-N1-C4
6	A	1905	ETM	C1-C2-N1-C5

There are no ring outliers.

9 monomers are involved in 22 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	701	NAG	1	0
6	A	1909[B]	ETM	3	0
5	A	1902	AT3	1	0
6	B	1959[B]	ETM	1	0
4	B	1953	ACT	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1952	AT3	2	0
5	B	1951[A]	AT3	4	0
4	A	1903	ACT	2	0
6	B	1955	ETM	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	535/543 (98%)	0.13	21 (3%)	39 48	41, 53, 70, 98	0
1	B	533/543 (98%)	0.32	42 (7%)	12 17	43, 59, 78, 99	0
All	All	1068/1086 (98%)	0.22	63 (5%)	22 30	41, 55, 76, 99	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	495	SER	7.2
1	A	493	ARG	6.3
1	B	492	PRO	6.1
1	B	543	THR	5.8
1	B	495	SER	4.7
1	B	106	PRO	4.5
1	B	109	ALA	4.4
1	B	542	ALA	4.4
1	A	497	SER	4.1
1	A	492	PRO	4.1
1	A	216	LEU	3.9
1	A	494	ASP	3.8
1	B	540	LEU	3.8
1	A	542	ALA	3.7
1	B	138	LEU	3.6
1	B	497	SER	3.6
1	B	42	VAL	3.6
1	B	322	GLN	3.5
1	A	540	LEU	3.5
1	B	541	SER	3.4
1	B	146	LEU	3.4
1	B	45[A]	ARG	3.3
1	B	101	VAL	3.3
1	A	496	LYS	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	493	ARG	3.2
1	A	541	SER	3.1
1	B	461	PRO	3.0
1	B	135	GLY	3.0
1	A	165	ARG	2.9
1	B	455	PHE	2.9
1	B	323	ASP	2.9
1	A	323[A]	ASP	2.9
1	B	318	THR	2.8
1	B	451	ILE	2.8
1	B	4	GLU	2.8
1	B	99	LEU	2.7
1	A	317	ASN	2.7
1	B	41	PRO	2.6
1	B	147	VAL	2.6
1	A	75	THR	2.6
1	A	322	GLN	2.5
1	A	454	ILE	2.5
1	B	281	LEU	2.5
1	A	226	VAL	2.5
1	B	265	ASN	2.5
1	B	47	PHE	2.5
1	B	156	PHE	2.4
1	B	102	TRP	2.3
1	B	117	TRP	2.3
1	A	480	TRP	2.3
1	B	35	ILE	2.3
1	B	467	THR	2.2
1	B	457	LEU	2.2
1	B	268	GLU	2.2
1	B	200	PHE	2.2
1	B	319	GLY	2.2
1	B	155	THR	2.2
1	B	13	ARG	2.1
1	A	491	ASP	2.1
1	A	318	THR	2.1
1	B	480	TRP	2.1
1	A	78	PRO	2.0
1	B	393	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FUC	C	2	10/11	0.47	0.63	102,104,105,105	0
2	NAG	C	1	14/15	0.77	0.43	83,90,94,98	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	ETM	A	1905	7/7	0.71	0.30	94,95,96,98	0
3	NAG	A	701	14/15	0.74	0.38	81,86,89,90	0
6	ETM	B	1955	7/7	0.75	0.34	93,93,95,97	7
5	AT3	A	1901[A]	10/10	0.80	0.49	80,81,83,84	10
5	AT3	B	1952	10/10	0.85	0.26	86,90,91,91	0
5	AT3	A	1902	10/10	0.85	0.28	78,80,81,81	0
5	AT3	B	1951[A]	10/10	0.85	0.33	78,79,83,84	10
6	ETM	A	1909[B]	7/7	0.86	0.28	58,58,60,62	7
4	ACT	B	1953	4/4	0.88	0.25	62,64,64,64	0
6	ETM	B	1959[B]	7/7	0.89	0.20	73,74,75,77	7
7	P6G	B	2901	19/19	0.89	0.16	67,72,77,77	0
4	ACT	A	1903	4/4	0.91	0.21	59,60,61,61	0
5	AT3	B	1961	10/10	0.95	0.17	68,70,71,73	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around P6G B 2901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.