



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 10:07 am BST

PDB ID : 2HAW  
Title : Crystal structure of family II Inorganic pyrophosphatase in complex with PNP  
Authors : Fabrichniy, I.P.; Lehtio, L.; Oksanen, E.; Goldman, A.  
Deposited on : 2006-06-13  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	<div> <div>19%</div> <div> <div></div> <div>94%</div> <div>5% ..</div> </div> </div>
1	B	309	<div> <div>18%</div> <div> <div></div> <div>93%</div> <div>5% •</div> </div> </div>

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 5483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Manganese-dependent inorganic pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	307	Total	C	N	O	S	0	8	0
			2431	1533	403	488	7			
1	B	308	Total	C	N	O	S	0	16	0
			2507	1575	413	511	8			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	4	Total	Mg	0	0
			4	4		
2	A	4	Total	Mg	0	0
			4	4		

- Molecule 3 is FLUORIDE ION (three-letter code: F) (formula: F).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	F	0	0
			1	1		
3	A	1	Total	F	0	0
			1	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	1
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	1
			5	4	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			13	8	5		
6	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 7 is IMIDODIPHOSPHORIC ACID (three-letter code: 2PN) (formula:  $\text{H}_5\text{NO}_6\text{P}_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	N	O	P	0	1
			9	1	6	2		
7	B	1	Total	N	O	P	0	1
			9	1	6	2		

- Molecule 8 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula:  $C_{10}H_{22}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			16	10	6		

- Molecule 9 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			6	3	3		
9	B	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	166	Total	O	0	0
			166	166		
10	B	270	Total	O	0	0
			270	270		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Manganese-dependent inorganic pyrophosphatase



- Molecule 1: Manganese-dependent inorganic pyrophosphatase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.01Å 115.87Å 147.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.84 – 1.75 19.80 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.84-1.75) 100.0 (19.80-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.45 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.166 , 0.186 0.170 , 0.188	Depositor DCC
$R_{free}$ test set	5228 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.3	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 55.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5483	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, F, CL, 1PE, PG4, SO4, 2PN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	1/2456 (0.0%)	0.68	1/3324 (0.0%)
1	B	0.80	1/2532 (0.0%)	0.78	2/3426 (0.1%)
All	All	0.71	2/4988 (0.0%)	0.73	3/6750 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	51	ASP	CG-OD1	6.18	1.39	1.25
1	B	114	GLU	CB-CG	5.09	1.61	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	79	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	B	191	LEU	CA-CB-CG	5.22	127.31	115.30
1	B	288	LEU	CB-CG-CD2	5.04	119.56	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2431	0	2472	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2507	0	2530	7	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
3	A	1	0	0	1	0
3	B	1	0	0	0	0
4	A	10	0	0	0	0
4	B	10	0	0	0	0
5	A	1	0	0	0	0
6	A	13	0	18	1	0
6	B	13	0	18	0	0
7	A	9	0	1	0	0
7	B	9	0	1	0	0
8	B	16	0	22	1	0
9	B	18	0	24	1	0
10	A	166	0	0	0	0
10	B	270	0	0	0	0
All	All	5483	0	5086	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:ASN:HD21	1:A:34:ALA:H	1.30	0.79
1:A:101:ALA:HA	1:B:104[B]:GLU:HG2	1.69	0.73
1:B:66:ASN:H	1:B:66:ASN:HD22	1.39	0.71
1:B:66:ASN:H	1:B:66:ASN:ND2	2.03	0.54
1:A:79:ARG:NH2	1:A:102[B]:ASN:OD1	2.42	0.53
8:B:1001:1PE:H251	8:B:1001:1PE:H122	1.94	0.49
1:B:191:LEU:C	1:B:193:LYS:H	2.18	0.47
1:A:15:ASP:HB3	3:A:3001:F:F	2.06	0.45
1:B:96:ASP:O	1:B:113:ALA:HA	2.16	0.45
1:A:133:LYS:CD	1:A:133:LYS:H	2.28	0.45
6:A:6002:PG4:H62	6:A:6002:PG4:H42	1.52	0.44
1:B:186:LYS:HE2	1:B:186:LYS:HB3	1.73	0.43
1:B:57[B]:SER:HB3	9:B:5003:GOL:H2	2.01	0.42
1:A:42:VAL:HG13	1:A:46:THR:HB	2.01	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/309 (101%)	305 (97%)	8 (3%)	0	100	100
1	B	322/309 (104%)	314 (98%)	8 (2%)	0	100	100
All	All	635/618 (103%)	619 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/262 (102%)	259 (97%)	9 (3%)	37	14
1	B	277/262 (106%)	260 (94%)	17 (6%)	18	4
All	All	545/524 (104%)	519 (95%)	26 (5%)	30	7

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLU
1	A	16	THR
1	A	97	HIS
1	A	99	ARG
1	A	133	LYS
1	A	219[A]	VAL
1	A	219[B]	VAL
1	A	227	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	263	LEU
1	B	16	THR
1	B	33	ASN
1	B	66	ASN
1	B	84	LYS
1	B	97	HIS
1	B	99	ARG
1	B	178[A]	GLU
1	B	178[B]	GLU
1	B	186	LYS
1	B	191	LEU
1	B	202[A]	LEU
1	B	202[B]	LEU
1	B	225[A]	GLU
1	B	225[B]	GLU
1	B	280	VAL
1	B	288	LEU
1	B	307	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	183	ASN
1	A	269	ASN
1	B	33	ASN
1	B	66	ASN
1	B	90	GLN
1	B	284	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 11 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	1PE	B	1001	-	15,15,15	0.54	0	14,14,14	1.05	1 (7%)
9	GOL	B	5003	-	5,5,5	0.43	0	5,5,5	0.78	0
6	PG4	A	6002	-	12,12,12	0.47	0	11,11,11	0.35	0
4	SO4	B	4004[B]	2	4,4,4	0.18	0	6,6,6	0.09	0
7	2PN	A	2001[A]	2	8,8,8	5.03	6 (75%)	8,13,13	3.07	2 (25%)
9	GOL	B	5002	-	5,5,5	0.39	0	5,5,5	0.73	0
6	PG4	B	1002	-	12,12,12	0.62	0	11,11,11	0.48	0
4	SO4	A	4002	-	4,4,4	0.17	0	6,6,6	0.36	0
4	SO4	A	4003[B]	2	4,4,4	0.14	0	6,6,6	0.24	0
9	GOL	B	5001	-	5,5,5	0.56	0	5,5,5	1.24	1 (20%)
4	SO4	B	4001	-	4,4,4	0.18	0	6,6,6	0.97	1 (16%)
7	2PN	B	2002[A]	2	8,8,8	4.32	5 (62%)	8,13,13	2.28	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	GOL	B	5003	-	-	2/4/4/4	-
6	PG4	A	6002	-	-	6/10/10/10	-
9	GOL	B	5002	-	-	3/4/4/4	-
6	PG4	B	1002	-	-	2/10/10/10	-
7	2PN	B	2002[A]	2	-	2/2/6/6	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	1PE	B	1001	-	-	6/13/13/13	-
9	GOL	B	5001	-	-	2/4/4/4	-
7	2PN	A	2001[A]	2	-	2/2/6/6	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	2002[A]	2PN	P1-O1	9.00	1.60	1.46
7	A	2001[A]	2PN	P2-O4	8.00	1.58	1.46
7	A	2001[A]	2PN	P1-N1	7.74	1.83	1.63
7	A	2001[A]	2PN	P2-N1	-5.96	1.47	1.63
7	A	2001[A]	2PN	P2-O5	-4.67	1.44	1.56
7	B	2002[A]	2PN	P2-O6	-4.55	1.44	1.56
7	B	2002[A]	2PN	P2-N1	4.38	1.74	1.63
7	B	2002[A]	2PN	P2-O4	4.05	1.52	1.46
7	A	2001[A]	2PN	P1-O1	3.88	1.52	1.46
7	B	2002[A]	2PN	P1-O2	2.37	1.63	1.56
7	A	2001[A]	2PN	P1-O2	2.27	1.62	1.56

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	2001[A]	2PN	O4-P2-N1	-6.87	101.65	111.77
7	B	2002[A]	2PN	O4-P2-N1	-5.06	104.32	111.77
7	A	2001[A]	2PN	O2-P1-O1	-4.96	100.99	113.45
7	B	2002[A]	2PN	O3-P1-O1	-2.45	107.29	113.45
8	B	1001	1PE	OH4-C24-C14	2.18	120.21	110.39
4	B	4001	SO4	O4-S-O3	2.07	117.89	109.06
9	B	5001	GOL	O1-C1-C2	-2.03	100.44	110.20

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	2001[A]	2PN	P2-N1-P1-O1
7	A	2001[A]	2PN	P1-N1-P2-O4
7	B	2002[A]	2PN	P2-N1-P1-O1
7	B	2002[A]	2PN	P1-N1-P2-O4
9	B	5003	GOL	C1-C2-C3-O3
9	B	5003	GOL	O2-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
9	B	5001	GOL	C1-C2-C3-O3
9	B	5001	GOL	O2-C2-C3-O3
6	A	6002	PG4	C6-C5-O3-C4
8	B	1001	1PE	C14-C24-OH4-C13
8	B	1001	1PE	OH5-C14-C24-OH4
9	B	5002	GOL	O1-C1-C2-C3
6	B	1002	PG4	O1-C1-C2-O2
6	A	6002	PG4	O2-C3-C4-O3
6	A	6002	PG4	O4-C7-C8-O5
9	B	5002	GOL	O1-C1-C2-O2
6	B	1002	PG4	O2-C3-C4-O3
8	B	1001	1PE	OH6-C15-C25-OH5
6	A	6002	PG4	C5-C6-O4-C7
8	B	1001	1PE	C23-C13-OH4-C24
6	A	6002	PG4	O1-C1-C2-O2
6	A	6002	PG4	C8-C7-O4-C6
9	B	5002	GOL	C1-C2-C3-O3
8	B	1001	1PE	C24-C14-OH5-C25
8	B	1001	1PE	OH2-C12-C22-OH3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	1001	1PE	1	0
9	B	5003	GOL	1	0
6	A	6002	PG4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	307/309 (99%)	0.94	59 (19%)	1 1	30, 41, 57, 70	0
1	B	308/309 (99%)	0.97	56 (18%)	1 1	33, 40, 64, 80	0
All	All	615/618 (99%)	0.96	115 (18%)	1 1	30, 40, 60, 80	0

All (115) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	258	LEU	6.9
1	B	192	SER	6.8
1	A	308	ALA	5.1
1	B	308	ALA	4.8
1	A	258	LEU	4.7
1	A	143	LEU	4.5
1	A	193	LYS	4.4
1	B	191	LEU	4.3
1	A	147	ILE	4.3
1	A	211	SER	4.3
1	A	171	GLU	4.2
1	B	226[A]	ASP	4.2
1	B	279[A]	ASN	4.1
1	A	145	ALA	4.1
1	A	305	ASP	4.0
1	B	291	GLY	3.9
1	A	279	ASN	3.8
1	B	210	GLY	3.7
1	A	225	GLU	3.7
1	B	223	ASP	3.6
1	A	257	ILE	3.6
1	A	222	VAL	3.6
1	A	176	ASP	3.6
1	A	283	GLU	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	222	VAL	3.5
1	B	211	SER	3.5
1	B	260	ASN	3.5
1	A	251	LEU	3.5
1	A	32	PHE	3.5
1	A	209	LEU	3.4
1	B	257	ILE	3.4
1	B	252	LEU	3.4
1	B	306	ALA	3.4
1	A	192	SER	3.3
1	A	31	GLY	3.3
1	B	188	GLY	3.3
1	A	179	GLU	3.3
1	A	191	LEU	3.3
1	A	41	GLN	3.3
1	A	146	ILE	3.3
1	A	182	LEU	3.1
1	B	132	VAL	3.1
1	B	146	ILE	3.1
1	B	141	LEU	3.1
1	B	251	LEU	3.0
1	B	283	GLU	3.0
1	A	281	THR	3.0
1	B	193	LYS	3.0
1	A	174	GLY	3.0
1	A	229	LYS	3.0
1	B	143	LEU	3.0
1	A	173	ALA	2.9
1	B	224	ILE	2.9
1	B	7	PHE	2.8
1	B	5	LEU	2.8
1	B	225[A]	GLU	2.8
1	B	169	LEU	2.8
1	B	68	VAL	2.8
1	A	142	MET	2.8
1	A	167	LYS	2.8
1	A	234	LEU	2.8
1	B	73	LEU	2.7
1	A	120	ALA	2.7
1	B	281	THR	2.7
1	B	139	ALA	2.6
1	B	216	ILE	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	253	VAL	2.6
1	A	178	GLU	2.6
1	B	37	VAL	2.6
1	A	52	TYR	2.6
1	A	136	LYS	2.5
1	B	209	LEU	2.5
1	A	223	ASP	2.5
1	A	56	GLU	2.4
1	A	226	ASP	2.4
1	B	259	GLU	2.4
1	A	7	PHE	2.4
1	A	227	VAL	2.4
1	A	238	ILE	2.4
1	A	249	LEU	2.4
1	A	183	ASN	2.4
1	B	190	ASP	2.4
1	B	6	ILE	2.4
1	B	72	ILE	2.4
1	B	134	ILE	2.4
1	A	133	LYS	2.4
1	A	137	GLU	2.4
1	A	95	ILE	2.3
1	A	267	ILE	2.3
1	B	179	GLU	2.3
1	B	229	LYS	2.3
1	A	151	LEU	2.3
1	A	88[A]	GLU	2.3
1	A	160	ASP	2.3
1	A	276	LYS	2.3
1	B	276	LYS	2.3
1	A	131	ASN	2.2
1	A	300	VAL	2.2
1	B	305	ASP	2.2
1	B	300	VAL	2.2
1	A	265	LEU	2.2
1	A	202[A]	LEU	2.1
1	A	291	GLY	2.1
1	B	140	GLY	2.1
1	A	119	THR	2.1
1	B	64	ALA	2.1
1	B	142	MET	2.1
1	B	238	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	17	ILE	2.1
1	A	260	ASN	2.0
1	B	87[A]	GLU	2.0
1	B	267	ILE	2.0
1	B	299	VAL	2.0
1	B	109	LEU	2.0
1	B	249	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	CL	A	6001	1/1	0.76	0.11	86,86,86,86	0
6	PG4	B	1002	13/13	0.76	0.21	48,56,81,82	0
6	PG4	A	6002	13/13	0.77	0.22	64,72,82,82	0
9	GOL	B	5001	6/6	0.83	0.35	42,54,60,66	0
9	GOL	B	5003	6/6	0.84	0.24	53,62,68,69	0
9	GOL	B	5002	6/6	0.84	0.26	72,77,77,79	0
4	SO4	A	4002	5/5	0.85	0.35	84,85,87,88	0
8	1PE	B	1001	16/16	0.87	0.16	44,54,66,69	0
2	MG	A	1004	1/1	0.91	0.48	42,42,42,42	1
2	MG	A	1001	1/1	0.93	0.15	32,32,32,32	0
2	MG	A	1002	1/1	0.95	0.10	33,33,33,33	0
2	MG	B	1008	1/1	0.96	0.26	33,33,33,33	1
4	SO4	B	4001	5/5	0.97	0.29	51,55,60,63	0
3	F	A	3001	1/1	0.98	0.33	35,35,35,35	0
2	MG	A	1003	1/1	0.98	0.15	35,35,35,35	1
7	2PN	A	2001[A]	9/9	0.98	0.12	29,32,34,34	9

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	B	1006	1/1	0.98	0.07	27,27,27,27	0
4	SO4	B	4004[B]	5/5	0.99	0.08	31,31,32,32	5
3	F	B	3002	1/1	0.99	0.19	27,27,27,27	0
2	MG	B	1005	1/1	0.99	0.10	27,27,27,27	0
7	2PN	B	2002[A]	9/9	0.99	0.09	23,28,29,31	9
2	MG	B	1007	1/1	0.99	0.16	32,32,32,32	1
4	SO4	A	4003[B]	5/5	0.99	0.10	28,29,30,30	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.