



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2021 – 12:15 PM EDT

PDB ID : 3HAB  
Title : The structure of DPP4 in complex with piperidine fused benzimidazole 25  
Authors : Scapin, G.  
Deposited on : 2009-05-01  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	728	<div> <div>5%</div> <div>94%</div> <div>6%</div> </div>
1	B	728	<div> <div>4%</div> <div>93%</div> <div>7%</div> </div>
2	C	2	<div> <div>100%</div> </div>
2	D	2	<div> <div>100%</div> </div>
2	E	2	<div> <div>50%</div> <div>50%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	2	 100%
2	G	2	 100%
2	H	2	 100%
2	I	2	 100%
2	J	2	 100%
2	K	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	2	X	-	-	-
2	NAG	D	2	-	-	-	X
2	NAG	F	2	X	-	-	-
2	NAG	G	2	-	-	-	X
3	NAG	B	801	X	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 13644 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 4 soluble form.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	728	Total	C	N	O	S	0	12	0
			6014	3860	990	1137	27			
1	B	728	Total	C	N	O	S	0	13	0
			6023	3865	993	1138	27			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	THR	SER	engineered mutation	UNP P27487
B	39	THR	SER	engineered mutation	UNP P27487

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	J	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	K	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is (2R,3R)-7-(methylsulfonyl)-3-(2,4,5-trifluorophenyl)-1,2,3,4-tetrahydropyrido[1,2-a]benzimidazol-2-amine (three-letter code: 677) (formula: C<sub>18</sub>H<sub>16</sub>F<sub>3</sub>N<sub>3</sub>O<sub>2</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	F	N	O	S	0	0
			27	18	3	3	2	1		
5	B	1	Total	C	F	N	O	S	0	0
			27	18	3	3	2	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	625	Total	O	0	0
			625	625		
6	B	619	Total	O	0	0
			619	619		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dipeptidyl peptidase 4 soluble form



- Molecule 1: Dipeptidyl peptidase 4 soluble form



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1  
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50%  50%


MAG1  
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%


MAG1  
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

MAG1  
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%


MAG1  
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%

MAG1  
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  100%

MAG1  
MAG2



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:

100%

MAG1  
MAG2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	117.91Å 125.64Å 136.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 29.96 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-2.10) 99.9 (29.96-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.89 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.169 , 0.211 0.184 , 0.218	Depositor DCC
$R_{free}$ test set	5950 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.8	Xtriage
Anisotropy	0.121	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	13644	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 677, NAG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.57	0/6231	0.63	1/8472 (0.0%)
1	B	0.56	0/6247	0.61	0/8495
All	All	0.56	0/12478	0.62	1/16967 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	184	ARG	NE-CZ-NH1	5.55	123.08	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6014	0	5723	28	0
1	B	6023	0	5738	31	0
2	C	28	0	25	0	0
2	D	28	0	25	1	0
2	E	28	0	25	0	0
2	F	28	0	25	0	0
2	G	28	0	25	0	0
2	H	28	0	25	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	I	28	0	25	1	0
2	J	28	0	25	0	0
2	K	28	0	25	0	0
3	A	14	0	13	0	0
3	B	42	0	39	1	0
4	A	1	0	0	0	0
5	A	27	0	16	0	0
5	B	27	0	16	0	0
6	A	625	0	0	2	0
6	B	619	0	0	4	0
All	All	13644	0	11770	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:258:LYS:HZ1	1:B:712:HIS:HD2	1.06	0.92
1:B:214[B]:LEU:HD23	1:B:225:TYR:HB3	1.56	0.87
1:A:69:LEU:HD23	1:A:78:VAL:HG22	1.61	0.82
1:A:184:ARG:HD2	1:A:186:THR:O	1.88	0.74
1:B:466:LYS:HE2	6:B:1449:HOH:O	1.90	0.71
1:A:258:LYS:HZ1	1:A:712:HIS:HD2	1.39	0.70
1:B:172:ILE:HD13	1:B:214[B]:LEU:HD21	1.76	0.68
1:A:258:LYS:NZ	1:A:712:HIS:HD2	1.92	0.68
1:B:258:LYS:HZ1	1:B:712:HIS:CD2	1.99	0.68
1:B:47:ASP:HA	1:B:52[B]:THR:HG23	1.78	0.64
1:A:78:VAL:HG23	1:A:89:PHE:HB2	1.78	0.64
1:B:105:TYR:HB2	1:B:114:ILE:HD11	1.78	0.64
1:B:47:ASP:HA	1:B:52[B]:THR:CG2	2.28	0.63
1:B:75:ASN:HD21	3:B:801:NAG:HN2	1.46	0.62
1:B:726:VAL:HG23	1:B:728:VAL:HG23	1.81	0.62
1:B:272:ASN:C	1:B:272:ASN:HD22	2.02	0.62
1:B:696:LYS:HG3	1:B:728:VAL:HG22	1.81	0.62
1:A:69:LEU:HD23	1:A:78:VAL:CG2	2.31	0.60
1:B:272:ASN:HD21	1:B:274:ASP:HB2	1.66	0.60
1:A:69:LEU:CD2	1:A:78:VAL:HG22	2.33	0.59
1:A:153:GLN:HE22	1:A:170:ASN:ND2	2.01	0.59
1:A:640:LEU:HD11	1:A:650:GLY:HA3	1.86	0.58
1:A:184:ARG:HG2	1:A:187:TRP:CE2	2.39	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:GLU:OE1	1:A:253[B]:ARG:HD2	2.05	0.56
1:B:114:ILE:HG23	1:B:135:TYR:HB3	1.90	0.54
1:A:55:LEU:HD12	1:A:500:LEU:CD2	2.37	0.54
1:A:253[A]:ARG:HH21	1:B:253[A]:ARG:HH21	1.56	0.54
1:A:471[A]:ARG:HD2	6:A:1197:HOH:O	2.09	0.53
6:B:933:HOH:O	2:I:1:NAG:H82	2.10	0.51
1:A:539:LYS:HE2	1:A:617:GLY:O	2.12	0.49
1:A:154:TRP:CE2	1:A:212[A]:SER:HB3	2.48	0.48
1:B:471[B]:ARG:HG3	1:B:480:TYR:CE2	2.47	0.48
1:B:50:LYS:HD3	6:B:1048:HOH:O	2.14	0.48
2:D:1:NAG:H4	2:D:2:NAG:H2	1.66	0.47
1:A:471[B]:ARG:HD2	1:A:480:TYR:HE2	1.80	0.47
1:A:248:TYR:CZ	1:B:234:PRO:HB2	2.50	0.46
1:A:471[A]:ARG:HG3	1:A:480:TYR:CE2	2.51	0.46
1:B:546:VAL:HG21	1:B:626:ILE:HD11	1.98	0.46
1:B:214[B]:LEU:HD22	1:B:223:LEU:HD21	1.96	0.46
1:B:154:TRP:CE2	1:B:212[A]:SER:HB2	2.51	0.46
1:B:146:GLU:OE2	6:B:901:HOH:O	2.20	0.45
1:B:69:LEU:HD11	1:B:107:ILE:HG13	1.98	0.45
1:A:170:ASN:HD22	1:A:170:ASN:N	2.15	0.44
1:B:501:ASP:O	1:B:505:GLN:HG2	2.18	0.44
1:A:726:VAL:HG23	1:A:728:VAL:HG23	1.98	0.44
1:B:237:GLU:HA	1:B:252:VAL:O	2.17	0.44
1:A:154:TRP:CE2	1:A:212[B]:SER:HB2	2.53	0.43
1:B:65:ASP:OD2	1:B:466:LYS:HB2	2.18	0.43
1:A:95:PHE:HB3	1:A:98:PHE:HB2	1.99	0.43
1:A:224:ALA:HB1	1:A:268:PHE:CZ	2.53	0.43
1:A:510:PRO:HD3	1:A:569:SER:HB2	1.99	0.43
1:A:332:GLU:HG3	6:A:1347:HOH:O	2.19	0.43
1:B:640:LEU:HD11	1:B:650:GLY:HA3	2.01	0.42
1:A:40:ARG:HB2	1:A:506:ASN:O	2.19	0.42
1:B:272:ASN:HD22	1:B:274:ASP:H	1.68	0.41
1:A:453:ARG:HG3	1:A:476:GLY:HA3	2.03	0.41
1:B:323[A]:SER:OG	1:B:347:GLU:HB3	2.20	0.41
1:B:345:HIS:HE1	1:B:389:ILE:O	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	737/728 (101%)	715 (97%)	21 (3%)	1 (0%)	51	54
1	B	739/728 (102%)	714 (97%)	25 (3%)	0	100	100
All	All	1476/1456 (101%)	1429 (97%)	46 (3%)	1 (0%)	51	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	714	GLN

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	664/653 (102%)	654 (98%)	10 (2%)	65	71
1	B	666/653 (102%)	653 (98%)	13 (2%)	55	60
All	All	1330/1306 (102%)	1307 (98%)	23 (2%)	60	67

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	LEU
1	A	97	GLU
1	A	116	LEU
1	A	170	ASN
1	A	263	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	366	LEU
1	A	385	CYS
1	A	391	LYS
1	A	472	CYS
1	A	566	TYR
1	B	51	ASN
1	B	66	HIS
1	B	97	GLU
1	B	230	ASP
1	B	272	ASN
1	B	385	CYS
1	B	450[A]	ASN
1	B	450[B]	ASN
1	B	472	CYS
1	B	487	ASN
1	B	627	TRP
1	B	685	ASN
1	B	704	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	123	GLN
1	A	141	GLN
1	A	169	ASN
1	A	170	ASN
1	A	263	ASN
1	A	388	GLN
1	A	435	GLN
1	A	572	ASN
1	A	694	ASN
1	A	712	HIS
1	B	74	ASN
1	B	169	ASN
1	B	272	ASN
1	B	345	HIS
1	B	388	GLN
1	B	487	ASN
1	B	533	HIS
1	B	572	ASN
1	B	685	ASN
1	B	694	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	712	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	C	1	1,2	14,14,15	0.61	0	17,19,21	2.14	8 (47%)
2	NAG	C	2	2	14,14,15	0.57	0	17,19,21	1.14	1 (5%)
2	NAG	D	1	1,2	14,14,15	0.55	0	17,19,21	1.51	3 (17%)
2	NAG	D	2	2	14,14,15	0.84	0	17,19,21	1.61	5 (29%)
2	NAG	E	1	1,2	14,14,15	0.55	0	17,19,21	0.69	0
2	NAG	E	2	2	14,14,15	0.48	0	17,19,21	0.87	1 (5%)
2	NAG	F	1	1,2	14,14,15	0.63	0	17,19,21	1.14	1 (5%)
2	NAG	F	2	2	14,14,15	0.82	0	17,19,21	1.38	3 (17%)
2	NAG	G	1	1,2	14,14,15	0.63	0	17,19,21	1.48	4 (23%)
2	NAG	G	2	2	14,14,15	0.77	0	17,19,21	1.37	3 (17%)
2	NAG	H	1	1,2	14,14,15	0.61	0	17,19,21	1.32	2 (11%)
2	NAG	H	2	2	14,14,15	0.78	0	17,19,21	1.51	2 (11%)
2	NAG	I	1	1,2	14,14,15	0.59	0	17,19,21	0.88	0
2	NAG	I	2	2	14,14,15	0.61	0	17,19,21	1.16	2 (11%)
2	NAG	J	1	1,2	14,14,15	0.62	0	17,19,21	1.04	1 (5%)



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	J	2	2	14,14,15	0.57	0	17,19,21	1.18	1 (5%)
2	NAG	K	1	1,2	14,14,15	0.64	0	17,19,21	1.10	1 (5%)
2	NAG	K	2	2	14,14,15	0.50	0	17,19,21	1.28	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	1/1/5/7	2/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	D	2	2	-	5/6/23/26	0/1/1/1
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	1/1/5/7	3/6/23/26	0/1/1/1
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	6/6/23/26	0/1/1/1
2	NAG	H	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	H	2	2	-	4/6/23/26	0/1/1/1
2	NAG	I	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	I	2	2	-	2/6/23/26	0/1/1/1
2	NAG	J	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	J	2	2	-	2/6/23/26	0/1/1/1
2	NAG	K	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	K	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	2	NAG	C4-C3-C2	4.30	117.33	111.02
2	C	1	NAG	O4-C4-C3	4.14	119.92	110.35
2	C	1	NAG	O5-C5-C6	3.87	113.27	107.20
2	D	1	NAG	C1-O5-C5	3.65	117.13	112.19
2	H	1	NAG	C4-C3-C2	3.64	116.35	111.02
2	C	1	NAG	C4-C3-C2	-3.42	106.01	111.02

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	NAG	O4-C4-C3	-3.38	102.53	110.35
2	H	2	NAG	C3-C4-C5	3.29	116.10	110.24
2	G	2	NAG	C4-C3-C2	3.18	115.68	111.02
2	G	1	NAG	O4-C4-C3	-3.18	103.01	110.35
2	F	2	NAG	C2-N2-C7	3.16	127.40	122.90
2	F	1	NAG	C1-O5-C5	3.11	116.41	112.19
2	K	2	NAG	O5-C5-C6	3.06	112.01	107.20
2	C	2	NAG	O5-C1-C2	-3.05	106.47	111.29
2	J	2	NAG	C1-O5-C5	3.03	116.30	112.19
2	D	2	NAG	C2-N2-C7	3.03	127.21	122.90
2	D	2	NAG	C1-O5-C5	2.99	116.24	112.19
2	F	2	NAG	C4-C3-C2	2.96	115.35	111.02
2	D	2	NAG	C4-C3-C2	2.86	115.22	111.02
2	K	2	NAG	C1-O5-C5	2.83	116.03	112.19
2	G	2	NAG	C3-C4-C5	2.80	115.23	110.24
2	G	1	NAG	C4-C3-C2	2.57	114.78	111.02
2	J	1	NAG	O5-C1-C2	-2.53	107.29	111.29
2	H	1	NAG	O5-C5-C6	2.48	111.10	107.20
2	I	2	NAG	C4-C3-C2	2.47	114.64	111.02
2	G	1	NAG	C1-O5-C5	2.46	115.53	112.19
2	C	1	NAG	O5-C5-C4	-2.45	104.87	110.83
2	G	2	NAG	C2-N2-C7	2.45	126.39	122.90
2	E	2	NAG	C1-O5-C5	2.42	115.47	112.19
2	D	2	NAG	C3-C4-C5	2.39	114.50	110.24
2	C	1	NAG	O4-C4-C5	2.36	115.15	109.30
2	G	1	NAG	O5-C1-C2	-2.28	107.69	111.29
2	C	1	NAG	C3-C4-C5	-2.27	106.19	110.24
2	C	1	NAG	O5-C1-C2	2.27	114.87	111.29
2	K	1	NAG	O5-C1-C2	-2.16	107.87	111.29
2	C	1	NAG	C1-O5-C5	2.16	115.12	112.19
2	F	2	NAG	O5-C5-C4	-2.15	105.61	110.83
2	D	2	NAG	C1-C2-N2	2.10	114.07	110.49
2	I	2	NAG	C3-C4-C5	2.05	113.89	110.24
2	D	1	NAG	C4-C3-C2	2.03	114.00	111.02

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	2	NAG	C1
2	F	2	NAG	C1

All (32) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2
2	F	2	NAG	C8-C7-N2-C2
2	F	2	NAG	O7-C7-N2-C2
2	G	2	NAG	C8-C7-N2-C2
2	G	2	NAG	O7-C7-N2-C2
2	H	2	NAG	C8-C7-N2-C2
2	H	2	NAG	O7-C7-N2-C2
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
2	E	2	NAG	O5-C5-C6-O6
2	H	2	NAG	O5-C5-C6-O6
2	E	2	NAG	C4-C5-C6-O6
2	I	2	NAG	C4-C5-C6-O6
2	I	2	NAG	O5-C5-C6-O6
2	K	2	NAG	C4-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
2	G	2	NAG	O5-C5-C6-O6
2	H	2	NAG	C4-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	K	2	NAG	O5-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	G	2	NAG	C4-C5-C6-O6
2	G	2	NAG	C1-C2-N2-C7
2	J	2	NAG	C4-C5-C6-O6
2	F	2	NAG	C3-C2-N2-C7
2	G	2	NAG	C3-C2-N2-C7
2	J	1	NAG	C4-C5-C6-O6
2	J	2	NAG	O5-C5-C6-O6
2	D	2	NAG	C1-C2-N2-C7
2	D	1	NAG	C4-C5-C6-O6


There are no ring outliers.


3 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2	NAG	1	0
2	I	1	NAG	1	0
2	D	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

























## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	B	803	1	14,14,15	0.51	0	17,19,21	1.47	1 (5%)
3	NAG	A	801	1	14,14,15	0.52	0	17,19,21	1.32	2 (11%)
5	677	B	804	-	27,30,30	2.65	6 (22%)	33,47,47	2.39	9 (27%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	677	A	803	-	27,30,30	2.68	6 (22%)	33,47,47	2.11	7 (21%)
3	NAG	B	802	1	14,14,15	0.53	0	17,19,21	0.94	1 (5%)
3	NAG	B	801	1	14,14,15	0.54	0	17,19,21	1.28	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	803	1	-	0/6/23/26	0/1/1/1
3	NAG	A	801	1	-	2/6/23/26	0/1/1/1
5	677	B	804	-	-	4/10/22/22	0/3/4/4
5	677	A	803	-	-	4/10/22/22	0/3/4/4
3	NAG	B	802	1	-	0/6/23/26	0/1/1/1
3	NAG	B	801	1	1/1/5/7	3/6/23/26	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	804	677	C21-S24	-11.61	1.64	1.77
5	A	803	677	C21-S24	-11.58	1.64	1.77
5	A	803	677	C29-C19	4.19	1.49	1.40
5	B	804	677	C29-C19	3.06	1.46	1.40
5	B	804	677	C27-S24	-2.77	1.64	1.75
5	A	803	677	C27-S24	-2.68	1.64	1.75
5	A	803	677	C23-C22	2.68	1.42	1.36
5	B	804	677	C10-C11	2.61	1.53	1.50
5	B	804	677	C23-C22	2.57	1.42	1.36
5	B	804	677	C20-C21	2.56	1.41	1.36
5	A	803	677	C20-C21	2.23	1.40	1.36
5	A	803	677	C10-C11	2.08	1.52	1.50

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	804	677	C27-S24-C21	7.22	113.11	104.58
5	A	803	677	C27-S24-C21	6.68	112.47	104.58
5	B	804	677	O25-S24-O26	-6.24	107.17	117.92
5	A	803	677	O25-S24-O26	-6.09	107.43	117.92

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	803	NAG	C1-O5-C5	4.56	118.37	112.19
3	B	801	NAG	C1-O5-C5	4.13	117.79	112.19
5	B	804	677	C4-C3-C2	4.01	119.82	116.48
5	A	803	677	O25-S24-C21	-3.69	105.24	108.25
5	B	804	677	O26-S24-C21	3.55	111.14	108.25
5	B	804	677	C11-C10-C9	-3.48	109.34	113.99
5	B	804	677	O25-S24-C21	-3.45	105.43	108.25
5	B	804	677	C1-C2-C3	-3.12	120.10	123.83
3	A	801	NAG	O5-C5-C6	3.09	112.05	107.20
3	A	801	NAG	C1-O5-C5	2.92	116.14	112.19
5	B	804	677	O26-S24-C27	2.84	112.57	108.49
5	A	803	677	O26-S24-C27	2.61	112.25	108.49
5	A	803	677	C4-C3-C2	2.20	118.31	116.48
3	B	802	NAG	O5-C5-C6	2.16	110.59	107.20
5	B	804	677	C22-C21-C20	-2.14	118.43	121.05
5	A	803	677	C11-C10-C9	-2.04	111.27	113.99
5	A	803	677	C23-C22-C21	2.02	122.53	119.74

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	B	801	NAG	C1

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	801	NAG	C8-C7-N2-C2
3	B	801	NAG	O7-C7-N2-C2
5	A	803	677	C20-C21-S24-C27
5	B	804	677	C22-C21-S24-C27
5	B	804	677	C20-C21-S24-C27
5	A	803	677	C22-C21-S24-C27
5	A	803	677	C22-C21-S24-O25
5	B	804	677	C22-C21-S24-O25
5	B	804	677	C20-C21-S24-O25
3	A	801	NAG	C4-C5-C6-O6
5	A	803	677	C20-C21-S24-O25
3	A	801	NAG	O5-C5-C6-O6
3	B	801	NAG	C1-C2-N2-C7


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	801	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	728/728 (100%)	0.03	34 (4%) 31 37	16, 24, 37, 51	0
1	B	728/728 (100%)	-0.02	29 (3%) 38 44	17, 24, 38, 54	0
All	All	1456/1456 (100%)	0.01	63 (4%) 35 41	16, 24, 38, 54	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	39	THR	7.4
1	B	39	THR	5.8
1	A	279	VAL	5.5
1	B	766	PRO	5.2
1	A	99	GLY	4.5
1	A	97	GLU	4.4
1	A	74	ASN	4.4
1	A	96	ASP	4.1
1	A	73	GLU	4.0
1	B	73	GLU	4.0
1	B	138	ASN	4.0
1	A	138	ASN	3.9
1	A	766	PRO	3.9
1	B	74	ASN	3.9
1	B	92	ASN	3.9
1	B	97	GLU	3.8
1	A	92	ASN	3.4
1	A	93	SER	3.3
1	A	333	SER	3.2
1	B	333	SER	3.2
1	B	96	ASP	3.1
1	A	75	ASN	3.1
1	A	280	THR	3.0
1	A	40	ARG	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	40	ARG	3.0
1	B	506	ASN	3.0
1	A	140	ARG	2.9
1	B	71	LYS	2.8
1	A	278	SER	2.8
1	A	98	PHE	2.7
1	B	91	GLU	2.7
1	B	502	LYS	2.7
1	B	765	LEU	2.7
1	A	521	GLU	2.6
1	A	546[A]	VAL	2.6
1	B	98	PHE	2.6
1	B	537	SER	2.6
1	A	335	GLY	2.6
1	B	334	SER	2.6
1	B	99	GLY	2.6
1	B	140	ARG	2.5
1	B	93	SER	2.5
1	B	521	GLU	2.5
1	B	100	HIS	2.5
1	A	336	ARG	2.4
1	B	95	PHE	2.4
1	A	141	GLN	2.4
1	B	72	GLN	2.3
1	A	656	VAL	2.3
1	A	144	THR	2.3
1	A	506	ASN	2.3
1	A	137	LEU	2.3
1	A	765	LEU	2.3
1	B	88	VAL	2.2
1	B	505	GLN	2.2
1	B	358[A]	ARG	2.2
1	B	378	GLU	2.1
1	A	665	VAL	2.1
1	A	520	ASN	2.1
1	A	332	GLU	2.1
1	A	330	TYR	2.1
1	A	334	SER	2.0
1	A	667	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	H	2	14/15	0.43	0.38	65,68,70,71	0
2	NAG	G	2	14/15	0.60	0.41	58,61,62,63	0
2	NAG	C	2	14/15	0.60	0.36	64,67,68,68	0
2	NAG	D	2	14/15	0.61	0.47	67,69,71,71	0
2	NAG	K	2	14/15	0.62	0.28	48,52,53,53	0
2	NAG	F	2	14/15	0.69	0.33	56,60,65,65	0
2	NAG	D	1	14/15	0.76	0.32	51,56,59,64	0
2	NAG	E	2	14/15	0.76	0.29	50,52,56,57	0
2	NAG	I	2	14/15	0.83	0.28	41,46,49,50	0
2	NAG	G	1	14/15	0.84	0.25	41,45,48,54	0
2	NAG	H	1	14/15	0.85	0.18	41,50,57,60	0
2	NAG	J	2	14/15	0.86	0.31	41,45,47,47	0
2	NAG	E	1	14/15	0.88	0.21	37,43,47,47	0
2	NAG	C	1	14/15	0.88	0.17	44,48,55,60	0
2	NAG	K	1	14/15	0.90	0.13	30,33,40,44	0
2	NAG	F	1	14/15	0.93	0.15	35,39,44,51	0
2	NAG	I	1	14/15	0.93	0.20	30,37,41,43	0
2	NAG	J	1	14/15	0.94	0.16	23,27,33,37	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain G:**



$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around Chain H:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain I:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain J:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	B	802	14/15	0.68	0.36	49,53,55,56	0
3	NAG	B	801	14/15	0.76	0.46	66,69,70,71	0
3	NAG	A	801	14/15	0.79	0.22	42,46,48,48	0
3	NAG	B	803	14/15	0.81	0.25	39,43,44,44	0
4	NA	A	802	1/1	0.93	0.06	25,25,25,25	0
5	677	A	803	27/27	0.96	0.08	13,18,21,25	0
5	677	B	804	27/27	0.96	0.08	15,20,23,26	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around 677 A 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 677 B 804:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.