



wwPDB X-ray Structure Validation Summary Report ⓘ

May 17, 2020 – 03:48 pm BST

PDB ID : 4HDH
Title : Crystal Structure of viral RdRp in complex with ATP
Authors : Surana, P.; Nair, D.T.
Deposited on : 2012-10-02
Resolution : 2.28 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	639	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>16%</div> <div>• • •</div> </div> </div>
1	B	639	<div> <div>4%</div> <div> <div></div> <div>80%</div> <div>13%</div> <div>• •</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10309 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	613	Total	C	N	O	S	0	0	0
			4920	3104	886	899	31			
1	B	613	Total	C	N	O	S	0	0	0
			4904	3095	879	899	31			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	267	GLY	-	EXPRESSION TAG	UNP G3LHD9
A	268	PRO	-	EXPRESSION TAG	UNP G3LHD9
A	269	LEU	-	EXPRESSION TAG	UNP G3LHD9
A	270	GLY	-	EXPRESSION TAG	UNP G3LHD9
A	271	SER	-	EXPRESSION TAG	UNP G3LHD9
A	373	ARG	LYS	CONFLICT	UNP G3LHD9
A	429	ASN	ASP	CONFLICT	UNP G3LHD9
A	836	ALA	THR	CONFLICT	UNP G3LHD9
B	267	GLY	-	EXPRESSION TAG	UNP G3LHD9
B	268	PRO	-	EXPRESSION TAG	UNP G3LHD9
B	269	LEU	-	EXPRESSION TAG	UNP G3LHD9
B	270	GLY	-	EXPRESSION TAG	UNP G3LHD9
B	271	SER	-	EXPRESSION TAG	UNP G3LHD9
B	373	ARG	LYS	CONFLICT	UNP G3LHD9
B	429	ASN	ASP	CONFLICT	UNP G3LHD9
B	836	ALA	THR	CONFLICT	UNP G3LHD9

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
2	B	1	Total 31	C 10	N 5	O 13	P 3	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Zn 2 2	0	0
3	A	2	Total Zn 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	200	Total O 200 200	0	0
4	B	219	Total O 219 219	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	144.84Å 86.51Å 112.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	144.84 – 2.28 44.92 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.0 (144.84-2.28) 99.1 (44.92-2.28)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.06 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.225 , 0.272 0.220 , 0.267	Depositor DCC
R_{free} test set	4577 reflections (7.05%)	wwPDB-VP
Wilson B-factor (Å ²)	34.3	Xtriage
Anisotropy	0.619	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10309	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.58 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.3114e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	0/5037	0.84	12/6818 (0.2%)
1	B	0.89	0/5021	0.83	9/6800 (0.1%)
All	All	0.89	0/10058	0.83	21/13618 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	B	0	5
All	All	0	11

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	502	ARG	NE-CZ-NH2	-10.43	115.08	120.30
1	B	502	ARG	NE-CZ-NH2	-9.35	115.63	120.30
1	A	502	ARG	NE-CZ-NH1	8.82	124.71	120.30
1	B	502	ARG	NE-CZ-NH1	7.53	124.06	120.30
1	A	500	LEU	CA-CB-CG	-7.08	99.01	115.30

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	408	ASN	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	414	VAL	Peptide
1	A	415	PHE	Peptide
1	A	416	ALA	Peptide
1	A	417	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4920	0	4819	83	0
1	B	4904	0	4786	74	0
2	A	31	0	12	4	0
2	B	31	0	12	5	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	200	0	0	4	0
4	B	219	0	0	3	0
All	All	10309	0	9629	160	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 160 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1001:ATP:H4'	2:A:1001:ATP:O1B	1.52	1.10
1:B:531:LYS:HG2	1:B:706:HIS:HD2	1.22	0.99
1:A:602:ARG:HG3	1:A:604:SER:H	1.29	0.98
1:B:531:LYS:HG2	1:B:706:HIS:CD2	2.04	0.92
1:B:602:ARG:HG3	1:B:604:SER:H	1.33	0.90

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	609/639 (95%)	572 (94%)	30 (5%)	7 (1%)	14	14
1	B	609/639 (95%)	575 (94%)	29 (5%)	5 (1%)	19	22
All	All	1218/1278 (95%)	1147 (94%)	59 (5%)	12 (1%)	15	16

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	420	GLN
1	B	408	ASN
1	B	411	LEU
1	A	297	ASP
1	A	408	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	515/544 (95%)	483 (94%)	32 (6%)	18	23
1	B	512/544 (94%)	478 (93%)	34 (7%)	16	20
All	All	1027/1088 (94%)	961 (94%)	66 (6%)	17	21

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	861	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	332	LYS
1	B	847	ARG
1	A	874	GLN
1	B	299	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	443	ASN
1	A	577	HIS
1	A	706	HIS
1	A	803	HIS
1	B	706	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ATP	A	1001	-	26,33,33	1.64	2 (7%)	31,52,52	2.54	9 (29%)
2	ATP	B	1001	-	26,33,33	1.63	4 (15%)	31,52,52	2.32	10 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	1001	-	-	5/18/38/38	0/3/3/3
2	ATP	B	1001	-	-	7/18/38/38	0/3/3/3

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	ATP	O4'-C1'	6.56	1.50	1.41
2	B	1001	ATP	O4'-C1'	5.95	1.49	1.41
2	A	1001	ATP	PG-O3G	3.01	1.66	1.54
2	B	1001	ATP	PG-O2G	2.85	1.65	1.54
2	B	1001	ATP	C5'-C4'	2.42	1.59	1.51

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	ATP	PB-O3B-PG	-8.45	103.84	132.83
2	B	1001	ATP	PB-O3B-PG	-6.03	112.13	132.83
2	B	1001	ATP	N3-C2-N1	-5.78	119.64	128.68
2	A	1001	ATP	N3-C2-N1	-5.52	120.05	128.68
2	B	1001	ATP	PA-O3A-PB	-4.48	117.45	132.83

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1001	ATP	C4'-C5'-O5'-PA
2	A	1001	ATP	O4'-C4'-C5'-O5'
2	A	1001	ATP	C3'-C4'-C5'-O5'
2	B	1001	ATP	C5'-O5'-PA-O1A
2	B	1001	ATP	C5'-O5'-PA-O2A


There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	ATP	4	0
2	B	1001	ATP	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	613/639 (95%)	0.04	28 (4%) 32 38	25, 45, 74, 94	0
1	B	613/639 (95%)	-0.01	23 (3%) 40 45	26, 46, 73, 92	0
All	All	1226/1278 (95%)	0.01	51 (4%) 36 41	25, 45, 73, 94	0

The worst 5 of 51 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	419	ASN	5.8
1	A	748	GLY	5.7
1	A	416	ALA	5.1
1	B	416	ALA	4.9
1	B	890	SER	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ATP	B	1001	31/31	0.83	0.23	48,63,68,71	0
2	ATP	A	1001	31/31	0.84	0.21	58,68,73,74	0
3	ZN	A	1002	1/1	0.99	0.02	28,28,28,28	0
3	ZN	B	1002	1/1	0.99	0.02	33,33,33,33	0
3	ZN	A	1003	1/1	0.99	0.01	39,39,39,39	0
3	ZN	B	1003	1/1	0.99	0.02	39,39,39,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.