



Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 07:35 pm BST

PDB ID : 1HEH
Title : C-terminal xylan binding domain from Cellulomonas fimi xylanase 11A
Authors : Simpson, P.J.; Hefang, X.; Bolam, D.N.; White, P.; Hancock, S.M.; Gilbert, H.J.; Williamson, M.P.
Deposited on : 2000-11-22

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 57%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	C	88	

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1185 atoms, of which 570 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called ENDO-1,4-BETA-XYLANASE D.


Mol	Chain	Residues	Atoms						Trace
1	C	88	Total	C	H	N	O	S	0
			1185	372	570	109	132	2	

4 Residue-property plots [i](#)

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: ENDO-1,4-BETA-XYLANASE D

Chain C:



5 Refinement protocol and experimental data overview

The models were refined using the following method: *HYBRID DISTANCE GEOMETRY/ SIMULATED ANNEALING*.

Of the 50 calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.1
Felix	structure solution	97.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	622
Number of shifts mapped to atoms	622
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	57%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	C	0	3
All	All	0	3

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	C	566	ARG	Sidechain
1	C	573	ARG	Sidechain
1	C	616	ARG	Sidechain

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	C	615	570	567	80
All	All	615	570	567	80

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 68.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:C:588:VAL:HG23	1:C:640:THR:O	1.04	1.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:C:580:VAL:HG21	1:C:586:TRP:CZ2	0.99	1.92
1:C:592:LEU:HD13	1:C:596:GLN:O	0.88	1.68
1:C:580:VAL:HG21	1:C:586:TRP:CH2	0.86	2.05
1:C:607:THR:O	1:C:613:VAL:HG22	0.85	1.71
1:C:601:SER:OG	1:C:606:LEU:HD21	0.80	1.77
1:C:636:THR:HG22	1:C:637:PRO:HD2	0.79	1.54
1:C:588:VAL:CG2	1:C:640:THR:O	0.77	2.33
1:C:586:TRP:CE3	1:C:615:ALA:HB3	0.74	2.17
1:C:580:VAL:CG2	1:C:586:TRP:CH2	0.72	2.72
1:C:565:VAL:O	1:C:576:VAL:HG13	0.67	1.89
1:C:588:VAL:HG21	1:C:639:ALA:HB1	0.67	1.67
1:C:603:ASN:O	1:C:604:ALA:HB2	0.66	1.91
1:C:586:TRP:CE3	1:C:615:ALA:CB	0.64	2.81
1:C:574:PHE:CZ	1:C:628:PHE:CB	0.63	2.81
1:C:559:SER:OG	1:C:643:THR:HG23	0.62	1.95
1:C:560:CYS:SG	1:C:562:VAL:CG1	0.61	2.89
1:C:636:THR:HG22	1:C:637:PRO:CD	0.61	2.26
1:C:565:VAL:O	1:C:576:VAL:CG1	0.60	2.49
1:C:588:VAL:CG2	1:C:589:THR:N	0.60	2.65
1:C:606:LEU:N	1:C:606:LEU:HD23	0.59	2.11
1:C:580:VAL:CG2	1:C:586:TRP:CZ2	0.58	2.79
1:C:588:VAL:HG22	1:C:589:THR:N	0.58	2.12
1:C:562:VAL:CG1	1:C:641:CYS:SG	0.58	2.92
1:C:562:VAL:HG11	1:C:641:CYS:SG	0.57	2.39
1:C:588:VAL:O	1:C:614:THR:HG23	0.57	2.00
1:C:601:SER:HB3	1:C:626:VAL:HG22	0.56	1.77
1:C:574:PHE:CE1	1:C:628:PHE:HB2	0.55	2.37
1:C:603:ASN:O	1:C:604:ALA:CB	0.54	2.56
1:C:586:TRP:O	1:C:587:VAL:CG2	0.54	2.56
1:C:636:THR:CG2	1:C:637:PRO:HD2	0.54	2.32
1:C:602:TRP:O	1:C:603:ASN:CB	0.53	2.57
1:C:574:PHE:CZ	1:C:628:PHE:HB2	0.53	2.39
1:C:559:SER:CB	1:C:643:THR:HG23	0.53	2.34
1:C:627:THR:O	1:C:628:PHE:CG	0.52	2.61
1:C:619:GLY:O	1:C:620:SER:CB	0.52	2.58
1:C:574:PHE:CZ	1:C:628:PHE:HB3	0.52	2.40
1:C:613:VAL:CG1	1:C:614:THR:N	0.51	2.74
1:C:559:SER:N	1:C:643:THR:CG2	0.50	2.74
1:C:587:VAL:HG12	1:C:587:VAL:O	0.50	2.05
1:C:588:VAL:O	1:C:614:THR:HA	0.50	2.07
1:C:580:VAL:HG11	1:C:583:SER:HB2	0.49	1.83
1:C:566:ARG:O	1:C:567:GLY:O	0.49	2.30

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:C:583:SER:O	1:C:584:SER:C	0.48	2.51
1:C:570:TRP:CZ3	1:C:575:ASN:HB2	0.48	2.44
1:C:586:TRP:C	1:C:587:VAL:HG23	0.48	2.29
1:C:588:VAL:HG23	1:C:640:THR:N	0.48	2.23
1:C:606:LEU:N	1:C:606:LEU:CD2	0.47	2.76
1:C:567:GLY:O	1:C:568:GLU:O	0.47	2.33
1:C:588:VAL:CG2	1:C:640:THR:N	0.47	2.78
1:C:568:GLU:O	1:C:575:ASN:HB3	0.46	2.11
1:C:624:PHE:CD1	1:C:624:PHE:C	0.46	2.89
1:C:593:ASN:O	1:C:594:GLY:O	0.46	2.33
1:C:574:PHE:CE1	1:C:636:THR:HG22	0.46	2.46
1:C:602:TRP:O	1:C:603:ASN:HB2	0.45	2.11
1:C:561:SER:CB	1:C:581:SER:HB2	0.45	2.41
1:C:567:GLY:O	1:C:568:GLU:C	0.45	2.54
1:C:601:SER:OG	1:C:606:LEU:CD2	0.45	2.60
1:C:609:SER:O	1:C:610:SER:CB	0.44	2.66
1:C:559:SER:CA	1:C:643:THR:CG2	0.44	2.96
1:C:563:SER:O	1:C:578:TYR:HA	0.43	2.12
1:C:588:VAL:HG23	1:C:640:THR:C	0.43	2.29
1:C:630:LYS:O	1:C:631:ASN:C	0.43	2.56
1:C:586:TRP:C	1:C:587:VAL:CG2	0.42	2.87
1:C:605:ALA:C	1:C:606:LEU:HD23	0.42	2.34
1:C:607:THR:O	1:C:613:VAL:HA	0.42	2.14
1:C:627:THR:HG22	1:C:628:PHE:N	0.42	2.30
1:C:559:SER:CA	1:C:643:THR:HG23	0.42	2.44
1:C:630:LYS:HB3	1:C:633:SER:OG	0.42	2.15
1:C:574:PHE:CZ	1:C:637:PRO:HD3	0.42	2.50
1:C:595:GLY:O	1:C:596:GLN:C	0.42	2.57
1:C:633:SER:O	1:C:634:SER:OG	0.42	2.33
1:C:580:VAL:HB	1:C:622:ASN:O	0.41	2.15
1:C:572:ASP:O	1:C:630:LYS:HG3	0.41	2.16
1:C:634:SER:O	1:C:635:ALA:C	0.41	2.57
1:C:624:PHE:CD1	1:C:624:PHE:O	0.41	2.74
1:C:575:ASN:ND2	1:C:627:THR:HA	0.41	2.30
1:C:570:TRP:HB2	1:C:573:ARG:O	0.41	2.16
1:C:607:THR:O	1:C:613:VAL:CG2	0.41	2.58
1:C:618:ASN:C	1:C:618:ASN:OD1	0.40	2.58

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	86/88 (98%)	57 (66%)	19 (22%)	10 (12%)	1	7
All	All	86/88 (98%)	57 (66%)	19 (22%)	10 (12%)	1	7

All 10 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	C	610	SER
1	C	620	SER
1	C	558	GLY
1	C	568	GLU
1	C	604	ALA
1	C	634	SER
1	C	594	GLY
1	C	592	LEU
1	C	567	GLY
1	C	587	VAL

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	66/66 (100%)	46 (70%)	20 (30%)	1	16
All	All	66/66 (100%)	46 (70%)	20 (30%)	1	16

All 20 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	C	610	SER
1	C	581	SER
1	C	623	SER
1	C	606	LEU
1	C	585	SER
1	C	590	LEU
1	C	620	SER
1	C	573	ARG
1	C	589	THR
1	C	559	SER
1	C	640	THR
1	C	600	SER
1	C	597	SER
1	C	560	CYS
1	C	609	SER
1	C	636	THR
1	C	584	SER
1	C	566	ARG
1	C	603	ASN
1	C	630	LYS

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.6 Ligand geometry ⓘ

There are no ligands in this entry.

6.7 Other polymers ⓘ

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 57% for the well-defined parts and 57% for the entire structure.

7.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	622
Number of shifts mapped to atoms	622
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	0	—	None (insufficient data)
$^{13}\text{C}_\beta$	0	—	None (insufficient data)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	83	-0.50 ± 0.56	None needed (imprecise)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 57%, i.e. 512 atoms were assigned a chemical shift out of a possible 898. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	256/436 (59%)	173/174 (99%)	0/176 (0%)	83/86 (97%)
Sidechain	212/383 (55%)	201/223 (90%)	0/142 (0%)	11/18 (61%)

Continued on next page...

Continued from previous page...

	Total	¹H	¹³C	¹⁵N
Aromatic	44/79 (56%)	41/41 (100%)	0/35 (0%)	3/3 (100%)
Overall	512/898 (57%)	415/438 (95%)	0/353 (0%)	97/107 (91%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 57%, i.e. 512 atoms were assigned a chemical shift out of a possible 898. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	256/436 (59%)	173/174 (99%)	0/176 (0%)	83/86 (97%)
Sidechain	212/383 (55%)	201/223 (90%)	0/142 (0%)	11/18 (61%)
Aromatic	44/79 (56%)	41/41 (100%)	0/35 (0%)	3/3 (100%)
Overall	512/898 (57%)	415/438 (95%)	0/353 (0%)	97/107 (91%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
???	C	637	PRO	HG2	0.07	3.48 – 0.38	-6.0
???	C	599	GLN	HB2	0.56	3.30 – 0.80	-6.0
???	C	578	TYR	HB2	0.86	4.76 – 1.06	-5.5

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain C:

