



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 11:25 pm BST

PDB ID : 5HE0  
Title : Bovine GRK2 in complex with Gbetagamma subunits and CCG215022  
Authors : Cato, M.C.; Waninger-Saroni, J.; Tesmer, J.J.G.  
Deposited on : 2016-01-05  
Resolution : 2.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1279 (2.58-2.54)
Clashscore	141614	1327 (2.58-2.54)
Ramachandran outliers	138981	1312 (2.58-2.54)
Sidechain outliers	138945	1312 (2.58-2.54)
RSRZ outliers	127900	1269 (2.58-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	641	<div> <div>10%</div> <div> <div></div> <div>81%</div> <div>14%</div> <div>• •</div> </div> </div>
2	B	339	<div> <div>6%</div> <div> <div></div> <div>85%</div> <div>14%</div> <div>•</div> </div> </div>
3	G	71	<div> <div>8%</div> <div> <div></div> <div>72%</div> <div>11%</div> <div>• 15%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8248 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-adrenergic receptor kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	621	Total	C	N	O	S	0	0	0
			5094	3254	886	918	36			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	670	ALA	SER	engineered mutation	UNP P21146


- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	339	Total	C	N	O	S	0	2	0
			2619	1613	470	513	23			

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	60	Total	C	N	O	S	0	0	0
			462	288	82	89	3			

- Molecule 4 is (4S)-4-{4-fluoro-3-[(pyridin-2-ylmethyl)carbamoyl]phenyl}-N-(1H-indazol-5-yl)-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxamide (three-letter code: 453) (formula: C<sub>26</sub>H<sub>22</sub>FN<sub>7</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0	0
			37	26	1	7	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	17	Total O 17 17	0	0
5	B	19	Total O 19 19	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-adrenergic receptor kinase 1




- Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1



- Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.55Å 239.93Å 208.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.56 29.83 – 2.56	Depositor EDS
% Data completeness (in resolution range)	94.5 (30.00-2.56) 94.6 (29.83-2.56)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 2.57Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.202 , 0.264 0.205 , 0.262	Depositor DCC
$R_{free}$ test set	2316 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	58.3	Xtriage
Anisotropy	0.434	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 31.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8248	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 453

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	0/5210	0.81	1/7001 (0.0%)
2	B	0.74	1/2666 (0.0%)	0.92	5/3613 (0.1%)
3	G	0.64	0/468	0.77	0/631
All	All	0.69	1/8344 (0.0%)	0.84	6/11245 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	297	TRP	CB-CG	-5.22	1.40	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	212	ASP	CB-CG-OD1	8.21	125.69	118.30
2	B	118	ASP	CB-CG-OD2	-6.06	112.84	118.30
1	A	392	ARG	NE-CZ-NH1	5.76	123.18	120.30
2	B	118	ASP	CB-CG-OD1	5.63	123.36	118.30
2	B	212	ASP	CB-CG-OD2	-5.61	113.25	118.30
2	B	247	ASP	CB-CG-OD1	5.40	123.16	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5094	0	5102	56	0
2	B	2619	0	2518	22	0
3	G	462	0	473	5	0
4	A	37	0	22	1	0
5	A	17	0	0	1	0
5	B	19	0	0	0	0
All	All	8248	0	8115	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:ASP:HB3	1:A:213:THR:HG23	1.60	0.81
3:G:54:VAL:HG22	3:G:55:PRO:HD2	1.66	0.74
1:A:565:MET:HE1	1:A:632:LEU:HD13	1.76	0.66
2:B:32:GLN:O	2:B:35:ASN:HB2	1.95	0.66
1:A:361:VAL:HA	1:A:368:TYR:CE2	2.34	0.62
1:A:237:LEU:O	1:A:241:ILE:HD12	2.02	0.59
2:B:180:PHE:CE1	2:B:216:GLY:HA2	2.36	0.59
2:B:51:LEU:HB2	2:B:336:LEU:HB2	1.84	0.58
1:A:470:LEU:HD23	1:A:470:LEU:C	2.26	0.56
1:A:430:GLN:HG3	1:A:435:ARG:O	2.06	0.55
2:B:247:ASP:O	2:B:248:ALA:HB3	2.07	0.54
1:A:644:LYS:NZ	5:A:801:HOH:O	2.40	0.53
1:A:256:CYS:N	1:A:272:ASP:OD2	2.37	0.53
3:G:54:VAL:CG2	3:G:55:PRO:HD2	2.37	0.53
1:A:240:ARG:CZ	1:A:511:LEU:HD21	2.38	0.53
1:A:245:LEU:O	1:A:311:ARG:HD3	2.09	0.52
2:B:205:ASP:O	2:B:206:ALA:HB3	2.10	0.52
1:A:252:PRO:HG2	1:A:253:PHE:CE2	2.46	0.51
1:A:570:ASN:HB2	1:A:573:LEU:HB3	1.93	0.51
2:B:79:LEU:HG	2:B:95:LEU:HD21	1.93	0.51
2:B:261:LEU:HD22	3:G:30:VAL:HG13	1.94	0.50
4:A:701:453:H18	4:A:701:453:OAL	2.12	0.49
1:A:603:MET:HB3	1:A:651:TYR:HA	1.94	0.49
1:A:262:HIS:HB2	1:A:266:LYS:O	2.14	0.48
1:A:570:ASN:CB	1:A:573:LEU:HB3	2.44	0.47
3:G:52:THR:O	3:G:53:PRO:C	2.51	0.47
1:A:125:SER:O	1:A:129:ILE:HD12	2.14	0.47
1:A:385:LEU:HD13	1:A:421:LEU:HD21	1.95	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:68:ARG:O	2:B:84:SER:OG	2.32	0.47
1:A:625:ARG:HD2	1:A:625:ARG:HA	1.77	0.47
1:A:242:MET:HA	1:A:242:MET:HE2	1.97	0.47
1:A:315:TYR:O	1:A:316:ARG:HB2	2.12	0.47
1:A:325:LEU:N	1:A:325:LEU:HD12	2.30	0.47
1:A:457:ASP:OD1	1:A:457:ASP:C	2.53	0.47
2:B:276:VAL:HA	2:B:286:LEU:O	2.15	0.46
1:A:294:MET:C	1:A:294:MET:SD	2.94	0.46
1:A:470:LEU:HD23	1:A:471:ILE:N	2.30	0.46
1:A:315:TYR:OH	1:A:322:ASN:HB3	2.16	0.45
1:A:349:ALA:O	1:A:368:TYR:OH	2.29	0.45
1:A:509:PHE:HB3	1:A:510:PRO:HD3	1.99	0.45
1:A:436:ARG:O	1:A:440:LEU:HD13	2.17	0.45
1:A:566:SER:HA	1:A:577:GLN:O	2.16	0.45
1:A:319:LYS:O	1:A:320:PRO:C	2.55	0.45
1:A:472:PRO:HA	1:A:473:PRO:HD3	1.89	0.44
1:A:525:VAL:O	1:A:529:ILE:HG12	2.17	0.44
2:B:145:TYR:OH	2:B:186:ASP:OD2	2.23	0.44
2:B:104:ALA:HB2	2:B:149:CYS:O	2.17	0.44
1:A:188:MET:HE1	1:A:221:CYS:SG	2.58	0.44
1:A:525:VAL:O	1:A:529:ILE:CG1	2.66	0.43
1:A:69:ARG:HB2	1:A:83:VAL:HG11	2.00	0.43
1:A:147:GLN:N	1:A:148:PRO:HD2	2.34	0.43
1:A:196:ILE:HD12	1:A:196:ILE:H	1.83	0.43
2:B:301:LYS:O	2:B:302:ALA:HB3	2.18	0.43
2:B:49:ARG:HB2	2:B:338:ILE:HG12	1.99	0.43
2:B:34:THR:HG21	2:B:300:LEU:HB3	2.01	0.43
1:A:223:ASP:OD2	1:A:226:ARG:HG3	2.18	0.42
1:A:430:GLN:HA	1:A:430:GLN:OE1	2.19	0.42
2:B:146:LEU:HD11	2:B:159:THR:HB	2.00	0.42
1:A:259:TYR:OH	1:A:517:TRP:HD1	2.02	0.42
1:A:278:ASP:OD2	1:A:321:ALA:HA	2.19	0.42
1:A:80:LYS:N	1:A:81:PRO:CD	2.82	0.42
2:B:291:ASP:C	2:B:291:ASP:OD1	2.58	0.42
1:A:314:VAL:HG12	1:A:370:SER:HA	2.02	0.42
1:A:468:PRO:HA	1:A:469:PRO:HD3	1.85	0.42
1:A:66:LEU:HD23	1:A:66:LEU:HA	1.87	0.42
1:A:80:LYS:HB3	1:A:81:PRO:HD3	2.02	0.42
2:B:269:ILE:HG21	2:B:289:TYR:CE2	2.55	0.41
1:A:279:LEU:HB2	1:A:320:PRO:O	2.20	0.41
2:B:333:ASP:O	2:B:334:SER:CB	2.68	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:ILE:HD13	1:A:380:MET:HE1	2.03	0.41
2:B:99:TRP:CB	2:B:117:LEU:HB2	2.51	0.41
2:B:219:ARG:C	2:B:220:GLN:HG3	2.40	0.41
2:B:166:CYS:HB2	2:B:180:PHE:HB2	2.02	0.41
1:A:377:LEU:O	1:A:381:LEU:HG	2.20	0.41
1:A:65:TYR:O	1:A:66:LEU:C	2.58	0.41
2:B:69:LEU:HA	2:B:82:TRP:O	2.20	0.41
1:A:246:VAL:O	1:A:307:HIS:HE1	2.04	0.40
1:A:199:ARG:HG3	1:A:204:GLU:HG3	2.02	0.40
1:A:305:LEU:HD23	1:A:305:LEU:HA	1.94	0.40
1:A:357:MET:CE	1:A:362:LEU:HD21	2.51	0.40
1:A:509:PHE:N	1:A:510:PRO:CD	2.84	0.40
1:A:648:ARG:HA	1:A:651:TYR:CD2	2.56	0.40
3:G:46:LYS:HD2	3:G:46:LYS:N	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	617/641 (96%)	564 (91%)	46 (8%)	7 (1%)	14	19
2	B	339/339 (100%)	319 (94%)	19 (6%)	1 (0%)	41	50
3	G	58/71 (82%)	53 (91%)	5 (9%)	0	100	100
All	All	1014/1051 (96%)	936 (92%)	70 (7%)	8 (1%)	19	27

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	139	LYS
1	A	212	ASP
2	B	302	ALA

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	370	SER
1	A	625	ARG
1	A	76	LEU
1	A	361	VAL
1	A	495	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	559/574 (97%)	542 (97%)	17 (3%)	41	53
2	B	284/282 (101%)	277 (98%)	7 (2%)	47	61
3	G	49/58 (84%)	45 (92%)	4 (8%)	11	14
All	All	892/914 (98%)	864 (97%)	28 (3%)	40	52

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LYS
1	A	38	SER
1	A	69	ARG
1	A	237	LEU
1	A	240	ARG
1	A	244	SER
1	A	262	HIS
1	A	401	GLU
1	A	412	GLU
1	A	415	ASP
1	A	419	PRO
1	A	440	LEU
1	A	474	ARG
1	A	511	LEU
1	A	595	GLU
1	A	625	ARG
1	A	651	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	31	SER
2	B	61	MET
2	B	105	TYR
2	B	217	MET
2	B	268	ASN
2	B	292	PHE
2	B	294	CYS
3	G	6	THR
3	G	46	LYS
3	G	57	SER
3	G	62	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	133	GLN
1	A	275	ASN
1	A	307	HIS
3	G	11	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	453	A	701	-	39,41,41	2.05	4 (10%)	53,58,58	2.18	18 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsions and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	453	A	701	-	-	5/21/37/37	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	701	453	CAR-CAM	9.16	1.46	1.35
4	A	701	453	CAN-CAM	-4.83	1.47	1.51
4	A	701	453	CAR-NAQ	-4.57	1.32	1.38
4	A	701	453	CAP-NAQ	-2.95	1.32	1.37

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	701	453	NAQ-CAP-NAO	7.81	124.34	116.12
4	A	701	453	CAW-CAX-CAY	-3.91	118.85	123.11
4	A	701	453	CAM-CAN-NAO	3.53	111.91	109.09
4	A	701	453	CAS-CAR-CAM	-3.49	124.13	127.62
4	A	701	453	OAL-CAK-CAM	-3.48	115.42	121.19
4	A	701	453	CAI-CAE-NAA	3.36	135.65	130.19
4	A	701	453	CBF-CBE-NBD	-3.33	105.50	112.71
4	A	701	453	CAC-CAD-CAE	2.92	110.85	104.66
4	A	701	453	CAS-CAR-NAQ	2.83	116.82	113.45
4	A	701	453	OAT-CAP-NAQ	-2.79	116.56	121.82
4	A	701	453	CAR-NAQ-CAP	-2.65	121.64	123.71
4	A	701	453	CAX-CAY-CBB	-2.62	119.03	125.09
4	A	701	453	CBH-NBG-CBF	2.55	120.91	117.42
4	A	701	453	CAY-CBB-NBD	2.49	122.26	117.36
4	A	701	453	CAV-CAW-CAX	2.33	121.86	119.05
4	A	701	453	CAG-NAJ-CAK	-2.12	123.86	127.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	701	453	CAZ-CAY-CAX	2.10	119.21	116.66
4	A	701	453	CBI-CBH-NBG	-2.02	120.13	123.43

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	701	453	CAM-CAK-NAJ-CAG
4	A	701	453	NAO-CAN-CAU-CAZ
4	A	701	453	NAO-CAN-CAU-CAV
4	A	701	453	CAX-CAY-CBB-NBD
4	A	701	453	CAX-CAY-CBB-OBC

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	701	453	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	621/641 (96%)	0.44	64 (10%) 6 10	41, 71, 116, 155	0
2	B	339/339 (100%)	0.29	22 (6%) 18 23	41, 58, 87, 152	0
3	G	60/71 (84%)	0.51	6 (10%) 7 10	47, 65, 116, 138	0
All	All	1020/1051 (97%)	0.39	92 (9%) 9 13	41, 65, 112, 155	0

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	129	ARG	6.6
1	A	475	GLY	6.5
1	A	474	ARG	5.5
1	A	395	LYS	5.5
3	G	46	LYS	5.1
1	A	547	LEU	4.8
1	A	99	GLU	4.8
1	A	394	HIS	4.7
1	A	318	LEU	4.2
1	A	301	ILE	4.0
1	A	408	THR	3.8
3	G	52	THR	3.8
2	B	233[A]	CYS	3.7
1	A	550	GLU	3.6
1	A	499	LEU	3.6
1	A	549	HIS	3.5
1	A	409	MET	3.5
1	A	377	LEU	3.5
1	A	312	PHE	3.4
3	G	61	PHE	3.4
2	B	317	CYS	3.4
1	A	359	PRO	3.4
1	A	333	ILE	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	97	THR	3.3
1	A	543	LYS	3.3
2	B	61	MET	3.3
2	B	60	ALA	3.3
1	A	363	GLN	3.3
2	B	286	LEU	3.3
1	A	334	SER	3.2
1	A	668	PRO	3.2
2	B	318	LEU	3.1
1	A	572	PHE	3.1
3	G	5	ASN	3.0
1	A	393	GLN	3.0
1	A	548	GLY	3.0
1	A	376	SER	2.9
1	A	545	LYS	2.9
2	B	277	SER	2.9
1	A	473	PRO	2.8
2	B	127	LYS	2.8
2	B	103	CYS	2.8
1	A	315	TYR	2.7
1	A	380	MET	2.7
2	B	268	ASN	2.7
1	A	342	PHE	2.7
1	A	298	ALA	2.6
1	A	365	GLY	2.6
1	A	102	LEU	2.6
1	A	126	LYS	2.6
2	B	276	VAL	2.5
2	B	285	LEU	2.5
1	A	118	LEU	2.5
1	A	405	MET	2.5
2	B	131	GLY	2.5
1	A	501	SER	2.5
2	B	130	GLU	2.5
2	B	271[A]	CYS	2.4
3	G	47	GLU	2.4
1	A	297	TYR	2.4
1	A	494	LYS	2.4
1	A	540	LYS	2.4
1	A	618	LYS	2.4
1	A	375	PHE	2.4
1	A	505	LEU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	507	ARG	2.4
1	A	569	GLY	2.3
1	A	402	ILE	2.3
2	B	242	ALA	2.3
1	A	95	LEU	2.3
1	A	498	LEU	2.3
1	A	415	ASP	2.3
1	A	362	LEU	2.3
1	A	404	ARG	2.2
1	A	551	GLU	2.2
1	A	435	ARG	2.2
2	B	232	ILE	2.2
1	A	546	GLN	2.2
1	A	397	LYS	2.2
1	A	336	LEU	2.1
2	B	266	HIS	2.1
1	A	613	GLN	2.1
1	A	378	GLY	2.1
1	A	323	ILE	2.1
1	A	229	MET	2.1
2	B	252	LEU	2.1
2	B	270	ILE	2.1
3	G	63	GLU	2.1
2	B	278	PHE	2.0
1	A	77	GLU	2.0
1	A	364	LYS	2.0
1	A	379	CYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	453	A	701	37/37	0.95	0.17	45,70,81,82	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.