



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 08:43 am BST

PDB ID : 3HGF  
Title : Expression, purification, spectroscopical and crystallographical studies of segments of the nucleotide binding domain of the reticulocyte binding protein Py235 of Plasmodium yoelii  
Authors : Gruber, A.; Manimekalai, M.S.S.; Balakrishna, A.M.; Hunke, C.; Jeyakanthan, J.; Preiser, P.R.; Gruber, G.  
Deposited on : 2009-05-13  
Resolution : 4.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1087 (4.30-3.70)
Clashscore	141614	1148 (4.30-3.70)
Ramachandran outliers	138981	1108 (4.30-3.70)
Sidechain outliers	138945	1099 (4.30-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	107	
1	B	107	
1	C	107	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1375 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rhoptyr protein fragment.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	98	Total	C	N	O	0	0	0
			493	297	98	98			
1	B	89	Total	C	N	O	0	0	0
			446	268	89	89			
1	C	87	Total	C	N	O	0	0	0
			436	262	87	87			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	HIS	-	EXPRESSION TAG	UNP Q7RPU0
A	2	HIS	-	EXPRESSION TAG	UNP Q7RPU0
A	3	HIS	-	EXPRESSION TAG	UNP Q7RPU0
A	4	HIS	-	EXPRESSION TAG	UNP Q7RPU0
A	5	HIS	-	EXPRESSION TAG	UNP Q7RPU0
A	6	HIS	-	EXPRESSION TAG	UNP Q7RPU0
A	7	PRO	-	EXPRESSION TAG	UNP Q7RPU0
A	8	MET	-	EXPRESSION TAG	UNP Q7RPU0
A	9	VAL	-	EXPRESSION TAG	UNP Q7RPU0
B	1	HIS	-	EXPRESSION TAG	UNP Q7RPU0
B	2	HIS	-	EXPRESSION TAG	UNP Q7RPU0
B	3	HIS	-	EXPRESSION TAG	UNP Q7RPU0
B	4	HIS	-	EXPRESSION TAG	UNP Q7RPU0
B	5	HIS	-	EXPRESSION TAG	UNP Q7RPU0
B	6	HIS	-	EXPRESSION TAG	UNP Q7RPU0
B	7	PRO	-	EXPRESSION TAG	UNP Q7RPU0
B	8	MET	-	EXPRESSION TAG	UNP Q7RPU0
B	9	VAL	-	EXPRESSION TAG	UNP Q7RPU0
C	1	HIS	-	EXPRESSION TAG	UNP Q7RPU0
C	2	HIS	-	EXPRESSION TAG	UNP Q7RPU0
C	3	HIS	-	EXPRESSION TAG	UNP Q7RPU0
C	4	HIS	-	EXPRESSION TAG	UNP Q7RPU0
C	5	HIS	-	EXPRESSION TAG	UNP Q7RPU0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	6	HIS	-	EXPRESSION TAG	UNP Q7RPU0
C	7	PRO	-	EXPRESSION TAG	UNP Q7RPU0
C	8	MET	-	EXPRESSION TAG	UNP Q7RPU0
C	9	VAL	-	EXPRESSION TAG	UNP Q7RPU0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Rhoptry protein fragment

Chain A: 




- Molecule 1: Rhoptry protein fragment

Chain B: 



- Molecule 1: Rhoptry protein fragment

Chain C: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.22Å 70.22Å 193.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.00 – 4.00 33.00 – 4.00	Depositor EDS
% Data completeness (in resolution range)	95.4 (33.00-4.00) 95.4 (33.00-4.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.09 (at 3.99Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.339 , 0.373 0.387 , 0.320	Depositor DCC
$R_{free}$ test set	120 reflections (4.60%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	133.2	Xtriage
Anisotropy	0.466	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	1375	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/493	0.45	0/687
1	B	0.27	0/445	0.45	0/620
1	C	0.27	0/434	0.45	0/603
All	All	0.27	0/1372	0.45	0/1910

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	493	0	212	8	0
1	B	446	0	188	3	0
1	C	436	0	183	3	0
All	All	1375	0	583	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:99:GLN:HA	1:C:103:ALA:HB2	1.83	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:VAL:C	1:A:11:GLU:H	2.12	0.53
1:A:36:LEU:C	1:A:38:ASN:H	2.14	0.51
1:B:96:ILE:C	1:B:98:GLU:H	2.15	0.49
1:B:79:SER:C	1:B:81:ASN:H	2.18	0.47
1:A:27:LYS:C	1:A:29:ILE:H	2.21	0.44
1:A:11:GLU:C	1:A:13:GLU:H	2.21	0.43
1:A:46:ASP:O	1:A:47:LYS:C	2.57	0.43
1:C:97:LYS:O	1:C:101:PRO:HD2	2.19	0.42
1:A:98:GLU:C	1:A:101:PRO:HD2	2.40	0.42
1:C:100:SER:CB	1:C:101:PRO:HD3	2.50	0.42
1:A:104:GLU:C	1:A:106:ASN:H	2.23	0.41
1:B:93:LEU:C	1:B:95:GLU:H	2.23	0.41
1:A:61:GLY:O	1:A:63:SER:N	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/107 (88%)	70 (74%)	14 (15%)	10 (11%)	0	7
1	B	85/107 (79%)	67 (79%)	17 (20%)	1 (1%)	13	49
1	C	81/107 (76%)	63 (78%)	14 (17%)	4 (5%)	2	22
All	All	260/321 (81%)	200 (77%)	45 (17%)	15 (6%)	1	19

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	20	VAL
1	A	43	ILE
1	A	62	LYS
1	A	84	LYS

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	C	101	PRO
1	A	10	LYS
1	A	37	LEU
1	A	92	ASP
1	B	41	ALA
1	C	66	LYS
1	C	104	GLU
1	A	28	TYR
1	A	105	MET
1	C	100	SER
1	A	12	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	2/104 (2%)	2 (100%)	0	100	100
1	B	1/104 (1%)	1 (100%)	0	100	100
1	C	1/104 (1%)	1 (100%)	0	100	100
All	All	4/312 (1%)	4 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.