



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 02:57 am BST

PDB ID : 3HHD
Title : Structure of the Human Fatty Acid Synthase KS-MAT Didomain as a Framework for Inhibitor Design.
Authors : Pappenberger, G.M.; Benz, J.; Thoma, R.; Rudolph, M.G.
Deposited on : 2009-05-15
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	965	<div> <div>2%</div> <div> <div></div> <div>72%</div> <div>15%</div> <div>•</div> <div>12%</div> </div> </div>
1	B	965	<div> <div>4%</div> <div> <div></div> <div>71%</div> <div>15%</div> <div>•</div> <div>12%</div> </div> </div>
1	C	965	<div> <div>2%</div> <div> <div></div> <div>73%</div> <div>14%</div> <div>•</div> <div>11%</div> </div> </div>
1	D	965	<div> <div>3%</div> <div> <div></div> <div>72%</div> <div>14%</div> <div>•</div> <div>12%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 27836 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fatty acid synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	852	Total	C	N	O	S	0	2	0
			6497	4119	1144	1200	34			
1	B	852	Total	C	N	O	S	0	3	0
			6505	4125	1147	1199	34			
1	C	855	Total	C	N	O	S	0	1	0
			6512	4127	1147	1204	34			
1	D	854	Total	C	N	O	S	0	2	0
			6512	4129	1146	1203	34			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	SER	-	EXPRESSION TAG	UNP P49327
A	0	THR	-	EXPRESSION TAG	UNP P49327
A	1	GLY	-	EXPRESSION TAG	UNP P49327
B	-1	SER	-	EXPRESSION TAG	UNP P49327
B	0	THR	-	EXPRESSION TAG	UNP P49327
B	1	GLY	-	EXPRESSION TAG	UNP P49327
C	-1	SER	-	EXPRESSION TAG	UNP P49327
C	0	THR	-	EXPRESSION TAG	UNP P49327
C	1	GLY	-	EXPRESSION TAG	UNP P49327
D	-1	SER	-	EXPRESSION TAG	UNP P49327
D	0	THR	-	EXPRESSION TAG	UNP P49327
D	1	GLY	-	EXPRESSION TAG	UNP P49327

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	561	Total 561	O 561	0	0
3	B	371	Total 371	O 371	0	0
3	C	437	Total 437	O 437	0	0
3	D	439	Total 439	O 439	0	0





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	86.62Å 91.16Å 132.11Å 73.84° 86.83° 62.54°	Depositor
Resolution (Å)	45.64 – 2.15 45.64 – 2.15	Depositor EDS
% Data completeness (in resolution range)	97.3 (45.64-2.15) 97.5 (45.64-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 2.14Å)	Xtriage
Refinement program	BUSTER-TNT 2.5.1	Depositor
R, R_{free}	0.169 , 0.211 0.171 , 0.214	Depositor DCC
R_{free} test set	9152 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	23.0	Xtriage
Anisotropy	0.518	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	27836	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	1/6662 (0.0%)	0.68	0/9058
1	B	0.61	1/6673 (0.0%)	0.66	1/9074 (0.0%)
1	C	0.63	2/6674 (0.0%)	0.67	1/9075 (0.0%)
1	D	0.62	1/6678 (0.0%)	0.68	2/9083 (0.0%)
All	All	0.64	5/26687 (0.0%)	0.67	4/36290 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	593	CYS	CB-SG	-7.10	1.70	1.82
1	A	593	CYS	CB-SG	-5.64	1.72	1.81
1	D	223	CYS	CB-SG	-5.36	1.73	1.81
1	B	593	CYS	CB-SG	-5.31	1.73	1.81
1	C	161	CYS	CB-SG	-5.06	1.73	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	137	ARG	NE-CZ-NH1	-6.05	117.28	120.30
1	B	333	GLU	C-N-CD	5.47	139.88	128.40
1	C	468	ARG	NE-CZ-NH2	-5.34	117.63	120.30
1	D	802	ARG	NE-CZ-NH2	-5.03	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6497	0	6477	106	0
1	B	6505	0	6491	119	0
1	C	6512	0	6491	94	0
1	D	6512	0	6487	117	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	561	0	0	8	0
3	B	371	0	0	11	0
3	C	437	0	0	13	0
3	D	439	0	0	15	0
All	All	27836	0	25946	427	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 427 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:656:ALA:HB3	1:B:657:PRO:HD3	1.45	0.95
1:D:656:ALA:HB3	1:D:657:PRO:HD3	1.50	0.94
1:A:42:ALA:HB1	1:A:49:ARG:HD2	1.56	0.87
1:A:67:VAL:HG11	1:A:75:MET:HE1	1.56	0.85
1:D:556:LEU:HD23	1:D:582:LEU:HD23	1.62	0.81

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	852/965 (88%)	837 (98%)	14 (2%)	1 (0%)	51	53
1	B	853/965 (88%)	836 (98%)	16 (2%)	1 (0%)	51	53
1	C	854/965 (88%)	840 (98%)	13 (2%)	1 (0%)	51	53
1	D	854/965 (88%)	837 (98%)	16 (2%)	1 (0%)	51	53
All	All	3413/3860 (88%)	3350 (98%)	59 (2%)	4 (0%)	51	53

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	333	GLU
1	B	333	GLU
1	C	333	GLU
1	D	333	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	698/792 (88%)	666 (95%)	32 (5%)	27	23
1	B	698/792 (88%)	658 (94%)	40 (6%)	20	16
1	C	699/792 (88%)	663 (95%)	36 (5%)	23	19
1	D	699/792 (88%)	658 (94%)	41 (6%)	19	15
All	All	2794/3168 (88%)	2645 (95%)	149 (5%)	22	19

5 of 149 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	746	GLN
1	C	343	LYS
1	D	676	ARG
1	B	773	ARG
1	C	169	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	483	GLN
1	C	136	GLN
1	D	331	HIS
1	B	723	HIS
1	A	852	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	852/965 (88%)	-0.32	19 (2%) 62 69	8, 22, 56, 78	0
1	B	852/965 (88%)	-0.11	34 (3%) 38 47	15, 31, 63, 91	0
1	C	855/965 (88%)	-0.27	15 (1%) 68 75	13, 28, 60, 78	0
1	D	854/965 (88%)	-0.24	28 (3%) 46 55	10, 27, 60, 85	0
All	All	3413/3860 (88%)	-0.24	96 (2%) 53 62	8, 27, 60, 91	0

The worst 5 of 96 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	415	ALA	7.8
1	B	416	PRO	7.3
1	A	415	ALA	6.8
1	D	415	ALA	6.6
1	A	414	PRO	6.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	D	964	1/1	0.91	0.11	66,66,66,66	0
2	CL	C	964	1/1	0.94	0.16	76,76,76,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.