



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2021 – 02:44 AM EDT

PDB ID : 3HI7  
Title : Crystal structure of human diamine oxidase  
Authors : McGrath, A.P.; Guss, J.M.  
Deposited on : 2009-05-19  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	731	<div> <div>3%</div> <div>92%</div> <div>5%</div> </div>
1	B	731	<div> <div>3%</div> <div>92%</div> <div>5%</div> </div>
2	C	3	<div> <div>33%</div> <div>67%</div> </div>
3	D	2	<div> <div>100%</div> </div>
3	E	2	<div> <div>100%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	2	 50% 50%
3	G	2	 100%
3	H	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	E	2	-	-	-	X



In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Amiloride-sensitive amine oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	715	Total 5843	C 3780	N 994	O 1048	S 21	0	25	0
1	B	710	Total 5789	C 3748	N 984	O 1036	S 21	0	22	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	ARG	PRO	engineered mutation	UNP P19801
B	21	ARG	PRO	engineered mutation	UNP P19801

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cu	0	0
			1	1		
4	B	1	Total	Cu	0	0
			1	1		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Ca	0	0
			2	2		
5	B	2	Total	Ca	0	0
			2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	568	Total	O	0	0
			568	568		
7	B	568	Total	O	0	0
			568	568		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Amiloride-sensitive amine oxidase



- Molecule 1: Amiloride-sensitive amine oxidase



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1  
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1  
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%


MAG1  
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

MAG1  
MAG2

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%

MAG1  
MAG2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.51Å 94.78Å 196.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.57 – 1.80 45.02 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.8 (46.57-1.80) 98.9 (45.02-1.80)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.15 (at 1.79Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.161 , 0.189 0.167 , 0.193	Depositor DCC
$R_{free}$ test set	8007 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.7	Xtriage
Anisotropy	0.096	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 58.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.012 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12989	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU, GOL, CA, BMA, TPQ, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	0/6079	0.67	0/8305
1	B	0.56	2/6016 (0.0%)	0.66	1/8220 (0.0%)
All	All	0.55	2/12095 (0.0%)	0.66	1/16525 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	123[A]	CYS	CB-SG	-6.02	1.72	1.82
1	B	123[B]	CYS	CB-SG	-6.02	1.72	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	205	ARG	NE-CZ-NH2	-6.47	117.07	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5843	0	5543	33	0
1	B	5789	0	5491	30	0
2	C	39	0	34	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	28	0	25	0	0
3	E	28	0	25	0	0
3	F	28	0	25	0	0
3	G	28	0	25	0	0
3	H	28	0	25	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	18	0	24	0	0
6	B	18	0	24	0	0
7	A	568	0	0	1	0
7	B	568	0	0	0	0
All	All	12989	0	11241	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:302[A]:LEU:HD23	1:B:302[A]:LEU:HD23	1.50	0.89
1:A:598[B]:HIS:CD2	1:B:598[B]:HIS:CD2	2.65	0.84
1:B:449[B]:VAL:HG21	1:B:468:ILE:CG2	2.10	0.81
1:A:598[B]:HIS:HD2	1:B:598[B]:HIS:NE2	1.84	0.76
1:A:328:LEU:HD21	1:A:382:THR:HG21	1.66	0.75
1:A:598[B]:HIS:HD2	1:B:598[B]:HIS:CD2	2.11	0.69
1:A:449[B]:VAL:HG21	1:A:468:ILE:CG2	2.24	0.67
1:A:598[B]:HIS:CD2	1:B:598[B]:HIS:NE2	2.62	0.66
1:B:449[B]:VAL:HG21	1:B:468:ILE:HG23	1.82	0.62
1:B:449[B]:VAL:CG2	1:B:468:ILE:CG2	2.78	0.60
1:B:175[A]:GLN:HE21	1:B:175[A]:GLN:HA	1.66	0.60
1:B:449[B]:VAL:CG2	1:B:468:ILE:HG23	2.31	0.60
1:B:328:LEU:HD21	1:B:382:THR:HG21	1.84	0.59
1:A:719[B]:ARG:HD3	1:A:723:PRO:O	2.03	0.59
1:B:476:GLU:HB2	1:B:695[A]:LEU:HD23	1.87	0.57
1:A:662[B]:ASN:OD1	7:A:1065:HOH:O	2.17	0.57
1:A:449[B]:VAL:CG2	1:A:468:ILE:CG2	2.85	0.55
1:B:356[B]:VAL:HG12	1:B:372:LEU:HG	1.91	0.53
1:A:356[B]:VAL:HG22	1:A:513:LEU:HB2	1.90	0.52
1:A:302[A]:LEU:HD23	1:B:302[A]:LEU:CD2	2.32	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:160:LYS:HE2	1:B:175[B]:GLN:NE2	2.26	0.50
1:A:421[A]:MET:O	1:A:421[A]:MET:HG3	2.12	0.50
1:A:66:LYS:HE3	1:A:68[B]:THR:HG21	1.94	0.49
1:A:598[B]:HIS:NE2	1:B:598[B]:HIS:CD2	2.80	0.49
1:A:528[B]:SER:OG	1:A:580:THR:OG1	2.28	0.48
1:A:216:GLY:HA2	1:A:489:PHE:CG	2.48	0.48
1:B:57:GLN:O	1:B:68[B]:THR:HG22	2.13	0.48
1:A:719[B]:ARG:HD2	1:A:725:TYR:HB2	1.95	0.47
1:A:57:GLN:O	1:A:68[B]:THR:HG22	2.13	0.47
1:B:479[B]:MET:SD	1:B:514:VAL:HG21	2.54	0.47
1:B:66:LYS:HE3	1:B:68[B]:THR:HG21	1.96	0.47
1:A:362:HIS:CG	1:A:494:GLY:HA2	2.50	0.46
1:B:461:TPQ:H6	1:B:482:THR:O	2.15	0.46
1:A:302[A]:LEU:CD2	1:B:302[A]:LEU:HD23	2.35	0.46
1:A:163:HIS:ND1	1:A:175:GLN:HG2	2.31	0.46
1:B:359:TYR:CD1	1:B:510:HIS:HB3	2.50	0.45
2:C:2:NAG:H4	2:C:3:BMA:O2	2.16	0.45
1:A:359:TYR:CD1	1:A:510:HIS:HB3	2.52	0.44
1:A:356[B]:VAL:HG12	1:A:372:LEU:HG	1.99	0.44
1:A:115:ALA:HB3	1:A:126:ARG:HG2	1.98	0.44
1:B:160:LYS:HE2	1:B:175[B]:GLN:HE22	1.83	0.43
1:A:449[B]:VAL:HG21	1:A:468:ILE:HG22	2.00	0.43
1:A:681:ASP:OD1	1:A:685[B]:THR:HG22	2.18	0.43
1:B:216:GLY:HA2	1:B:489:PHE:CG	2.54	0.42
1:B:362:HIS:CG	1:B:494:GLY:HA2	2.55	0.42
1:B:237:TRP:CH2	1:B:363:THR:HG22	2.55	0.41
1:A:449[B]:VAL:CG2	1:A:468:ILE:HG23	2.51	0.41
1:A:461:TPQ:H6	1:A:482:THR:O	2.20	0.41
1:A:675:HIS:HE1	1:A:685[B]:THR:HG21	1.85	0.41
1:B:356[B]:VAL:CG1	1:B:372:LEU:HG	2.49	0.41
1:A:302[B]:LEU:HD11	1:A:421[B]:MET:HE3	2.03	0.41
1:B:681:ASP:OD1	1:B:685[B]:THR:HG22	2.21	0.41
1:B:449[B]:VAL:HG22	1:B:468:ILE:HG23	2.03	0.40
1:A:356[B]:VAL:CG2	1:A:513:LEU:HB2	2.51	0.40
1:B:66:LYS:HG3	1:B:68[B]:THR:HG23	2.02	0.40
1:A:563[B]:ARG:HA	1:A:563[B]:ARG:HD3	1.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	735/731 (100%)	714 (97%)	21 (3%)	0	100	100
1	B	727/731 (100%)	708 (97%)	18 (2%)	1 (0%)	51	36
All	All	1462/1462 (100%)	1422 (97%)	39 (3%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	459	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	619/634 (98%)	614 (99%)	5 (1%)	81	78
1	B	615/634 (97%)	607 (99%)	8 (1%)	69	62
All	All	1234/1268 (97%)	1221 (99%)	13 (1%)	78	68

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	288	ARG
1	A	337	LEU
1	A	429	ARG
1	A	439	PHE
1	A	711	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	175[A]	GLN
1	B	175[B]	GLN
1	B	288	ARG
1	B	429	ARG
1	B	439	PHE
1	B	525[A]	THR
1	B	525[B]	THR
1	B	733	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	TPQ	B	461	4,1	13,14,15	1.35	2 (15%)	15,19,21	1.32	2 (13%)
1	TPQ	A	461	4,1	13,14,15	1.39	1 (7%)	15,19,21	1.08	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPQ	B	461	4,1	-	4/5/22/24	0/1/1/1
1	TPQ	A	461	4,1	-	5/5/22/24	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	461	TPQ	O4-C4	-3.50	1.24	1.34
1	B	461	TPQ	O4-C4	-2.79	1.26	1.34
1	B	461	TPQ	C3-C4	2.50	1.39	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	461	TPQ	CB-CA-C	-3.07	105.72	111.47
1	B	461	TPQ	C3-C4-C5	-2.88	118.28	121.26
1	B	461	TPQ	CB-CA-C	-2.80	106.21	111.47

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	461	TPQ	C-CA-CB-C1
1	A	461	TPQ	O-C-CA-CB
1	A	461	TPQ	C2-C1-CB-CA
1	B	461	TPQ	C-CA-CB-C1
1	B	461	TPQ	O-C-CA-CB
1	B	461	TPQ	C2-C1-CB-CA
1	A	461	TPQ	N-CA-CB-C1
1	B	461	TPQ	N-CA-CB-C1
1	A	461	TPQ	C6-C1-CB-CA

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	461	TPQ	1	0
1	A	461	TPQ	1	0

## 5.5 Carbohydrates

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	C	1	2,1	14,14,15	0.52	0	17,19,21	0.76	0
2	NAG	C	2	2	14,14,15	0.53	0	17,19,21	1.06	1 (5%)
2	BMA	C	3	2	11,11,12	0.77	0	15,15,17	1.13	1 (6%)
3	NAG	D	1	3,1	14,14,15	0.42	0	17,19,21	1.02	1 (5%)
3	NAG	D	2	3	14,14,15	0.52	0	17,19,21	0.90	1 (5%)
3	NAG	E	1	3,1	14,14,15	0.51	0	17,19,21	0.91	0
3	NAG	E	2	3	14,14,15	0.45	0	17,19,21	0.91	0
3	NAG	F	1	3,1	14,14,15	0.60	0	17,19,21	1.50	3 (17%)
3	NAG	F	2	3	14,14,15	0.59	0	17,19,21	0.91	0
3	NAG	G	1	3,1	14,14,15	0.54	0	17,19,21	0.87	0
3	NAG	G	2	3	14,14,15	0.63	0	17,19,21	0.90	0
3	NAG	H	1	3,1	14,14,15	0.45	0	17,19,21	1.20	1 (5%)
3	NAG	H	2	3	14,14,15	0.71	0	17,19,21	1.18	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	BMA	C	3	2	-	2/2/19/22	0/1/1/1
3	NAG	D	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	NAG	E	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1
3	NAG	F	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
3	NAG	G	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	0/6/23/26	0/1/1/1
3	NAG	H	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	H	2	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (10) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	1	NAG	C1-O5-C5	3.52	116.97	112.19
3	F	1	NAG	C4-C3-C2	2.87	115.22	111.02
3	D	1	NAG	O5-C1-C2	-2.77	106.92	111.29
3	F	1	NAG	C8-C7-N2	2.68	120.64	116.10
3	F	1	NAG	O5-C5-C6	2.42	111.00	107.20
3	D	2	NAG	C1-O5-C5	2.30	115.30	112.19
3	H	2	NAG	C1-O5-C5	-2.23	109.16	112.19
2	C	2	NAG	C4-C3-C2	2.11	114.11	111.02
2	C	3	BMA	C1-C2-C3	2.10	112.25	109.67
3	H	2	NAG	O5-C5-C6	2.08	110.46	107.20

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	3	BMA	C4-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6
2	C	3	BMA	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
3	E	2	NAG	O5-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3	BMA	1	0
2	C	2	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	GOL	A	804	-	5,5,5	0.30	0	5,5,5	0.29	0
6	GOL	B	804	-	5,5,5	0.32	0	5,5,5	0.31	0
6	GOL	B	806	-	5,5,5	0.43	0	5,5,5	0.43	0
6	GOL	A	805	-	5,5,5	0.31	0	5,5,5	0.48	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	GOL	A	806	-	5,5,5	0.30	0	5,5,5	0.22	0
6	GOL	B	805	-	5,5,5	0.35	0	5,5,5	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	804	-	-	0/4/4/4	-
6	GOL	B	804	-	-	1/4/4/4	-
6	GOL	B	806	-	-	3/4/4/4	-
6	GOL	A	805	-	-	0/4/4/4	-
6	GOL	A	806	-	-	2/4/4/4	-
6	GOL	B	805	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	806	GOL	C1-C2-C3-O3
6	B	805	GOL	O1-C1-C2-C3
6	B	806	GOL	O1-C1-C2-C3
6	B	806	GOL	C1-C2-C3-O3
6	A	806	GOL	O2-C2-C3-O3
6	B	806	GOL	O1-C1-C2-O2
6	B	805	GOL	O1-C1-C2-O2
6	B	805	GOL	C1-C2-C3-O3
6	B	804	GOL	C1-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	714/731 (97%)	0.02	23 (3%)	47 41	7, 15, 28, 43	2 (0%)
1	B	709/731 (96%)	0.01	20 (2%)	53 47	8, 16, 27, 39	1 (0%)
All	All	1423/1462 (97%)	0.01	43 (3%)	50 44	7, 15, 27, 43	3 (0%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	267	LEU	6.5
1	A	27	PRO	5.4
1	A	721	ASN	4.5
1	A	123[A]	CYS	4.3
1	B	721	ASN	4.3
1	B	720	ASP	4.3
1	A	49	TRP	4.2
1	A	266	PRO	4.1
1	A	720	ASP	3.8
1	B	723	PRO	3.6
1	B	279	PRO	3.5
1	B	278	GLU	3.2
1	B	123[A]	CYS	3.2
1	B	722	GLY	3.2
1	B	129	SER	3.1
1	A	28	ARG	3.0
1	A	279	PRO	2.8
1	A	257	GLU	2.7
1	B	598[A]	HIS	2.7
1	B	356[A]	VAL	2.6
1	B	730	ILE	2.6
1	B	89	GLY	2.5
1	A	278	GLU	2.4
1	B	571	ARG	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	28	ARG	2.4
1	A	598[A]	HIS	2.4
1	A	265	ASP	2.3
1	B	465	TRP	2.3
1	A	152[A]	TYR	2.3
1	A	722	GLY	2.3
1	A	571	ARG	2.2
1	A	295	ILE	2.2
1	B	92	HIS	2.2
1	A	572	LYS	2.2
1	B	49	TRP	2.2
1	A	50	SER	2.1
1	B	355	ALA	2.1
1	A	723	PRO	2.1
1	A	378	LEU	2.1
1	B	378	LEU	2.0
1	B	572	LYS	2.0
1	A	734	ARG	2.0
1	A	256	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPQ	A	461	14/15	0.97	0.14	11,14,17,21	2
1	TPQ	B	461	14/15	0.98	0.10	9,12,14,20	2

## 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	E	2	14/15	0.63	0.42	43,46,49,50	0
3	NAG	H	2	14/15	0.65	0.30	28,30,32,34	14

*Continued on next page...*

*Continued from previous page...*



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BMA	C	3	11/12	0.75	0.27	44,46,46,48	0
3	NAG	D	2	14/15	0.76	0.30	35,38,42,43	0
3	NAG	G	2	14/15	0.85	0.28	32,35,37,38	0
3	NAG	F	2	14/15	0.85	0.28	28,32,38,39	0
2	NAG	C	2	14/15	0.87	0.21	30,34,41,41	0
3	NAG	H	1	14/15	0.90	0.17	22,25,31,31	0
3	NAG	F	1	14/15	0.90	0.14	20,24,31,37	0
3	NAG	E	1	14/15	0.93	0.16	22,26,29,36	0
3	NAG	G	1	14/15	0.94	0.14	18,21,26,27	0
3	NAG	D	1	14/15	0.95	0.16	21,23,27,30	0
2	NAG	C	1	14/15	0.95	0.11	22,27,30,31	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain F:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	GOL	A	806	6/6	0.84	0.16	27,33,35,35	0
6	GOL	B	806	6/6	0.90	0.23	33,38,39,40	0
6	GOL	B	805	6/6	0.91	0.11	27,32,33,34	0
6	GOL	A	805	6/6	0.97	0.14	20,20,20,22	0
6	GOL	A	804	6/6	0.97	0.09	14,15,18,20	0
6	GOL	B	804	6/6	0.98	0.12	17,20,22,24	0
5	CA	B	803	1/1	0.99	0.04	19,19,19,19	0
5	CA	A	802	1/1	1.00	0.04	8,8,8,8	0
5	CA	A	803	1/1	1.00	0.05	8,8,8,8	0
5	CA	B	802	1/1	1.00	0.06	14,14,14,14	0
4	CU	A	801	1/1	1.00	0.07	13,13,13,13	0
4	CU	B	801	1/1	1.00	0.08	11,11,11,11	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.