



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 07:00 am BST

PDB ID : 5HKD  
Title : Bacterial sodium channel neck 7G mutant  
Authors : Rohaim, A.; Minor, D.L.  
Deposited on : 2016-01-13  
Resolution : 3.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1212 (4.00-3.60)
Clashscore	141614	1288 (4.00-3.60)
Ramachandran outliers	138981	1243 (4.00-3.60)
Sidechain outliers	138945	1237 (4.00-3.60)
RSRZ outliers	127900	1121 (4.00-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	
1	B	152	
1	C	152	
1	D	152	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ion transport protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	121	Total	C	N	O	S	0	0	0
			929	625	138	160	6			
1	B	116	Total	C	N	O	S	0	0	0
			913	617	137	153	6			
1	C	123	Total	C	N	O	S	0	0	0
			924	624	139	155	6			
1	D	123	Total	C	N	O	S	0	0	0
			928	629	140	153	6			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	137	GLY	-	expression tag	UNP Q0ABW0
A	138	PRO	-	expression tag	UNP Q0ABW0
A	139	SER	-	expression tag	UNP Q0ABW0
A	140	SER	-	expression tag	UNP Q0ABW0
A	141	PRO	-	expression tag	UNP Q0ABW0
A	142	SER	-	expression tag	UNP Q0ABW0
A	248	GLY	ALA	engineered mutation	UNP Q0ABW0
A	249	GLY	GLU	engineered mutation	UNP Q0ABW0
A	250	GLY	ASP	engineered mutation	UNP Q0ABW0
A	251	GLY	ALA	engineered mutation	UNP Q0ABW0
A	252	GLY	LYS	engineered mutation	UNP Q0ABW0
A	253	GLY	ARG	engineered mutation	UNP Q0ABW0
A	254	GLY	ILE	engineered mutation	UNP Q0ABW0
B	137	GLY	-	expression tag	UNP Q0ABW0
B	138	PRO	-	expression tag	UNP Q0ABW0
B	139	SER	-	expression tag	UNP Q0ABW0
B	140	SER	-	expression tag	UNP Q0ABW0
B	141	PRO	-	expression tag	UNP Q0ABW0
B	142	SER	-	expression tag	UNP Q0ABW0
B	248	GLY	ALA	engineered mutation	UNP Q0ABW0
B	249	GLY	GLU	engineered mutation	UNP Q0ABW0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	250	GLY	ASP	engineered mutation	UNP Q0ABW0
B	251	GLY	ALA	engineered mutation	UNP Q0ABW0
B	252	GLY	LYS	engineered mutation	UNP Q0ABW0
B	253	GLY	ARG	engineered mutation	UNP Q0ABW0
B	254	GLY	ILE	engineered mutation	UNP Q0ABW0
C	137	GLY	-	expression tag	UNP Q0ABW0
C	138	PRO	-	expression tag	UNP Q0ABW0
C	139	SER	-	expression tag	UNP Q0ABW0
C	140	SER	-	expression tag	UNP Q0ABW0
C	141	PRO	-	expression tag	UNP Q0ABW0
C	142	SER	-	expression tag	UNP Q0ABW0
C	248	GLY	ALA	engineered mutation	UNP Q0ABW0
C	249	GLY	GLU	engineered mutation	UNP Q0ABW0
C	250	GLY	ASP	engineered mutation	UNP Q0ABW0
C	251	GLY	ALA	engineered mutation	UNP Q0ABW0
C	252	GLY	LYS	engineered mutation	UNP Q0ABW0
C	253	GLY	ARG	engineered mutation	UNP Q0ABW0
C	254	GLY	ILE	engineered mutation	UNP Q0ABW0
D	137	GLY	-	expression tag	UNP Q0ABW0
D	138	PRO	-	expression tag	UNP Q0ABW0
D	139	SER	-	expression tag	UNP Q0ABW0
D	140	SER	-	expression tag	UNP Q0ABW0
D	141	PRO	-	expression tag	UNP Q0ABW0
D	142	SER	-	expression tag	UNP Q0ABW0
D	248	GLY	ALA	engineered mutation	UNP Q0ABW0
D	249	GLY	GLU	engineered mutation	UNP Q0ABW0
D	250	GLY	ASP	engineered mutation	UNP Q0ABW0
D	251	GLY	ALA	engineered mutation	UNP Q0ABW0
D	252	GLY	LYS	engineered mutation	UNP Q0ABW0
D	253	GLY	ARG	engineered mutation	UNP Q0ABW0
D	254	GLY	ILE	engineered mutation	UNP Q0ABW0


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ca 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	O	0	0
			2	2		





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	137.10Å 137.99Å 173.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 3.80 14.99 – 3.80	Depositor EDS
% Data completeness (in resolution range)	96.0 (15.00-3.80) 97.9 (14.99-3.80)	Depositor EDS
$R_{merge}$	0.66	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.23 (at 3.77Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, $R_{free}$	0.273 , 0.312 0.270 , 0.305	Depositor DCC
$R_{free}$ test set	795 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	129.7	Xtriage
Anisotropy	0.153	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.22 , 155.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.39$ , $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	0.100 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3697	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	169.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.67	0/953	0.80	1/1301 (0.1%)
1	B	0.64	1/939 (0.1%)	1.04	3/1281 (0.2%)
1	C	0.66	1/948 (0.1%)	1.10	3/1295 (0.2%)
1	D	0.62	0/954	1.07	3/1305 (0.2%)
All	All	0.65	2/3794 (0.1%)	1.01	10/5182 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3
1	C	0	2
1	D	0	3
All	All	0	8

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	279	ASP	CB-CG	5.38	1.63	1.51
1	C	152	TRP	CB-CG	5.01	1.59	1.50

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	173	ALA	CB-CA-C	-25.48	71.88	110.10
1	B	173	ALA	CB-CA-C	-24.76	72.95	110.10
1	D	173	ALA	CB-CA-C	-24.28	73.68	110.10
1	D	173	ALA	C-N-CA	8.56	143.10	121.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	173	ALA	C-N-CA	8.26	142.34	121.70

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	173	ALA	Peptide
1	B	276	SER	Peptide
1	B	278	VAL	Peptide
1	C	173	ALA	Peptide
1	C	281	LEU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	929	0	891	57	0
1	B	913	0	889	54	0
1	C	924	0	882	42	0
1	D	928	0	884	48	0
2	B	1	0	0	0	0
3	A	2	0	0	0	0
All	All	3697	0	3546	167	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 167 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:281:LEU:HD22	1:D:282:GLU:HA	1.64	0.80
1:D:205:ARG:HB2	1:D:206:PRO:HD3	1.66	0.78
1:A:267:MET:HB2	1:B:268:LEU:HD21	1.66	0.77
1:B:205:ARG:HB2	1:B:206:PRO:HD3	1.68	0.76
1:B:260:ALA:O	1:B:263:GLU:HG2	1.85	0.74

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	117/152 (77%)	97 (83%)	19 (16%)	1 (1%)	17	54
1	B	112/152 (74%)	91 (81%)	20 (18%)	1 (1%)	17	54
1	C	119/152 (78%)	94 (79%)	25 (21%)	0	100	100
1	D	119/152 (78%)	97 (82%)	22 (18%)	0	100	100
All	All	467/608 (77%)	379 (81%)	86 (18%)	2 (0%)	34	70

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	173	ALA
1	B	258	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	92/126 (73%)	81 (88%)	11 (12%)	5	25
1	B	92/126 (73%)	80 (87%)	12 (13%)	4	23
1	C	88/126 (70%)	79 (90%)	9 (10%)	7	31
1	D	88/126 (70%)	78 (89%)	10 (11%)	5	28
All	All	360/504 (71%)	318 (88%)	42 (12%)	5	27

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	230	LEU
1	C	161	PHE
1	D	219	VAL
1	B	257	GLU
1	B	266	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	192	GLN
1	C	192	GLN
1	D	192	GLN
1	D	261	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	121/152 (79%)	-0.82	1 (0%) 86 81	84, 151, 249, 285	0
1	B	116/152 (76%)	-0.78	1 (0%) 84 79	93, 162, 262, 296	0
1	C	123/152 (80%)	-0.78	1 (0%) 86 81	78, 160, 259, 321	0
1	D	123/152 (80%)	-0.83	1 (0%) 86 81	87, 168, 268, 371	0
All	All	483/608 (79%)	-0.80	4 (0%) 86 81	78, 163, 266, 371	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	242	GLN	6.2
1	A	242	GLN	5.2
1	D	242	GLN	4.0
1	B	242	GLN	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	B	301	1/1	0.97	0.34	30,30,30,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.