



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 07:15 am BST

PDB ID : 6HND
Title : Crystal structure of the aromatic aminotransferase Aro9 from *C. Albicans*
Authors : Kiliszek, A.; Rząd, K.; Rypniewski, W.; Milewski, S.; Gabriel, I.
Deposited on : 2018-09-14
Resolution : 2.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2391 (2.26-2.22)
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	529	 2% 82% 6% 12%
1	B	529	 2% 80% 8% 11%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8124 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aromatic-amino-acid:2-oxoglutarate transaminase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	467	Total	C	N	O	S	0	0	0
			3767	2442	617	702	6			
1	B	470	Total	C	N	O	S	0	0	0
			3790	2456	622	706	6			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	524	HIS	-	expression tag	UNP A0A1D8PMC5
A	525	HIS	-	expression tag	UNP A0A1D8PMC5
A	526	HIS	-	expression tag	UNP A0A1D8PMC5
A	527	HIS	-	expression tag	UNP A0A1D8PMC5
A	528	HIS	-	expression tag	UNP A0A1D8PMC5
A	529	HIS	-	expression tag	UNP A0A1D8PMC5
B	524	HIS	-	expression tag	UNP A0A1D8PMC5
B	525	HIS	-	expression tag	UNP A0A1D8PMC5
B	526	HIS	-	expression tag	UNP A0A1D8PMC5
B	527	HIS	-	expression tag	UNP A0A1D8PMC5
B	528	HIS	-	expression tag	UNP A0A1D8PMC5
B	529	HIS	-	expression tag	UNP A0A1D8PMC5

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			8	6	2		



- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	K	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	254	Total	O	0	0
			254	254		
5	B	266	Total	O	0	0
			266	266		

- Molecule 1: Aromatic-amino-acid:2-oxoglutarate transaminase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.38 Å 89.28 Å 161.91 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.93 – 2.23 46.93 – 2.23	Depositor EDS
% Data completeness (in resolution range)	99.7 (46.93-2.23) 99.7 (46.93-2.23)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.31 (at 2.22 Å)	Xtriage
Refinement program	REFMAC 5.8.0232	Depositor
R, R_{free}	0.169 , 0.230 0.175 , 0.234	Depositor DCC
R_{free} test set	2696 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	29.7	Xtriage
Anisotropy	0.530	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 30.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8124	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7014e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, MPD, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/3863	0.83	0/5243
1	B	0.70	0/3888	0.83	0/5277
All	All	0.70	0/7751	0.83	0/10520

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3767	0	3713	13	0
1	B	3790	0	3734	17	0
2	A	15	0	7	0	0
2	B	15	0	7	0	0
3	A	8	0	14	0	0
3	B	8	0	14	0	0
4	B	1	0	0	0	0
5	A	254	0	0	1	0
5	B	266	0	0	2	0
All	All	8124	0	7489	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:421:GLN:HG3	5:B:941:HOH:O	1.72	0.89
1:B:389:PHE:O	1:B:394:GLY:HA3	1.91	0.70
1:A:430:VAL:HG12	1:A:439:VAL:CG1	2.24	0.66
1:B:430:VAL:HG12	1:B:439:VAL:CG1	2.37	0.55
1:A:389:PHE:O	1:A:394:GLY:HA3	2.07	0.54
1:A:68:ILE:C	5:A:888:HOH:O	2.46	0.53
1:A:343:PHE:CD1	1:A:347:LEU:HD11	2.44	0.51
1:A:160:VAL:HG12	1:A:353:VAL:HG22	1.91	0.51
1:B:120:LYS:HB3	1:B:386:GLN:HG2	1.92	0.51
1:B:347:LEU:HD13	1:B:381:VAL:HG21	1.96	0.47
1:B:221:LEU:HD21	1:B:265:LYS:HD2	1.97	0.46
1:B:11:ARG:NH2	1:B:178:LEU:O	2.46	0.45
1:A:495:LEU:HB2	5:B:919:HOH:O	2.17	0.43
1:A:128:LEU:O	1:B:344:ALA:HB1	2.18	0.43
1:A:254:THR:HA	1:A:440:THR:OG1	2.19	0.43
1:A:483:LYS:HG2	1:A:484:ASP:N	2.34	0.43
1:B:137:ALA:HB3	1:B:138:PRO:HD3	1.99	0.43
1:B:342:LEU:HD22	1:B:399:ILE:HG12	2.00	0.42
1:B:254:THR:HA	1:B:440:THR:OG1	2.19	0.42
1:B:256:PHE:CE1	1:B:431:ILE:HG21	2.54	0.42
1:B:3:ASP:OD1	1:B:3:ASP:C	2.58	0.42
1:B:142:PHE:CE1	1:B:385:ILE:HG21	2.55	0.41
1:A:171:ASN:HA	1:A:197:PHE:HB3	2.01	0.41
1:A:334:LEU:HD23	1:A:334:LEU:C	2.41	0.41
1:B:524:HIS:C	1:B:524:HIS:CD2	2.94	0.41
1:B:428:VAL:CG1	1:B:513:VAL:HG11	2.50	0.41
1:B:363:LYS:NZ	1:B:367:ASP:OD2	2.48	0.41
1:A:310:LEU:HD23	1:A:310:LEU:HA	1.85	0.40
1:A:183:VAL:HG21	1:A:238:LEU:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	459/529 (87%)	438 (95%)	21 (5%)	0	100	100
1	B	462/529 (87%)	442 (96%)	18 (4%)	2 (0%)	34	35
All	All	921/1058 (87%)	880 (96%)	39 (4%)	2 (0%)	47	53

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	521	LEU
1	B	219	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/467 (88%)	397 (96%)	16 (4%)	32	35
1	B	416/467 (89%)	399 (96%)	17 (4%)	30	33
All	All	829/934 (89%)	796 (96%)	33 (4%)	31	34

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	18	VAL
1	A	31	ASN
1	A	37	LYS
1	A	45	MET
1	A	68	ILE
1	A	180	ASP
1	A	283	TYR
1	A	293	LYS
1	A	295	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	313	ASP
1	A	317	LYS
1	A	405	ASN
1	A	430	VAL
1	A	502	ILE
1	A	515	GLU
1	B	26	ASP
1	B	43	TYR
1	B	105	THR
1	B	180	ASP
1	B	204	LYS
1	B	283	TYR
1	B	310	LEU
1	B	356	LYS
1	B	363	LYS
1	B	386	GLN
1	B	390	LYS
1	B	402	MET
1	B	410	LYS
1	B	430	VAL
1	B	483	LYS
1	B	495	LEU
1	B	524	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	295	ASN
1	A	474	ASN
1	B	388	ASN
1	B	524	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MPD	B	602	-	7,7,7	0.18	0	9,10,10	0.52	0
3	MPD	A	602	-	7,7,7	0.16	0	9,10,10	0.42	0
2	PLP	B	601	1	15,15,16	0.86	0	20,22,23	1.14	2 (10%)
2	PLP	A	601	1	15,15,16	0.89	1 (6%)	20,22,23	1.14	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	B	602	-	-	0/5/5/5	-
3	MPD	A	602	-	-	2/5/5/5	-
2	PLP	B	601	1	-	1/6/6/8	0/1/1/1
2	PLP	A	601	1	-	3/6/6/8	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	PLP	C5-C4	2.02	1.42	1.40

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	PLP	O4P-C5A-C5	3.19	115.43	109.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	PLP	O4P-C5A-C5	2.97	115.01	109.35
2	A	601	PLP	C4A-C4-C5	2.67	123.68	120.94
2	B	601	PLP	O2P-P-O4P	-2.14	101.04	106.73

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	PLP	C5A-O4P-P-O2P
2	A	601	PLP	C5A-O4P-P-O3P
3	A	602	MPD	CM-C2-C3-C4
2	A	601	PLP	C5A-O4P-P-O1P
3	A	602	MPD	O2-C2-C3-C4
2	B	601	PLP	C5A-O4P-P-O2P

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	467/529 (88%)	-0.42	11 (2%) 59 60	30, 40, 66, 105	0
1	B	470/529 (88%)	-0.44	11 (2%) 60 61	30, 41, 64, 112	0
All	All	937/1058 (88%)	-0.43	22 (2%) 60 61	30, 40, 66, 112	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	295	ASN	3.6
1	A	30	ALA	3.6
1	B	309	ASP	3.5
1	A	237	ASN	3.5
1	A	68	ILE	3.0
1	B	18	VAL	2.9
1	A	43	TYR	2.9
1	B	31	ASN	2.8
1	A	36	GLU	2.8
1	A	31	ASN	2.7
1	A	236	PRO	2.5
1	A	105	THR	2.5
1	A	32	PHE	2.5
1	B	25	SER	2.4
1	B	104	HIS	2.4
1	A	106	PRO	2.4
1	B	30	ALA	2.3
1	B	105	THR	2.3
1	B	68	ILE	2.3
1	A	232	GLU	2.2
1	B	106	PRO	2.2
1	B	24	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MPD	B	602	8/8	0.83	0.22	77,79,81,83	0
3	MPD	A	602	8/8	0.87	0.15	82,83,87,87	0
2	PLP	B	601	15/16	0.98	0.12	34,42,50,56	0
2	PLP	A	601	15/16	0.98	0.12	34,41,46,52	0
4	K	B	603	1/1	0.99	0.17	53,53,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.