



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 01:18 am BST

PDB ID : 1HR0  
Title : CRYSTAL STRUCTURE OF INITIATION FACTOR IF1 BOUND TO THE 30S RIBOSOMAL SUBUNIT  
Authors : Carter, A.P.; Clemons Jr., W.M.; Brodersen, D.E.; Morgan-Warren, R.J.; Wimberly, B.T.; Ramakrishnan, V.  
Deposited on : 2000-12-20  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)
RNA backbone	3102	1010 (3.50-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1522	
2	X	6	
3	B	256	
4	C	239	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	D	209	
6	E	162	
7	F	101	
8	G	156	
9	H	138	
10	I	128	
11	J	105	
12	K	129	
13	L	135	
14	M	126	
15	N	61	
16	O	89	
17	P	88	
18	Q	105	
19	R	88	
20	S	93	
21	T	106	
22	V	26	
23	W	71	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
24	MG	A	1547	-	-	-	X
24	MG	A	1548	-	-	-	X
24	MG	A	1549	-	-	-	X
24	MG	A	1561	-	-	-	X
24	MG	A	1565	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
24	MG	A	1570	-	-	-	X
24	MG	A	1573	-	-	-	X
24	MG	A	1584	-	-	-	X
24	MG	A	1591	-	-	-	X
24	MG	A	1592	-	-	-	X
24	MG	A	1596	-	-	-	X

## 2 Entry composition [i](#)

There are 25 unique types of molecules in this entry. The entry contains 52365 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 16S RIBOSOMAL RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1507	Total	C	N	O	P	22	0	0
			32382	14418	6002	10459	1503			

- Molecule 2 is a RNA chain called FRAGMENT OF MESSENGER RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	6	Total	C	N	O	P	0	0	0
			117	54	14	44	5			

- Molecule 3 is a protein called 30S RIBOSOMAL PROTEIN S2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	234	Total	C	N	O	S	0	0	0
			1900	1213	341	341	5			

- Molecule 4 is a protein called 30S RIBOSOMAL PROTEIN S3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	C	206	Total	C	N	O	S	0	0	0
			1612	1016	314	281	1			

- Molecule 5 is a protein called 30S RIBOSOMAL PROTEIN S4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	D	208	Total	C	N	O	S	0	0	0
			1703	1066	339	291	7			

- Molecule 6 is a protein called 30S RIBOSOMAL PROTEIN S5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	E	150	Total	C	N	O	S	0	0	0
			1146	724	217	201	4			

- Molecule 7 is a protein called 30S RIBOSOMAL PROTEIN S6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	F	101	Total	C	N	O	S	0	0	0
			843	531	155	154	3			

- Molecule 8 is a protein called 30S RIBOSOMAL PROTEIN S7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	G	155	Total	C	N	O	S	0	0	0
			1257	781	252	218	6			

- Molecule 9 is a protein called 30S RIBOSOMAL PROTEIN S8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	H	138	Total	C	N	O	S	0	0	0
			1116	705	215	193	3			

- Molecule 10 is a protein called 30S RIBOSOMAL PROTEIN S9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	I	127	Total	C	N	O	S	0	0	0
			1011	639	198	174				

- Molecule 11 is a protein called 30S RIBOSOMAL PROTEIN S10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	J	98	Total	C	N	O	S	0	0	0
			792	498	156	137	1			

- Molecule 12 is a protein called 30S RIBOSOMAL PROTEIN S11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	K	119	Total	C	N	O	S	0	0	0
			885	549	168	165	3			

- Molecule 13 is a protein called 30S RIBOSOMAL PROTEIN S12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	L	124	Total	C	N	O	S	0	0	0
			970	611	195	163	1			

- Molecule 14 is a protein called 30S RIBOSOMAL PROTEIN S13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	M	125	Total	C	N	O	S	0	0	0
			997	617	207	171	2			

- Molecule 15 is a protein called 30S RIBOSOMAL PROTEIN S14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	N	60	Total	C	N	O	S	0	0	0
			492	312	104	72	4			

- Molecule 16 is a protein called 30S RIBOSOMAL PROTEIN S15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
16	O	88	Total	C	N	O	S	0	0	0
			734	459	147	126	2			

- Molecule 17 is a protein called 30S RIBOSOMAL PROTEIN S16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
17	P	83	Total	C	N	O	S	0	0	0
			700	443	139	117	1			

- Molecule 18 is a protein called 30S RIBOSOMAL PROTEIN S17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
18	Q	104	Total	C	N	O	S	0	0	0
			857	547	161	147	2			

- Molecule 19 is a protein called 30S RIBOSOMAL PROTEIN S18.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
19	R	73	Total	C	N	O	0	0	0
			597	380	118	99			

- Molecule 20 is a protein called 30S RIBOSOMAL PROTEIN S19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
20	S	80	Total	C	N	O	S	0	0	0
			647	414	119	112	2			

- Molecule 21 is a protein called 30S RIBOSOMAL PROTEIN S20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
21	T	99	Total	C	N	O	S	0	0	0
			762	469	162	129	2			

- Molecule 22 is a protein called 30S RIBOSOMAL PROTEIN THX.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
22	V	24	Total	C	N	O	0	0	0
			208	128	50	30			

- Molecule 23 is a protein called TRANSLATION INITIATION FACTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
23	W	71	Total	C	N	O	S	0	0	0
			570	362	103	103	2			

- Molecule 24 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
24	W	1	Total	Mg	0	0
			1	1		
24	A	64	Total	Mg	0	0
			64	64		

- Molecule 25 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
25	D	1	Total	Zn	0	0
			1	1		
25	N	1	Total	Zn	0	0
			1	1		














• Molecule 6: 30S RIBOSOMAL PROTEIN S5




• Molecule 7: 30S RIBOSOMAL PROTEIN S6




• Molecule 8: 30S RIBOSOMAL PROTEIN S7




• Molecule 9: 30S RIBOSOMAL PROTEIN S8




- Molecule 10: 30S RIBOSOMAL PROTEIN S9




- Molecule 11: 30S RIBOSOMAL PROTEIN S10



- Molecule 12: 30S RIBOSOMAL PROTEIN S11




- Molecule 13: 30S RIBOSOMAL PROTEIN S12



- Molecule 14: 30S RIBOSOMAL PROTEIN S13






• Molecule 15: 30S RIBOSOMAL PROTEIN S14

Chain N: 26% 59% 13%




• Molecule 16: 30S RIBOSOMAL PROTEIN S15

Chain O: 2% 33% 61% 6%




• Molecule 17: 30S RIBOSOMAL PROTEIN S16

Chain P: 2% 36% 51% 7% 6%



• Molecule 18: 30S RIBOSOMAL PROTEIN S17



Chain Q: 5% 34% 55% 10%




• Molecule 19: 30S RIBOSOMAL PROTEIN S18

Chain R: 3% 18% 53% 10% 17%

• Molecule 20: 30S RIBOSOMAL PROTEIN S19




● Molecule 21: 30S RIBOSOMAL PROTEIN S20



● Molecule 22: 30S RIBOSOMAL PROTEIN THX



● Molecule 23: TRANSLATION INITIATION FACTOR



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	399.58Å 399.58Å 176.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.90 – 3.20 29.75 – 3.08	Depositor EDS
% Data completeness (in resolution range)	91.5 (28.90-3.20) 89.6 (29.75-3.08)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 3.06Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.218 , 0.261 0.202 , 0.243	Depositor DCC
$R_{free}$ test set	11800 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	77.4	Xtriage
Anisotropy	0.307	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 86.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	52365	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.51	0/36247	0.75	45/56569 (0.1%)
2	X	0.56	0/128	0.73	0/196
3	B	0.37	0/1935	0.70	0/2609
4	C	0.38	0/1636	0.63	0/2205
5	D	0.40	0/1733	0.65	0/2318
6	E	0.47	0/1162	0.75	0/1564
7	F	0.34	0/856	0.61	0/1154
8	G	0.36	0/1276	0.61	0/1709
9	H	0.45	0/1136	0.77	1/1527 (0.1%)
10	I	0.36	0/1029	0.67	0/1378
11	J	0.38	0/805	0.73	1/1082 (0.1%)
12	K	0.38	0/900	0.70	0/1213
13	L	0.42	0/986	0.74	0/1320
14	M	0.36	0/1008	0.69	0/1347
15	N	0.45	0/501	0.85	0/664
16	O	0.38	0/745	0.62	0/992
17	P	0.44	0/716	0.77	0/963
18	Q	0.45	0/870	0.78	0/1159
19	R	0.39	0/603	0.66	0/799
20	S	0.35	0/661	0.69	0/890
21	T	0.39	0/764	0.72	0/1006
22	V	0.45	0/212	0.67	0/277
23	W	0.39	0/580	0.70	0/782
All	All	0.47	0/56489	0.74	47/83723 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	3	39

There are no bond length outliers.

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	559	A	C2'-C3'-O3'	9.71	130.87	109.50
1	A	366	C	C2'-C3'-O3'	8.88	129.03	109.50
1	A	1498	U	C2'-C3'-O3'	8.70	128.64	109.50
1	A	1528	U	C2'-C3'-O3'	8.48	128.15	109.50
1	A	1302	U	C2'-C3'-O3'	8.21	127.56	109.50

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	559	A	C3'
1	A	1498	U	C3'
1	A	1528	U	C3'

5 of 39 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	115	G	Sidechain
1	A	197	A	Sidechain
1	A	203	U	Sidechain
1	A	37	U	Sidechain
1	A	77	G	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	32382	0	16356	1335	0
2	X	117	0	64	1	0
3	B	1900	0	1951	296	0
4	C	1612	0	1677	271	0
5	D	1703	0	1765	158	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	E	1146	0	1207	116	0
7	F	843	0	857	84	0
8	G	1257	0	1296	102	0
9	H	1116	0	1177	98	0
10	I	1011	0	1043	155	0
11	J	792	0	835	163	0
12	K	885	0	904	84	0
13	L	970	0	1057	134	0
14	M	997	0	1072	136	0
15	N	492	0	529	73	0
16	O	734	0	771	76	0
17	P	700	0	720	61	0
18	Q	857	0	930	103	0
19	R	597	0	668	94	0
20	S	647	0	673	99	0
21	T	762	0	859	85	0
22	V	208	0	221	25	0
23	W	570	0	599	82	0
24	A	64	0	0	0	0
24	W	1	0	0	0	0
25	D	1	0	0	0	0
25	N	1	0	0	0	0
All	All	52365	0	37231	3507	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 3507 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:I:8:GLY:HA2	10:I:79:LEU:HD12	1.19	1.18
3:B:77:ALA:HB2	3:B:211:ILE:HD13	1.22	1.17
5:D:36:ARG:H	5:D:37:PRO:HD3	1.04	1.17
13:L:33:ARG:HD3	13:L:62:SER:HB3	1.20	1.11
1:A:1256:A:H4'	1:A:1257:U:H5'	1.35	1.09

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	B	232/256 (91%)	156 (67%)	48 (21%)	28 (12%)	0	2
4	C	204/239 (85%)	134 (66%)	44 (22%)	26 (13%)	0	1
5	D	206/209 (99%)	159 (77%)	37 (18%)	10 (5%)	2	17
6	E	148/162 (91%)	130 (88%)	13 (9%)	5 (3%)	3	24
7	F	99/101 (98%)	79 (80%)	15 (15%)	5 (5%)	2	15
8	G	153/156 (98%)	111 (72%)	29 (19%)	13 (8%)	1	4
9	H	136/138 (99%)	122 (90%)	11 (8%)	3 (2%)	6	35
10	I	125/128 (98%)	84 (67%)	33 (26%)	8 (6%)	1	10
11	J	96/105 (91%)	62 (65%)	19 (20%)	15 (16%)	0	1
12	K	117/129 (91%)	87 (74%)	19 (16%)	11 (9%)	0	3
13	L	122/135 (90%)	91 (75%)	19 (16%)	12 (10%)	0	3
14	M	123/126 (98%)	84 (68%)	28 (23%)	11 (9%)	1	4
15	N	58/61 (95%)	38 (66%)	15 (26%)	5 (9%)	1	4
16	O	86/89 (97%)	66 (77%)	18 (21%)	2 (2%)	6	34
17	P	81/88 (92%)	58 (72%)	15 (18%)	8 (10%)	0	3
18	Q	102/105 (97%)	85 (83%)	9 (9%)	8 (8%)	1	6
19	R	71/88 (81%)	50 (70%)	16 (22%)	5 (7%)	1	8
20	S	78/93 (84%)	50 (64%)	19 (24%)	9 (12%)	0	2
21	T	97/106 (92%)	66 (68%)	21 (22%)	10 (10%)	0	3
22	V	22/26 (85%)	18 (82%)	2 (9%)	2 (9%)	1	3
23	W	69/71 (97%)	47 (68%)	15 (22%)	7 (10%)	0	3
All	All	2425/2611 (93%)	1777 (73%)	445 (18%)	203 (8%)	1	5

5 of 203 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	8	LYS
3	B	15	VAL
3	B	16	HIS
3	B	17	PHE
3	B	21	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	B	202/220 (92%)	174 (86%)	28 (14%)	3	16
4	C	160/188 (85%)	138 (86%)	22 (14%)	3	16
5	D	180/181 (99%)	170 (94%)	10 (6%)	21	57
6	E	115/123 (94%)	97 (84%)	18 (16%)	2	12
7	F	90/90 (100%)	83 (92%)	7 (8%)	12	43
8	G	126/127 (99%)	118 (94%)	8 (6%)	18	52
9	H	119/119 (100%)	107 (90%)	12 (10%)	7	29
10	I	98/99 (99%)	87 (89%)	11 (11%)	6	25
11	J	87/92 (95%)	77 (88%)	10 (12%)	5	24
12	K	90/99 (91%)	80 (89%)	10 (11%)	6	25
13	L	104/111 (94%)	96 (92%)	8 (8%)	13	44
14	M	100/101 (99%)	88 (88%)	12 (12%)	5	22
15	N	49/50 (98%)	42 (86%)	7 (14%)	3	15
16	O	79/80 (99%)	73 (92%)	6 (8%)	13	45
17	P	72/74 (97%)	70 (97%)	2 (3%)	43	74
18	Q	96/97 (99%)	91 (95%)	5 (5%)	23	59
19	R	64/77 (83%)	57 (89%)	7 (11%)	6	26
20	S	71/80 (89%)	67 (94%)	4 (6%)	21	57
21	T	76/82 (93%)	67 (88%)	9 (12%)	5	23
22	V	19/21 (90%)	18 (95%)	1 (5%)	22	58

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
23	W	62/62 (100%)	54 (87%)	8 (13%)	4	19
All	All	2059/2173 (95%)	1854 (90%)	205 (10%)	7	30

5 of 205 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
8	G	136	LYS
10	I	111	ARG
21	T	42	GLN
9	H	52	ASP
9	H	133	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 59 such sidechains are listed below:

Mol	Chain	Res	Type
7	F	32	ASN
10	I	23	ASN
20	S	53	ASN
7	F	94	GLN
8	G	37	ASN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	1505/1522 (98%)	224 (14%)	85 (5%)
2	X	5/6 (83%)	0	0
All	All	1510/1528 (98%)	224 (14%)	85 (5%)

5 of 224 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	6	G
1	A	8	A
1	A	9	G
1	A	31	G
1	A	32	A

5 of 85 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	575	G
1	A	965	A
1	A	1451	A
1	A	687	A
1	A	812	C

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 67 ligands modelled in this entry, 67 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1506/1522 (98%)	0.54	85 (5%) 24 13	38, 69, 147, 199	0
2	X	6/6 (100%)	1.27	2 (33%) 0 0	52, 71, 99, 119	0
3	B	234/256 (91%)	-0.00	9 (3%) 40 26	37, 90, 147, 184	0
4	C	206/239 (86%)	-0.09	1 (0%) 91 86	48, 89, 138, 162	0
5	D	208/209 (99%)	-0.21	0 100 100	41, 69, 117, 157	0
6	E	150/162 (92%)	-0.28	1 (0%) 87 81	37, 60, 95, 145	0
7	F	101/101 (100%)	-0.01	0 100 100	55, 94, 140, 153	0
8	G	155/156 (99%)	-0.06	5 (3%) 47 31	49, 81, 137, 170	0
9	H	138/138 (100%)	-0.28	0 100 100	33, 56, 87, 126	0
10	I	127/128 (99%)	-0.06	0 100 100	43, 93, 132, 163	0
11	J	98/105 (93%)	0.23	7 (7%) 16 9	54, 116, 167, 186	0
12	K	119/129 (92%)	-0.16	1 (0%) 86 78	36, 70, 116, 168	0
13	L	124/135 (91%)	-0.12	1 (0%) 86 78	28, 72, 120, 181	0
14	M	125/126 (99%)	0.13	7 (5%) 24 13	52, 85, 141, 188	0
15	N	60/61 (98%)	-0.07	0 100 100	44, 83, 132, 153	0
16	O	88/89 (98%)	-0.14	2 (2%) 60 47	44, 68, 123, 180	0
17	P	83/88 (94%)	-0.16	2 (2%) 59 44	43, 60, 83, 137	0
18	Q	104/105 (99%)	0.27	5 (4%) 30 18	31, 60, 132, 199	0
19	R	73/88 (82%)	0.06	3 (4%) 37 24	50, 75, 133, 181	0
20	S	80/93 (86%)	0.35	4 (5%) 28 16	74, 100, 149, 196	0
21	T	99/106 (93%)	-0.18	0 100 100	42, 70, 110, 169	0
22	V	24/26 (92%)	-0.22	0 100 100	53, 75, 122, 137	0
23	W	71/71 (100%)	0.40	3 (4%) 36 23	80, 112, 138, 172	0
All	All	3979/4139 (96%)	0.18	138 (3%) 44 28	28, 75, 140, 199	0



The worst 5 of 138 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
18	Q	105	ALA	17.8
18	Q	103	GLY	16.4
18	Q	102	GLY	13.7
1	A	1129	C	11.1
14	M	124	PRO	9.2

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
24	MG	A	1548	1/1	0.40	0.59	65,65,65,65	0
24	MG	A	1565	1/1	0.55	0.46	65,65,65,65	0
24	MG	A	1584	1/1	0.60	0.41	65,65,65,65	1
24	MG	A	1591	1/1	0.63	0.44	65,65,65,65	0
24	MG	A	1563	1/1	0.65	0.17	65,65,65,65	0
24	MG	A	1547	1/1	0.66	0.95	65,65,65,65	0
24	MG	A	1559	1/1	0.68	0.30	65,65,65,65	0
24	MG	A	1579	1/1	0.70	0.29	65,65,65,65	0
24	MG	A	1596	1/1	0.70	0.45	65,65,65,65	0
24	MG	A	1570	1/1	0.73	0.58	65,65,65,65	0
24	MG	A	1561	1/1	0.74	0.69	65,65,65,65	0
24	MG	A	1589	1/1	0.74	0.36	65,65,65,65	0
24	MG	A	1573	1/1	0.78	0.43	65,65,65,65	0
24	MG	A	1592	1/1	0.79	0.70	65,65,65,65	0
24	MG	A	1549	1/1	0.80	2.64	65,65,65,65	1
24	MG	A	1555	1/1	0.82	0.86	65,65,65,65	0
24	MG	A	1577	1/1	0.82	0.31	65,65,65,65	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
24	MG	A	1605	1/1	0.83	0.36	65,65,65,65	0
24	MG	A	1576	1/1	0.84	0.26	65,65,65,65	0
24	MG	A	1604	1/1	0.84	0.89	65,65,65,65	0
24	MG	A	1585	1/1	0.84	0.43	65,65,65,65	0
24	MG	A	1558	1/1	0.85	0.36	65,65,65,65	0
24	MG	A	1599	1/1	0.85	0.83	65,65,65,65	0
24	MG	A	1580	1/1	0.86	0.41	65,65,65,65	0
24	MG	A	1602	1/1	0.87	0.67	65,65,65,65	0
24	MG	A	1550	1/1	0.87	1.22	65,65,65,65	0
24	MG	A	1575	1/1	0.87	0.68	65,65,65,65	0
24	MG	A	1566	1/1	0.88	0.63	65,65,65,65	0
24	MG	A	1545	1/1	0.88	0.67	65,65,65,65	0
24	MG	A	87	1/1	0.88	0.44	65,65,65,65	0
24	MG	A	1557	1/1	0.88	0.83	65,65,65,65	0
24	MG	A	1586	1/1	0.89	0.32	65,65,65,65	0
24	MG	A	1588	1/1	0.89	0.50	65,65,65,65	0
24	MG	A	1587	1/1	0.89	0.39	65,65,65,65	0
24	MG	A	1582	1/1	0.89	0.25	65,65,65,65	0
24	MG	A	1590	1/1	0.89	0.79	65,65,65,65	0
24	MG	A	1569	1/1	0.90	0.25	65,65,65,65	1
24	MG	A	1556	1/1	0.90	0.74	65,65,65,65	0
24	MG	A	1567	1/1	0.91	0.66	65,65,65,65	0
24	MG	A	1594	1/1	0.91	0.57	65,65,65,65	0
24	MG	A	71	1/1	0.91	0.26	65,65,65,65	0
24	MG	A	1546	1/1	0.91	0.78	65,65,65,65	0
24	MG	A	1552	1/1	0.91	1.01	65,65,65,65	0
24	MG	A	1601	1/1	0.91	1.13	65,65,65,65	0
24	MG	A	1583	1/1	0.92	0.45	65,65,65,65	0
24	MG	W	72	1/1	0.92	0.47	65,65,65,65	1
24	MG	A	1554	1/1	0.93	1.07	65,65,65,65	0
24	MG	A	1571	1/1	0.93	0.82	65,65,65,65	0
24	MG	A	1598	1/1	0.93	0.22	65,65,65,65	0
24	MG	A	1600	1/1	0.93	0.34	65,65,65,65	0
24	MG	A	1551	1/1	0.93	0.29	65,65,65,65	0
24	MG	A	1560	1/1	0.93	0.35	65,65,65,65	0
24	MG	A	1574	1/1	0.93	0.29	65,65,65,65	0
24	MG	A	1597	1/1	0.93	0.24	65,65,65,65	0
24	MG	A	1553	1/1	0.94	1.25	65,65,65,65	0
24	MG	A	1562	1/1	0.94	0.16	65,65,65,65	0
24	MG	A	1603	1/1	0.94	0.10	65,65,65,65	0
24	MG	A	1564	1/1	0.95	0.86	65,65,65,65	0
24	MG	A	1595	1/1	0.95	0.50	65,65,65,65	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
24	MG	A	1578	1/1	0.95	0.24	65,65,65,65	0
25	ZN	D	300	1/1	0.96	0.42	65,65,65,65	0
24	MG	A	1568	1/1	0.97	0.21	65,65,65,65	1
24	MG	A	86	1/1	0.97	0.85	65,65,65,65	0
24	MG	A	1593	1/1	0.98	0.18	65,65,65,65	0
24	MG	A	1581	1/1	0.98	0.15	65,65,65,65	1
24	MG	A	1572	1/1	0.99	0.60	65,65,65,65	0
25	ZN	N	190	1/1	1.00	0.13	65,65,65,65	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.