



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 01:56 AM BST

PDB ID : 5HRO  
Title : STRUCTURE OF HIV-1 REVERSE TRANSCRIPTASE In COMPLEX  
WITH A DNA aptamer and an Alpha-carboxy nucleoside phosphonate in-  
hibitor (alpha-CNP)  
Authors : Das, K.; Arnold, E.  
Deposited on : 2016-01-23  
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	<div> <div>17%</div> <div> <div></div> <div>76%</div> <div>23%</div> <div>.</div> </div> </div>
1	C	555	<div> <div>24%</div> <div> <div></div> <div>75%</div> <div>24%</div> <div>..</div> </div> </div>
2	B	444	<div> <div>7%</div> <div> <div></div> <div>78%</div> <div>14%</div> <div>7%</div> </div> </div>
2	D	444	<div> <div>12%</div> <div> <div></div> <div>75%</div> <div>18%</div> <div>7%</div> </div> </div>
3	E	38	<div> <div></div> <div> <div></div> <div>58%</div> <div>32%</div> <div>.</div> <div>8%</div> </div> </div>
3	F	38	<div> <div>5%</div> <div> <div></div> <div>55%</div> <div>29%</div> <div>8%</div> <div>8%</div> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	G	2	<div><div></div><div>50%</div><div></div><div>50%</div></div>
4	H	2	<div><div></div><div>100%</div></div>

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 17349 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 REVERSE TRANSCRIPTASE P66 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	551	Total	C	N	O	S	0	0	0
			4488	2905	747	829	7			
1	C	551	Total	C	N	O	S	0	0	0
			4488	2905	747	829	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	280	SER	CYS	engineered mutation	UNP P03366
C	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 2 is a protein called HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	412	Total	C	N	O	S	0	0	0
			3400	2212	563	619	6			
2	D	412	Total	C	N	O	S	0	0	0
			3400	2212	563	619	6			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	MET	-	initiating methionine	UNP P03366
B	-14	ALA	-	expression tag	UNP P03366
B	-13	HIS	-	expression tag	UNP P03366
B	-12	HIS	-	expression tag	UNP P03366
B	-11	HIS	-	expression tag	UNP P03366
B	-10	HIS	-	expression tag	UNP P03366
B	-9	HIS	-	expression tag	UNP P03366
B	-8	HIS	-	expression tag	UNP P03366
B	-7	ALA	-	expression tag	UNP P03366
B	-6	LEU	-	expression tag	UNP P03366

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	GLU	-	expression tag	UNP P03366
B	-4	VAL	-	expression tag	UNP P03366
B	-3	LEU	-	expression tag	UNP P03366
B	-2	PHE	-	expression tag	UNP P03366
B	-1	GLN	-	expression tag	UNP P03366
B	0	GLY	-	expression tag	UNP P03366
B	280	SER	CYS	engineered mutation	UNP P03366
D	-15	MET	-	initiating methionine	UNP P03366
D	-14	ALA	-	expression tag	UNP P03366
D	-13	HIS	-	expression tag	UNP P03366
D	-12	HIS	-	expression tag	UNP P03366
D	-11	HIS	-	expression tag	UNP P03366
D	-10	HIS	-	expression tag	UNP P03366
D	-9	HIS	-	expression tag	UNP P03366
D	-8	HIS	-	expression tag	UNP P03366
D	-7	ALA	-	expression tag	UNP P03366
D	-6	LEU	-	expression tag	UNP P03366
D	-5	GLU	-	expression tag	UNP P03366
D	-4	VAL	-	expression tag	UNP P03366
D	-3	LEU	-	expression tag	UNP P03366
D	-2	PHE	-	expression tag	UNP P03366
D	-1	GLN	-	expression tag	UNP P03366
D	0	GLY	-	expression tag	UNP P03366
D	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is a DNA chain called DNA (38-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	35	Total	C	N	O	P	0	0	0
			720	340	130	215	35			
3	F	35	Total	C	N	O	P	0	0	0
			720	340	130	215	35			

- Molecule 4 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	G	2	Total	C	O	0	0	0
			23	12	11			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	H	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Mg	0	0
			3	3		
5	C	3	Total	Mg	0	0
			3	3		

- Molecule 6 is [(1R)-2-methoxy-1-{[(1S,3R)-3-(5-methyl-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)cyclopentyl]oxy}-2-oxoethyl]phosphonic acid (three-letter code: 3JY) (formula: C<sub>13</sub>H<sub>19</sub>N<sub>2</sub>O<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			23	12	2	8	1		
6	C	1	Total	C	N	O	P	0	0
			23	12	2	8	1		

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 8 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	O	0	0
			1	1		
8	B	1	Total	O	0	0
			1	1		
8	C	3	Total	O	0	0
			3	3		










• Molecule 2: HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT



• Molecule 2: HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT



• Molecule 3: DNA (38-MER)




• Molecule 3: DNA (38-MER)



- Molecule 4: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



- Molecule 4: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.25Å 128.96Å 131.05Å 90.00° 101.47° 90.00°	Depositor
Resolution (Å)	38.52 – 2.75 38.52 – 2.73	Depositor EDS
% Data completeness (in resolution range)	95.8 (38.52-2.75) 94.7 (38.52-2.73)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 2.72Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, $R_{free}$	0.235 , 0.265 0.235 , 0.264	Depositor DCC
$R_{free}$ test set	2960 reflections (3.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.1	Xtriage
Anisotropy	0.348	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 62.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	17349	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	96.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OMC, MG, 3JY, GLC, SO4, FRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/4605	0.49	0/6255
1	C	0.28	0/4605	0.49	1/6255 (0.0%)
2	B	0.27	0/3497	0.47	0/4751
2	D	0.29	0/3497	0.49	0/4751
3	E	0.64	1/759 (0.1%)	1.01	0/1170
3	F	0.72	3/759 (0.4%)	1.01	2/1170 (0.2%)
All	All	0.33	4/17722 (0.0%)	0.56	3/24352 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	26	DC	C4'-O4'	7.28	1.52	1.45
3	F	19	DC	C4'-O4'	5.75	1.50	1.45
3	E	24	DC	C4'-O4'	5.24	1.50	1.45
3	F	24	DC	C4'-O4'	5.10	1.50	1.45

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	7	DC	O4'-C4'-C3'	-6.52	101.89	104.50
3	F	19	DC	O4'-C4'-C3'	-6.06	102.08	104.50
1	C	55	PRO	N-CA-C	5.21	125.65	112.10

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4488	0	4545	86	0
1	C	4488	0	4545	95	0
2	B	3400	0	3431	41	0
2	D	3400	0	3430	54	0
3	E	720	0	396	13	0
3	F	720	0	396	12	0
4	G	23	0	21	0	0
4	H	23	0	21	0	0
5	A	3	0	0	0	0
5	C	3	0	0	0	0
6	A	23	0	14	0	0
6	C	23	0	14	4	0
7	A	5	0	0	0	0
7	B	20	0	0	0	0
7	C	5	0	0	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	3	0	0	0	0
All	All	17349	0	16813	277	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 277 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:ASN:HB2	1:A:289:LEU:HD21	1.27	1.09
1:A:255:ASN:HB2	1:A:289:LEU:CD2	1.98	0.92
1:C:221:HIS:HE1	1:C:223:LYS:HG3	1.39	0.88
1:A:255:ASN:CB	1:A:289:LEU:HD21	2.11	0.80
1:C:255:ASN:HB2	1:C:289:LEU:CD2	2.10	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	549/555 (99%)	536 (98%)	13 (2%)	0	100	100
1	C	549/555 (99%)	533 (97%)	16 (3%)	0	100	100
2	B	408/444 (92%)	396 (97%)	12 (3%)	0	100	100
2	D	408/444 (92%)	396 (97%)	12 (3%)	0	100	100
All	All	1914/1998 (96%)	1861 (97%)	53 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	492/495 (99%)	486 (99%)	6 (1%)	71	82
1	C	492/495 (99%)	483 (98%)	9 (2%)	59	75
2	B	374/403 (93%)	373 (100%)	1 (0%)	92	95
2	D	374/403 (93%)	372 (100%)	2 (0%)	88	92
All	All	1732/1796 (96%)	1714 (99%)	18 (1%)	76	85

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	42	GLU
1	C	109	LEU
1	C	450	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	11	LYS
1	C	5	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	255	ASN
2	D	208	HIS
1	C	487	GLN
2	B	428	GLN
1	C	507	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	OMC	F	8	3,2	15,22,23	0.91	1 (6%)	17,31,34	1.36	2 (11%)
3	OMC	E	8	3,2	15,22,23	0.86	1 (6%)	17,31,34	1.39	2 (11%)
3	OMC	E	6	3	15,22,23	0.93	1 (6%)	17,31,34	1.39	2 (11%)
3	OMC	F	6	3	15,22,23	0.94	1 (6%)	17,31,34	1.47	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OMC	F	8	3,2	-	1/7/27/28	0/2/2/2
3	OMC	E	8	3,2	-	0/7/27/28	0/2/2/2
3	OMC	E	6	3	-	1/7/27/28	0/2/2/2
3	OMC	F	6	3	-	0/7/27/28	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	8	OMC	C4-N4	2.73	1.43	1.35
3	E	6	OMC	C4-N4	2.71	1.43	1.35
3	F	6	OMC	C4-N4	2.70	1.43	1.35
3	F	8	OMC	C4-N4	2.68	1.43	1.35

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	8	OMC	C2-N3-C4	4.47	120.87	116.34
3	F	6	OMC	C2-N3-C4	4.47	120.87	116.34
3	E	6	OMC	C2-N3-C4	4.43	120.83	116.34
3	F	8	OMC	C2-N3-C4	4.18	120.58	116.34
3	F	6	OMC	N4-C4-N3	2.36	120.21	116.49

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	8	OMC	C1'-C2'-O2'-CM2
3	E	6	OMC	C1'-C2'-O2'-CM2

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	8	OMC	2	0
3	E	8	OMC	2	0
3	F	6	OMC	1	0

## 5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul



statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	GLC	G	1	4	11,11,12	0.61	0	15,15,17	0.83	1 (6%)
4	FRU	G	2	4	11,12,12	0.63	0	10,18,18	0.92	0
4	GLC	H	1	4	11,11,12	0.55	0	15,15,17	1.13	2 (13%)
4	FRU	H	2	4	11,12,12	0.67	0	10,18,18	1.16	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	G	1	4	-	0/2/19/22	0/1/1/1
4	FRU	G	2	4	-	3/5/24/24	0/1/1/1
4	GLC	H	1	4	-	0/2/19/22	0/1/1/1
4	FRU	H	2	4	-	3/5/24/24	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1	GLC	C1-C2-C3	2.51	112.75	109.67
4	G	1	GLC	C1-O5-C5	2.25	115.24	112.19
4	H	1	GLC	C1-O5-C5	2.22	115.20	112.19
4	H	2	FRU	C6-C5-C4	-2.17	109.85	115.09

There are no chirality outliers.


5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	G	2	FRU	O1-C1-C2-C3
4	G	2	FRU	O1-C1-C2-O2
4	G	2	FRU	O1-C1-C2-O5
4	H	2	FRU	O1-C1-C2-O2
4	H	2	FRU	O1-C1-C2-O5

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 6 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	3JY	C	605	5	15,24,25	2.56	5 (33%)	16,36,37	2.47	5 (31%)
7	SO4	B	503	-	4,4,4	0.12	0	6,6,6	0.12	0
7	SO4	B	504	-	4,4,4	0.13	0	6,6,6	0.21	0
7	SO4	A	605	-	4,4,4	0.13	0	6,6,6	0.11	0
6	3JY	A	603	5	15,24,25	2.45	4 (26%)	16,36,37	2.82	6 (37%)
7	SO4	C	604	-	4,4,4	0.13	0	6,6,6	0.17	0
7	SO4	B	502	-	4,4,4	0.18	0	6,6,6	0.36	0
7	SO4	B	505	-	4,4,4	0.14	0	6,6,6	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	3JY	C	605	5	-	5/6/27/29	0/2/2/2
6	3JY	A	603	5	-	6/6/27/29	0/2/2/2

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	605	3JY	O11-C10	-7.50	1.34	1.44
6	A	603	3JY	O11-C10	-7.47	1.34	1.44
6	C	605	3JY	O1-C2	3.92	1.34	1.24
6	A	603	3JY	O1-C2	3.90	1.34	1.24
6	C	605	3JY	P16-O18	2.51	1.53	1.49

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	605	3JY	C2-N21-C22	7.63	121.58	115.14
6	A	603	3JY	C2-N21-C22	7.52	121.49	115.14
6	A	603	3JY	C12-O11-C10	5.61	135.79	115.39
6	C	605	3JY	C12-O11-C10	2.82	125.66	115.39
6	C	605	3JY	C10-C20-C7	2.62	108.66	102.43

There are no chirality outliers.

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	C	605	3JY	C20-C10-O11-C12
6	C	605	3JY	C13-C12-P16-O18
6	C	605	3JY	C13-C12-P16-O19
6	C	605	3JY	C13-C12-P16-O17
6	C	605	3JY	O11-C12-P16-O18


There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	605	3JY	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	551/555 (99%)	1.08	93 (16%) <b>1</b> <b>1</b>	41, 96, 151, 190	0
1	C	551/555 (99%)	1.34	135 (24%) <b>0</b> <b>0</b>	43, 113, 162, 182	0
2	B	412/444 (92%)	0.61	33 (8%) <b>12</b> <b>15</b>	41, 81, 127, 142	0
2	D	412/444 (92%)	0.85	55 (13%) <b>3</b> <b>4</b>	44, 92, 136, 160	0
3	E	33/38 (86%)	0.12	0 <b>100</b> <b>100</b>	70, 97, 113, 152	0
3	F	33/38 (86%)	0.79	2 (6%) <b>21</b> <b>26</b>	82, 112, 131, 169	0
All	All	1992/2074 (96%)	0.98	318 (15%) <b>1</b> <b>2</b>	41, 93, 152, 190	0

The worst 5 of 318 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	134	SER	14.5
1	C	60	VAL	12.0
1	C	132	ILE	11.7
1	A	131	THR	11.3
1	C	135	ILE	10.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	OMC	F	6	21/22	0.90	0.24	85,98,111,120	0
3	OMC	F	8	21/22	0.92	0.25	95,103,106,118	0
3	OMC	E	8	21/22	0.94	0.23	67,73,76,90	0
3	OMC	E	6	21/22	0.97	0.20	82,91,95,98	0

## 6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	FRU	G	2	12/12	0.86	0.21	82,97,108,109	0
4	GLC	H	1	11/12	0.93	0.13	82,92,96,101	0
4	FRU	H	2	12/12	0.93	0.17	93,101,102,104	0
4	GLC	G	1	11/12	0.95	0.15	67,77,86,87	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	MG	C	603	1/1	0.56	0.14	172,172,172,172	0
7	SO4	B	505	5/5	0.81	0.40	78,78,84,111	0
6	3JY	C	605	23/24	0.84	0.24	98,119,137,145	0
5	MG	A	602	1/1	0.85	0.15	114,114,114,114	0
7	SO4	B	504	5/5	0.85	0.40	87,98,102,121	0
7	SO4	B	502	5/5	0.85	0.21	78,79,85,101	0
7	SO4	B	503	5/5	0.85	0.17	94,100,109,113	0
7	SO4	C	604	5/5	0.88	0.24	84,92,112,112	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	MG	A	604	1/1	0.92	0.14	70,70,70,70	0
6	3JY	A	603	23/24	0.94	0.20	103,107,123,129	0
7	SO4	A	605	5/5	0.95	0.12	98,102,110,116	0
5	MG	C	602	1/1	0.95	0.19	70,70,70,70	0
5	MG	C	601	1/1	0.96	0.28	122,122,122,122	0
5	MG	A	601	1/1	0.97	0.20	114,114,114,114	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around 3JY C 605:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 3JY A 603:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.