



wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 02:50 pm BST

PDB ID : 3HW4
Title : Crystal structure of avian influenza A virus in complex with TMP
Authors : Zhao, C.; Lou, Z.; Guo, Y.; Ma, M.; Chen, Y.; Rao, Z.
Deposited on : 2009-06-17
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	<div> <div>47%</div> <div>16%</div> <div>• •</div> <div>32%</div> </div>
1	B	261	<div>4%</div> <div>48%</div> <div>15%</div> <div>5%</div> <div>•</div> <div>31%</div>
1	C	261	<div>3%</div> <div>48%</div> <div>15%</div> <div>5%</div> <div></div> <div>32%</div>
1	D	261	<div>2%</div> <div>45%</div> <div>19%</div> <div>•</div> <div>33%</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TMP	A	257	-	-	X	-
3	TMP	B	257	-	-	X	-
3	TMP	D	257	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6359 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polymerase acidic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	0	0
			1470	928	252	279	11			
1	B	180	Total	C	N	O	S	0	0	0
			1496	948	258	279	11			
1	C	177	Total	C	N	O	S	0	0	0
			1463	925	253	274	11			
1	D	176	Total	C	N	O	S	0	0	0
			1453	919	250	273	11			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	EXPRESSION TAG	UNP Q9Q0U9
A	-3	PRO	-	EXPRESSION TAG	UNP Q9Q0U9
A	-2	LEU	-	EXPRESSION TAG	UNP Q9Q0U9
A	-1	GLY	-	EXPRESSION TAG	UNP Q9Q0U9
A	0	SER	-	EXPRESSION TAG	UNP Q9Q0U9
A	201	ILE	VAL	SEE REMARK 999	UNP Q9Q0U9
B	-4	GLY	-	EXPRESSION TAG	UNP Q9Q0U9
B	-3	PRO	-	EXPRESSION TAG	UNP Q9Q0U9
B	-2	LEU	-	EXPRESSION TAG	UNP Q9Q0U9
B	-1	GLY	-	EXPRESSION TAG	UNP Q9Q0U9
B	0	SER	-	EXPRESSION TAG	UNP Q9Q0U9
B	201	ILE	VAL	SEE REMARK 999	UNP Q9Q0U9
C	-4	GLY	-	EXPRESSION TAG	UNP Q9Q0U9
C	-3	PRO	-	EXPRESSION TAG	UNP Q9Q0U9
C	-2	LEU	-	EXPRESSION TAG	UNP Q9Q0U9
C	-1	GLY	-	EXPRESSION TAG	UNP Q9Q0U9
C	0	SER	-	EXPRESSION TAG	UNP Q9Q0U9
C	201	ILE	VAL	SEE REMARK 999	UNP Q9Q0U9
D	-4	GLY	-	EXPRESSION TAG	UNP Q9Q0U9
D	-3	PRO	-	EXPRESSION TAG	UNP Q9Q0U9
D	-2	LEU	-	EXPRESSION TAG	UNP Q9Q0U9

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	GLY	-	EXPRESSION TAG	UNP Q9Q0U9
D	0	SER	-	EXPRESSION TAG	UNP Q9Q0U9
D	201	ILE	VAL	SEE REMARK 999	UNP Q9Q0U9

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0

- Molecule 3 is THYMIDINE-5'-PHOSPHATE (three-letter code: TMP) (formula: C₁₀H₁₅N₂O₈P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O P 21 10 2 8 1	0	0
3	B	1	Total C N O P 21 10 2 8 1	0	0
3	D	1	Total C N O P 21 10 2 8 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	127	Total 127	O 127	0	0
4	B	104	Total 104	O 104	0	0
4	C	89	Total 89	O 89	0	0
4	D	90	Total 90	O 90	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Polymerase acidic protein



• Molecule 1: Polymerase acidic protein



• Molecule 1: Polymerase acidic protein



GLU	GLU	THR	ILE	GLU	GLU	ARG	PHE	GLU	ILE	THR	GLY	THR	THR	MET	CYS	ARG	LEU	ALA	ASP	GLN	SER	LEU	PRO	PRO	ASN	PHE	SER	SER	LEU	GLY	LYS	PHE	ARG	ALA	TYR	VAL	ASP	GLY	PHE	GLU	PRO	ASN	GLY	CYS	ILE	GLU	GLY	LYS	LEU	SER	GLN	MET	SER	LYS	VAL	ASN	ALA	ARG
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

● Molecule 1: Polymerase acidic protein



ALA	ARG	E195	R196	G197	GLU	THR	ILE	GLU	GLU	ARG	PHE	GLU	ILE	THR	GLY	THR	MET	CYS	ARG	LEU	ALA	ASP	GLN	SER	LEU	GLU	LYS	PHE	ARG	ALA	TVR	VAL	ASP	GLY	PHE	GLU	PRO	ASN	GLY	CYS	ILE	GLU	GLY	LYS	LEU	SER	GLN	MET	SER	LYS	GLU	VAL	ASN																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																	
GLY	P-3	T85	R86	A87	M88	T89	V90	N91	N92	S93	N96	K102	P103	K104	R116	R124	R125	E126	V127	H128	E133	M136	K137	I138	K139	S140	E141	K142	T143	H144	I145	H146	S149	E153	E154	M155	R168	F176	T177	Q180	E181	L187	W188	D189	S190	F191	R192	Q193	S194																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																					

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	51.32Å 60.03Å 67.11Å 96.12° 96.36° 109.51°	Depositor
Resolution (Å)	47.78 – 1.90 47.78 – 1.90	Depositor EDS
% Data completeness (in resolution range)	93.5 (47.78-1.90) 93.5 (47.78-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.90 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.213 , 0.276 0.215 , 0.271	Depositor DCC
R_{free} test set	2794 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6359	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TMP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.13	4/1499 (0.3%)	1.46	17/2011 (0.8%)
1	B	1.13	4/1528 (0.3%)	1.26	16/2050 (0.8%)
1	C	0.99	1/1493 (0.1%)	1.06	9/2003 (0.4%)
1	D	1.04	1/1482 (0.1%)	1.14	8/1988 (0.4%)
All	All	1.08	10/6002 (0.2%)	1.24	50/8052 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
All	All	0	2

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	116	ARG	CZ-NH1	7.19	1.42	1.33
1	B	17	ALA	CA-CB	6.88	1.67	1.52
1	C	166	GLU	CG-CD	6.25	1.61	1.51
1	B	116	ARG	CZ-NH1	6.12	1.41	1.33
1	D	36	ALA	CA-CB	5.69	1.64	1.52

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	116	ARG	NE-CZ-NH1	23.66	132.13	120.30
1	A	116	ARG	NE-CZ-NH2	-21.84	109.38	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	168	ARG	NE-CZ-NH2	-17.01	111.80	120.30
1	B	116	ARG	NE-CZ-NH2	-15.88	112.36	120.30
1	D	168	ARG	NE-CZ-NH1	15.24	127.92	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	-3	PRO	Peptide
1	C	139	LYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1470	0	1439	59	0
1	B	1496	0	1458	44	0
1	C	1463	0	1436	49	0
1	D	1453	0	1429	62	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	21	0	13	10	0
3	B	21	0	11	20	0
3	D	21	0	13	15	0
4	A	127	0	0	6	0
4	B	104	0	0	9	0
4	C	89	0	0	8	0
4	D	90	0	0	5	0
All	All	6359	0	5799	221	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:257:TMP:C5'	3:B:257:TMP:O5'	1.63	1.42
1:A:84:ARG:NH2	3:B:257:TMP:H51	1.35	1.39
1:A:105:PHE:O	1:A:106:LEU:CB	1.77	1.15
3:A:257:TMP:C6	3:A:257:TMP:H5'1	1.84	1.12
1:C:155:MET:HG3	4:C:1029:HOH:O	1.49	1.10

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	174/261 (67%)	169 (97%)	4 (2%)	1 (1%)	25	15
1	B	176/261 (67%)	168 (96%)	5 (3%)	3 (2%)	9	2
1	C	173/261 (66%)	167 (96%)	3 (2%)	3 (2%)	9	2
1	D	172/261 (66%)	161 (94%)	9 (5%)	2 (1%)	13	4
All	All	695/1044 (67%)	665 (96%)	21 (3%)	9 (1%)	12	4

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	106	LEU
1	B	140	SER
1	B	141	GLU
1	C	94	ILE
1	D	140	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/232 (69%)	148 (92%)	12 (8%)	13	5
1	B	162/232 (70%)	148 (91%)	14 (9%)	10	4
1	C	159/232 (68%)	145 (91%)	14 (9%)	10	4
1	D	158/232 (68%)	147 (93%)	11 (7%)	15	7
All	All	639/928 (69%)	588 (92%)	51 (8%)	12	5

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	187	LEU
1	C	42	LEU
1	D	139	LYS
1	B	193	GLN
1	C	49	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	92	ASN
1	B	136	ASN
1	D	92	ASN
1	B	52	HIS
1	D	96	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TMP	B	257	-	19,22,22	2.69	7 (36%)	23,33,33	3.38	8 (34%)
3	TMP	D	257	2	19,22,22	1.30	3 (15%)	23,33,33	2.95	4 (17%)
3	TMP	A	257	2	19,22,22	1.57	4 (21%)	23,33,33	2.21	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TMP	B	257	-	-	5/7/22/22	0/2/2/2
3	TMP	D	257	2	-	2/7/22/22	0/2/2/2
3	TMP	A	257	2	-	6/7/22/22	0/2/2/2

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	257	TMP	C1'-N1	-5.34	1.33	1.49
3	B	257	TMP	C5'-C4'	-5.28	1.35	1.51
3	B	257	TMP	O5'-C5'	4.90	1.63	1.44
3	A	257	TMP	C4-C5	4.43	1.51	1.41
3	B	257	TMP	C2-N3	-3.82	1.30	1.38

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	257	TMP	O4'-C4'-C5'	9.31	139.99	109.37
3	A	257	TMP	C4-N3-C2	8.96	122.71	115.14
3	D	257	TMP	C4-N3-C2	8.60	122.40	115.14
3	D	257	TMP	C2'-C1'-N1	8.01	132.74	114.27

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	257	TMP	C5-C6-N1	-6.89	114.77	122.19

There are no chirality outliers.

5 of 13 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	B	257	TMP	C5'-O5'-P-O2P
3	B	257	TMP	C5'-O5'-P-O3P
3	B	257	TMP	O4'-C4'-C5'-O5'
3	D	257	TMP	O4'-C1'-N1-C6
3	A	257	TMP	C5'-O5'-P-O2P


There are no ring outliers.

3 monomers are involved in 45 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	257	TMP	20	0
3	D	257	TMP	15	0
3	A	257	TMP	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	178/261 (68%)	0.07	3 (1%) 70 72	21, 31, 45, 49	0
1	B	180/261 (68%)	0.31	10 (5%) 24 27	21, 32, 49, 59	0
1	C	177/261 (67%)	0.26	8 (4%) 33 36	25, 36, 52, 66	0
1	D	176/261 (67%)	0.23	6 (3%) 45 48	23, 34, 53, 68	0
All	All	711/1044 (68%)	0.22	27 (3%) 40 43	21, 33, 50, 68	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	197	GLY	8.6
1	C	197	GLY	7.6
1	B	196	ARG	6.8
1	B	197	GLY	6.1
1	B	176	PHE	5.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	TMP	B	257	21/21	0.57	0.40	49,70,88,89	0
3	TMP	D	257	21/21	0.72	0.38	33,61,65,67	0
3	TMP	A	257	21/21	0.77	0.40	20,20,20,20	0
2	MG	A	999	1/1	0.94	0.09	28,28,28,28	0
2	MG	B	999	1/1	0.95	0.08	33,33,33,33	0
2	MG	D	999	1/1	0.96	0.07	36,36,36,36	0
2	MG	C	999	1/1	0.96	0.04	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around TMP D 257:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around TMP A 257:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.