



## Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 08:07 pm BST

PDB ID : 6I49  
Title : Structure of *P. aeruginosa* LpxC with compound 17a: (2R)-N-Hydroxy-2-methyl-2-(methylsulfonyl)-4(6((4(morpholinomethyl)phenyl)ethynyl)-3-oxo-1H-pyrrolo[1,2-c]imidazol-2(3H)yl)butanamide  
Authors : Surivet, J.-P.; Panchaud, P.; Specklin, J.-L.; Diethelm, S.; Blumstein, A.-C.; Gauvin, J.-C.; Jacob, L.; Masse, F.; Mathieu, G.; Mirre, A.; Schmitt, C.; Enderlin-Paput, M.; Lange, R.; Bur, D.; Tidten-Luksch, N.; Gnerre, C.; Seeland, S.; Hermann, C.; Locher, H.H.; Seiler, P.; Mac Sweeney, A.; Hub-schwerlen, C.; Ritz, D.; Rueedi, G.  
Deposited on : 2018-11-09  
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	299	<div> <div></div> <div>88%12%</div> </div>
1	BBB	299	<div> <div>%</div> <div>86%13%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9470 atoms, of which 4644 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-3-O-acyl-N-acetylglucosamine deacetylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AAA	299	Total	C	H	N	O	S	99	3	0
			4612	1465	2311	396	433	7			
1	BBB	298	Total	C	H	N	O	S	103	4	0
			4570	1453	2281	393	437	6			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	40	SER	CYS	engineered mutation	UNP B7UZI4
BBB	40	SER	CYS	engineered mutation	UNP B7UZI4

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	BBB	1	Total	Zn	0	0
			1	1		
2	AAA	1	Total	Zn	0	0
			1	1		

- Molecule 3 is (2 {R})-2-methyl-2-methylsulfonyl-4-[6-[2-[4-(morpholin-4-ylmethyl)phenyl]ethynyl]-3-oxidanylidene-1 {H}-pyrrolo[1,2-c]imidazol-2-yl]- {N}-oxidanyl-butanamide (three-letter code: H2Z) (formula: C<sub>25</sub>H<sub>30</sub>N<sub>4</sub>O<sub>6</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	AAA	1	Total	C	H	N	O	S	0	0
			66	25	30	4	6	1		
3	BBB	1	Total	C	H	N	O	S	1	0
			52	21	22	3	5	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	76	Total	O	0	0
			76	76		
4	BBB	92	Total	O	0	0
			92	92		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UDP-3-O-acyl-N-acetylglucosamine deacetylase



- Molecule 1: UDP-3-O-acyl-N-acetylglucosamine deacetylase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	167.31 Å   89.18 Å   35.38 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	43.12 – 1.94 43.09 – 1.94	Depositor EDS
% Data completeness (in resolution range)	98.2 (43.12-1.94) 98.2 (43.09-1.94)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.73 (at 1.94 Å)	Xtriage
Refinement program	REFMAC 5.8.0238 2018/15/10	Depositor
R, $R_{free}$	0.183   ,   0.243 0.191   ,   0.245	Depositor DCC
$R_{free}$ test set	1982 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.1	Xtriage
Anisotropy	0.120	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 36.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9470	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, H2Z

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	AAA	1.04	2/2341 (0.1%)	1.18	5/3171 (0.2%)
1	BBB	1.06	2/2334 (0.1%)	1.18	6/3162 (0.2%)
All	All	1.05	4/4675 (0.1%)	1.18	11/6333 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	BBB	49	GLU	CD-OE2	-8.40	1.16	1.25
1	BBB	282	GLU	CD-OE1	-5.96	1.19	1.25
1	AAA	77	GLU	C-O	-5.60	1.12	1.23
1	AAA	139	GLU	CD-OE1	5.06	1.31	1.25

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	187	ARG	NE-CZ-NH2	10.67	125.64	120.30
1	BBB	130	ARG	NE-CZ-NH2	-8.49	116.06	120.30
1	AAA	187	ARG	NE-CZ-NH1	-7.91	116.35	120.30
1	AAA	258	ARG	NE-CZ-NH2	-7.79	116.40	120.30
1	BBB	130	ARG	NE-CZ-NH1	7.45	124.03	120.30
1	BBB	1	MET	CG-SD-CE	-7.38	88.39	100.20
1	AAA	258	ARG	NE-CZ-NH1	7.30	123.95	120.30
1	BBB	133	ARG	NE-CZ-NH2	7.02	123.81	120.30
1	AAA	221	ARG	CG-CD-NE	6.47	125.39	111.80
1	BBB	5	ARG	CG-CD-NE	-5.57	100.11	111.80
1	BBB	277	ASP	CB-CG-OD2	-5.26	113.57	118.30

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2301	2311	2276	26	0
1	BBB	2289	2281	2247	23	0
2	AAA	1	0	0	0	0
2	BBB	1	0	0	0	0
3	AAA	36	30	0	1	0
3	BBB	30	22	0	1	0
4	AAA	76	0	0	2	0
4	BBB	92	0	0	3	0
All	All	4826	4644	4523	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:147[A]:VAL:HG21	1:AAA:256:GLU:OE1	1.66	0.94
1:BBB:88:ILE:HD11	1:BBB:114:ILE:HG21	1.49	0.94
1:AAA:130:ARG:HD2	4:AAA:471:HOH:O	1.79	0.83
1:BBB:19:HIS:HE1	3:BBB:302:H2Z:O2	1.66	0.78
1:AAA:83[A]:MET:SD	1:AAA:88:ILE:HD12	2.23	0.77
1:BBB:261:LYS:CB	4:BBB:491:HOH:O	2.33	0.76
1:AAA:147[A]:VAL:CG2	1:AAA:256:GLU:OE1	2.35	0.74
1:AAA:147[A]:VAL:HG22	1:AAA:256:GLU:HB3	1.77	0.66
1:AAA:134:GLU:OE1	1:BBB:169:THR:HA	1.96	0.66
1:BBB:174:VAL:HG13	1:BBB:179:THR:OG1	1.97	0.65
1:BBB:142:LYS:HE3	1:BBB:261:LYS:O	1.96	0.65
1:BBB:145:VAL:CG1	1:BBB:258:ARG:HB2	2.28	0.64
1:BBB:242:ALA:O	1:BBB:246:LEU:HG	1.98	0.63
1:AAA:37:ILE:HD12	1:AAA:83[A]:MET:CE	2.31	0.60
1:BBB:160:PHE:H	1:BBB:170:GLN:HE22	1.49	0.60
1:BBB:130:ARG:HD2	4:BBB:488:HOH:O	2.03	0.58
1:AAA:130:ARG:CD	4:AAA:471:HOH:O	2.45	0.57
1:BBB:83[A]:MET:SD	1:BBB:88:ILE:HD12	2.44	0.57
1:AAA:174:VAL:HG13	1:AAA:179:THR:OG1	2.06	0.55

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:197:ILE:O	1:BBB:201:ARG:HG3	2.07	0.55
1:AAA:129:ILE:HG12	1:AAA:251:ASN:HB2	1.89	0.54
1:AAA:37:ILE:HD12	1:AAA:83[A]:MET:HE2	1.90	0.52
1:AAA:232:ASP:OD1	1:AAA:236:LYS:HD2	2.10	0.52
1:BBB:240:LEU:HD23	1:BBB:240:LEU:C	2.30	0.52
1:AAA:145:VAL:CG1	1:AAA:258:ARG:HB2	2.40	0.51
1:AAA:11:ILE:HB	1:AAA:28:LEU:HB2	1.92	0.51
1:AAA:147[A]:VAL:CG2	1:AAA:256:GLU:HB3	2.40	0.51
1:AAA:240:LEU:C	1:AAA:240:LEU:HD23	2.30	0.51
3:AAA:302:H2Z:C11	3:AAA:302:H2Z:C19	2.91	0.48
1:AAA:88:ILE:HD11	1:AAA:114:ILE:HG21	1.95	0.47
1:AAA:153:LYS:HE2	4:BBB:421:HOH:O	2.15	0.47
1:AAA:182:VAL:O	1:AAA:186:SER:OG	2.29	0.45
1:BBB:137:VAL:HG13	1:BBB:273[A]:THR:HG21	1.98	0.45
1:BBB:162:HIS:CE1	1:BBB:164:VAL:HG23	2.51	0.45
1:AAA:37:ILE:HD12	1:AAA:83[A]:MET:HE3	1.98	0.44
1:BBB:83[A]:MET:SD	1:BBB:88:ILE:CD1	3.06	0.43
1:BBB:153:LYS:O	1:BBB:256:GLU:HA	2.19	0.43
1:AAA:45:ASP:HA	1:AAA:46:PRO:HA	1.88	0.43
1:BBB:145:VAL:HG13	1:BBB:258:ARG:HB2	1.98	0.43
1:AAA:120:GLN:HA	1:AAA:120:GLN:NE2	2.35	0.42
1:AAA:64:THR:HG21	1:AAA:248:LEU:HD11	2.02	0.42
1:BBB:102:ILE:HG13	1:BBB:102:ILE:O	2.21	0.41
1:BBB:27:THR:HB	1:BBB:94:GLU:HB2	2.01	0.41
1:BBB:83[B]:MET:HE2	1:BBB:88:ILE:HD12	2.03	0.41
1:AAA:11:ILE:HG13	1:AAA:117:ALA:HB2	2.01	0.40
1:BBB:191:PHE:HA	1:BBB:215:ILE:O	2.21	0.40
1:AAA:153:LYS:O	1:AAA:256:GLU:HA	2.22	0.40
1:AAA:7:LEU:HD13	1:AAA:11:ILE:CD1	2.50	0.40
1:BBB:83[B]:MET:HA	1:BBB:88:ILE:HD12	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	300/299 (100%)	290 (97%)	8 (3%)	2 (1%)	22	11
1	BBB	300/299 (100%)	288 (96%)	12 (4%)	0	100	100
All	All	600/598 (100%)	578 (96%)	20 (3%)	2 (0%)	41	32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	166	ARG
1	AAA	297	MET

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	241/253 (95%)	239 (99%)	2 (1%)	81	78
1	BBB	240/253 (95%)	236 (98%)	4 (2%)	60	49
All	All	481/506 (95%)	475 (99%)	6 (1%)	69	64

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	40	SER
1	AAA	171	GLN
1	BBB	8	LYS
1	BBB	136	SER
1	BBB	171	GLN
1	BBB	204	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	H2Z	BBB	302	2	28,32,39	3.97	14 (50%)	26,48,57	2.73	4 (15%)
3	H2Z	AAA	302	2	35,39,39	3.57	18 (51%)	35,57,57	2.28	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H2Z	BBB	302	2	-	1/20/40/52	0/3/3/4
3	H2Z	AAA	302	2	-	2/24/52/52	0/4/4/4

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	BBB	302	H2Z	C24-S	-11.13	1.62	1.76

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	BBB	302	H2Z	C4-N2	-9.27	1.18	1.36
3	BBB	302	H2Z	C1-C2	-8.62	1.40	1.53
3	AAA	302	H2Z	C15-C14	7.36	1.64	1.51
3	AAA	302	H2Z	C3-N	7.23	1.57	1.46
3	AAA	302	H2Z	C15-N1	6.62	1.60	1.47
3	AAA	302	H2Z	C5-C6	6.60	1.49	1.41
3	AAA	302	H2Z	C24-S	-6.44	1.68	1.76
3	AAA	302	H2Z	C3-C4	5.46	1.61	1.50
3	AAA	302	H2Z	C4-N2	-5.33	1.26	1.36
3	BBB	302	H2Z	O2-C21	5.04	1.28	1.22
3	AAA	302	H2Z	O5-S	-4.87	1.40	1.44
3	AAA	302	H2Z	C23-N3	-4.81	1.25	1.34
3	BBB	302	H2Z	C21-N	-4.38	1.30	1.37
3	BBB	302	H2Z	C9-C8	-4.25	1.34	1.44
3	AAA	302	H2Z	C1-C2	-4.16	1.47	1.53
3	AAA	302	H2Z	O-S	4.01	1.47	1.44
3	BBB	302	H2Z	C23-N3	-3.51	1.27	1.34
3	BBB	302	H2Z	O5-S	-3.40	1.41	1.44
3	AAA	302	H2Z	C6-C7	-3.34	1.36	1.44
3	BBB	302	H2Z	C5-C6	3.31	1.45	1.41
3	BBB	302	H2Z	C12-C13	3.29	1.44	1.38
3	AAA	302	H2Z	C12-C13	3.07	1.44	1.38
3	BBB	302	H2Z	C11-C10	3.02	1.44	1.38
3	BBB	302	H2Z	C3-C4	2.60	1.55	1.50
3	AAA	302	H2Z	O3-N3	-2.51	1.33	1.40
3	AAA	302	H2Z	C22-C	-2.44	1.50	1.53
3	AAA	302	H2Z	C11-C10	2.38	1.43	1.38
3	AAA	302	H2Z	C19-N1	-2.17	1.40	1.46
3	AAA	302	H2Z	C9-C8	-2.15	1.39	1.44
3	BBB	302	H2Z	C12-C9	2.10	1.43	1.39
3	BBB	302	H2Z	C3-N	2.06	1.50	1.46

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	BBB	302	H2Z	O-S-O5	-10.19	109.63	118.10
3	AAA	302	H2Z	O-S-O5	-9.74	110.00	118.10
3	BBB	302	H2Z	C3-N-C2	-7.68	114.88	123.26
3	AAA	302	H2Z	C3-N-C2	-7.40	115.19	123.26
3	BBB	302	H2Z	C2-N-C21	3.92	131.78	123.07
3	AAA	302	H2Z	C2-N-C21	3.82	131.56	123.07
3	AAA	302	H2Z	C20-N2-C21	2.09	132.74	127.82

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	BBB	302	H2Z	C20-N2-C21	2.04	132.61	127.82

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	AAA	302	H2Z	C6-C7-C8-C9
3	BBB	302	H2Z	C6-C7-C8-C9
3	AAA	302	H2Z	C14-C15-N1-C19

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	BBB	302	H2Z	1	0
3	AAA	302	H2Z	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	299/299 (100%)	-0.14	0 100 100	15, 24, 43, 66	0
1	BBB	298/299 (99%)	-0.14	2 (0%) 87 91	12, 24, 41, 59	0
All	All	597/598 (99%)	-0.14	2 (0%) 94 96	12, 24, 42, 66	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	295	SER	2.4
1	BBB	204	ASN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	H2Z	AAA	302	36/36	0.97	0.13	17,23,70,76	0
3	H2Z	BBB	302	30/36	0.98	0.11	15,23,40,46	1
2	ZN	AAA	301	1/1	1.00	0.09	16,16,16,16	0

*Continued on next page...*






*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	BBB	301	1/1	1.00	0.07	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around H2Z AAA 302:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around H2Z BBB 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.