



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 10:12 pm BST

PDB ID : 2I5F
Title : Crystal structure of the C-terminal PH domain of pleckstrin in complex with D-myo-Ins(1,2,3,5,6)P5
Authors : Jackson, S.G.; Haslam, R.J.; Junop, M.S.
Deposited on : 2006-08-24
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	109	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 999 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pleckstrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	99	843	547	145	147	4	0	7	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	239	GLY	-	CLONING ARTIFACT	UNP P08567
A	240	SER	-	CLONING ARTIFACT	UNP P08567
A	241	PHE	-	CLONING ARTIFACT	UNP P08567
A	242	THR	-	CLONING ARTIFACT	UNP P08567
A	243	GLY	-	CLONING ARTIFACT	UNP P08567

- Molecule 2 is (1R,2R,3R,4R,5S,6S)-6-HYDROXYCYCLOHEXANE-1,2,3,4,5-PENTAYL PENTAKIS[DIHYDROGEN (PHOSPHATE)] (three-letter code: 5IP) (formula: $C_6H_{17}O_{21}P_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			32	6	21	5		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	124	Total	O	0	0
			124	124		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pleckstrin

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	32.15Å 47.70Å 64.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.70 – 1.35 28.73 – 1.35	Depositor EDS
% Data completeness (in resolution range)	97.7 (28.70-1.35) 97.7 (28.73-1.35)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.179 , 0.208 0.267 , 0.289	Depositor DCC
R_{free} test set	1084 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	15.2	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	999	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 5IP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.46	8/882 (0.9%)	1.56	16/1186 (1.3%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	274	TYR	CE2-CZ	-8.66	1.27	1.38
1	A	322	GLU	CB-CG	7.08	1.65	1.52
1	A	328	GLN	CB-CG	-5.75	1.37	1.52
1	A	337	GLU	CD-OE2	5.72	1.31	1.25
1	A	284	GLU	CG-CD	5.71	1.60	1.51
1	A	335	ARG	CZ-NH2	5.22	1.39	1.33
1	A	335	ARG	NE-CZ	-5.20	1.26	1.33
1	A	328	GLN	CD-NE2	-5.11	1.20	1.32

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	258	ARG	NE-CZ-NH2	-13.97	113.31	120.30
1	A	269	ARG	NE-CZ-NH1	10.69	125.64	120.30
1	A	258	ARG	NE-CZ-NH1	9.89	125.25	120.30
1	A	335	ARG	NE-CZ-NH1	8.47	124.53	120.30
1	A	269	ARG	NE-CZ-NH2	-7.65	116.48	120.30
1	A	266	PHE	CB-CG-CD2	-6.98	115.91	120.80
1	A	287[A]	LEU	CB-CG-CD1	-6.77	99.49	111.00
1	A	287[B]	LEU	CB-CG-CD1	-6.77	99.49	111.00
1	A	278	TYR	CB-CG-CD1	-5.93	117.44	121.00
1	A	278	TYR	CB-CG-CD2	5.79	124.48	121.00
1	A	278	TYR	CG-CD1-CE1	5.49	125.69	121.30
1	A	266	PHE	CB-CG-CD1	5.38	124.56	120.80
1	A	257	ARG	NE-CZ-NH2	-5.34	117.63	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	293	ARG	NE-CZ-NH2	-5.31	117.64	120.30
1	A	257	ARG	NE-CZ-NH1	5.16	122.88	120.30
1	A	325	TYR	CB-CG-CD1	5.02	124.01	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	843	0	874	36	0
2	A	32	0	6	2	0
3	A	124	0	0	11	0
All	All	999	0	880	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266:PHE:HD2	1:A:275[B]:LEU:HD11	1.28	0.98
1:A:347:ARG:HG3	1:A:347:ARG:HH11	1.31	0.95
1:A:279:ASP:HB2	3:A:68:HOH:O	1.66	0.93
1:A:344[A]:MET:HG2	3:A:125:HOH:O	1.66	0.93
1:A:280:PRO:HD2	3:A:68:HOH:O	1.73	0.88
1:A:287[B]:LEU:HD11	3:A:4:HOH:O	1.75	0.86
1:A:344[A]:MET:HG2	3:A:95:HOH:O	1.79	0.82
1:A:347:ARG:CG	1:A:347:ARG:HH11	1.93	0.81
1:A:344[A]:MET:CG	3:A:95:HOH:O	2.31	0.79
1:A:257:ARG:HB2	2:A:550:5IP:O45	1.84	0.77
1:A:287[B]:LEU:CD1	3:A:4:HOH:O	2.38	0.70
1:A:278:TYR:HA	1:A:287[B]:LEU:HD13	1.77	0.66
1:A:344[A]:MET:CE	3:A:95:HOH:O	2.43	0.65
1:A:344[A]:MET:HE2	3:A:125:HOH:O	1.97	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:LEU:CD2	1:A:263:VAL:HG22	2.28	0.64
1:A:344[A]:MET:HE3	3:A:95:HOH:O	1.98	0.61
1:A:266:PHE:CD2	1:A:275[B]:LEU:HD11	2.21	0.61
1:A:252:LEU:HD21	1:A:263:VAL:HG22	1.84	0.58
1:A:252:LEU:HD21	1:A:263:VAL:CG2	2.40	0.52
1:A:241:PHE:HE2	1:A:274:TYR:CE1	2.28	0.52
1:A:252:LEU:CD2	1:A:263:VAL:CG2	2.89	0.51
1:A:343:GLN:HE21	1:A:347:ARG:HH21	1.58	0.50
1:A:252:LEU:HD23	1:A:263:VAL:HG22	1.94	0.49
1:A:265[A]:LYS:HE2	1:A:282:GLY:HA3	1.96	0.47
1:A:312:GLU:H	1:A:328:GLN:NE2	2.12	0.47
1:A:347:ARG:NH1	1:A:347:ARG:HG3	2.10	0.47
1:A:280:PRO:CD	3:A:68:HOH:O	2.48	0.44
1:A:252:LEU:HD23	1:A:263:VAL:HA	1.99	0.43
1:A:275[B]:LEU:HB2	1:A:292:LEU:HD11	2.00	0.43
1:A:312:GLU:O	1:A:328:GLN:HG3	2.21	0.41
1:A:257:ARG:HB2	2:A:550:5IP:P5	2.60	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/109 (94%)	100 (98%)	2 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	91/92 (99%)	88 (97%)	3 (3%)	38 7

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	311	GLU
1	A	326	PHE
1	A	347	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	328	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	5IP	A	550	-	32,32,32	2.34	9 (28%)	48,53,53	2.94	19 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5IP	A	550	-	-	7/25/49/49	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	550	5IP	P5-O25	9.05	1.89	1.54
2	A	550	5IP	C4-C5	4.15	1.63	1.52
2	A	550	5IP	O15-C5	-3.87	1.30	1.44
2	A	550	5IP	P6-O16	3.14	1.65	1.59
2	A	550	5IP	O14-C4	3.09	1.50	1.43
2	A	550	5IP	P6-O36	2.91	1.59	1.50
2	A	550	5IP	C2-C3	2.64	1.57	1.52
2	A	550	5IP	C2-C1	2.60	1.57	1.52
2	A	550	5IP	P1-O11	2.35	1.63	1.59

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	550	5IP	O12-C2-C1	-10.14	84.78	108.69
2	A	550	5IP	C3-C2-C1	8.18	128.32	110.41
2	A	550	5IP	C4-C5-C6	6.49	126.49	111.66
2	A	550	5IP	O35-P5-O15	-5.86	79.72	105.99
2	A	550	5IP	O11-C1-C2	4.91	120.26	108.69
2	A	550	5IP	O35-P5-O45	3.60	124.79	110.68
2	A	550	5IP	O26-P6-O36	3.57	124.67	110.68
2	A	550	5IP	O13-C3-C4	-3.55	100.40	108.66
2	A	550	5IP	O13-C3-C2	3.43	116.77	108.69
2	A	550	5IP	O15-C5-C6	2.83	115.35	108.69
2	A	550	5IP	O14-C4-C3	-2.80	102.51	109.94
2	A	550	5IP	O15-P5-O45	2.73	119.94	109.39
2	A	550	5IP	C2-C1-C6	2.46	115.79	110.41
2	A	550	5IP	O22-P2-O12	2.42	116.85	105.99
2	A	550	5IP	O23-P3-O13	-2.35	95.45	105.99
2	A	550	5IP	O26-P6-O46	-2.28	98.92	107.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	550	5IP	O16-P6-O36	-2.19	100.93	109.39
2	A	550	5IP	O21-P1-O11	-2.12	96.51	105.99
2	A	550	5IP	O11-P1-O41	-2.08	101.35	109.39

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	550	5IP	C2-O12-P2-O22
2	A	550	5IP	C2-C1-O11-P1
2	A	550	5IP	C4-C3-O13-P3
2	A	550	5IP	C2-O12-P2-O42
2	A	550	5IP	C2-C3-O13-P3
2	A	550	5IP	C6-O16-P6-O46
2	A	550	5IP	C6-O16-P6-O26

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	550	5IP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.




6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 5IP A 550:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.