



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 06:23 am BST

PDB ID : 3I75
Title : Antibody Structure
Authors : Isaacs, N.W.; Riboldi-Tunnicliffe, A.
Deposited on : 2009-07-08
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	
2	B	225	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3648 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antibody heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	3	0
			1661	1035	282	337	7			

- Molecule 2 is a protein called Antibody light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	223	Total	C	N	O	S	0	2	0
			1692	1075	270	338	9			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	122	Total 122	O 122	0	0
6	B	130	Total 130	O 130	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Antibody heavy chain



• Molecule 2: Antibody light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	129.66 Å 129.66 Å 74.50 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	64.83 – 1.95 64.83 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.4 (64.83-1.95) 96.3 (64.83-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 1.95 Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.180 , 0.213 0.189 , 0.214	Depositor DCC
R_{free} test set	2571 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3648	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.36	10/1713 (0.6%)	1.20	13/2323 (0.6%)
2	B	1.43	13/1749 (0.7%)	1.14	6/2392 (0.3%)
All	All	1.39	23/3462 (0.7%)	1.17	19/4715 (0.4%)

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	199	TRP	CB-CG	-8.65	1.34	1.50
1	A	123	GLU	CB-CG	7.23	1.65	1.52
2	B	199	TRP	CG-CD1	-6.93	1.27	1.36
1	A	212	ALA	CA-CB	6.91	1.67	1.52
2	B	61	ASN	CB-CG	6.76	1.66	1.51
2	B	134	TYR	CD1-CE1	6.68	1.49	1.39
2	B	147	VAL	CB-CG1	-6.58	1.39	1.52
1	A	88	CYS	CB-SG	-6.04	1.72	1.82
2	B	202	GLU	CD-OE2	5.75	1.31	1.25
2	B	114	TYR	CD2-CE2	5.74	1.48	1.39
2	B	89	GLU	CD-OE1	-5.69	1.19	1.25
2	B	217	VAL	CB-CG2	-5.61	1.41	1.52
1	A	147	LYS	CD-CE	5.46	1.64	1.51
2	B	65	LYS	CB-CG	-5.31	1.38	1.52
1	A	160	LEU	CG-CD2	5.29	1.71	1.51
2	B	12	VAL	CB-CG1	-5.25	1.41	1.52
1	A	10	SER	CB-OG	-5.22	1.35	1.42
1	A	1	ASP	CB-CG	5.22	1.62	1.51
1	A	155	ARG	CG-CD	-5.20	1.39	1.51
2	B	159	GLU	CD-OE1	5.09	1.31	1.25
1	A	209	PHE	CD1-CE1	-5.07	1.29	1.39
2	B	84	ARG	NE-CZ	5.05	1.39	1.33
1	A	18	LYS	CD-CE	5.04	1.63	1.51

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	225	ASP	CB-CG-OD2	9.07	126.46	118.30
2	B	31	ASP	CB-CG-OD2	8.82	126.24	118.30
1	A	1	ASP	CB-CG-OD2	8.81	126.23	118.30
1	A	82	ASP	CB-CG-OD2	7.86	125.37	118.30
1	A	108	ARG	NE-CZ-NH1	7.45	124.03	120.30
1	A	11	LEU	CA-CB-CG	7.24	131.95	115.30
1	A	165	ASP	CB-CG-OD2	7.06	124.65	118.30
1	A	92	ASP	CB-CG-OD2	6.85	124.46	118.30
1	A	110	ASP	CB-CG-OD2	6.59	124.23	118.30
2	B	188	LEU	CA-CB-CG	6.15	129.45	115.30
1	A	108	ARG	NE-CZ-NH2	-5.84	117.38	120.30
2	B	67	ARG	NE-CZ-NH2	-5.79	117.41	120.30
2	B	4	LEU	CB-CG-CD1	5.69	120.67	111.00
1	A	207	LYS	CD-CE-NZ	-5.52	99.00	111.70
1	A	155	ARG	NE-CZ-NH2	-5.38	117.61	120.30
2	B	113	ASP	CB-CG-OD2	5.34	123.11	118.30
1	A	143	ASP	CB-CG-OD2	5.32	123.09	118.30
1	A	93	ASP	CB-CG-OD1	5.29	123.06	118.30
1	A	10	SER	CB-CA-C	-5.10	100.41	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1661	0	1608	18	0
2	B	1692	0	1632	17	0
3	A	10	0	0	0	0
3	B	20	0	0	1	0
4	A	1	0	0	0	0
5	B	12	0	16	0	0
6	A	122	0	0	2	0
6	B	130	0	0	5	0
All	All	3648	0	3256	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:81[B]:MET:CE	2:B:81[B]:MET:SD	2.02	1.48
2:B:1:GLU:HA	6:B:252:HOH:O	1.55	1.07
2:B:1:GLU:CA	6:B:252:HOH:O	2.11	0.95
1:A:79:GLU:OE1	1:A:81:GLU:OE1	1.89	0.91
2:B:1:GLU:CB	6:B:252:HOH:O	2.30	0.79
2:B:61:ASN:HD21	2:B:63:ASN:HB2	1.55	0.69
1:A:108:ARG:HD3	1:A:109:ALA:O	1.92	0.69
2:B:61:ASN:HB2	6:B:313:HOH:O	1.96	0.65
1:A:191:THR:HG23	6:A:297:HOH:O	1.99	0.62
2:B:61:ASN:HD22	2:B:64:PHE:HD2	1.47	0.62
2:B:173:GLY:N	3:B:230:SO4:O3	2.31	0.60
1:A:157:ASN:H	1:A:157:ASN:HD22	1.51	0.58
1:A:55[B]:ARG:HB3	1:A:57:ASP:OD1	2.04	0.58
2:B:61:ASN:ND2	2:B:63:ASN:HB2	2.19	0.55
1:A:147:LYS:NZ	6:A:320:HOH:O	2.41	0.52
2:B:1:GLU:HB2	6:B:252:HOH:O	2.03	0.51
1:A:119:PRO:HG2	2:B:224:ARG:HD2	1.93	0.51
1:A:37:GLN:HB2	1:A:47:LEU:HD11	1.93	0.50
1:A:78:LEU:HD13	1:A:106:LEU:CD2	2.42	0.49
1:A:38[B]:HIS:CE1	1:A:42:LYS:O	2.65	0.49
1:A:157:ASN:ND2	1:A:157:ASN:H	2.08	0.49
1:A:57:ASP:N	1:A:57:ASP:OD1	2.35	0.48
1:A:38[B]:HIS:HE1	1:A:42:LYS:O	1.95	0.48
1:A:55[A]:ARG:HB3	1:A:57:ASP:OD1	2.14	0.47
2:B:81[B]:MET:CE	2:B:81[B]:MET:CG	2.91	0.47
2:B:199:TRP:CG	2:B:200:PRO:HA	2.50	0.46
2:B:213:SER:O	2:B:214:SER:HB2	2.15	0.46
1:A:203:SER:HB2	1:A:204:PRO:HD2	1.96	0.46
2:B:83:LEU:HB3	2:B:86:LEU:HD21	1.98	0.46
2:B:48:ILE:HG21	2:B:81[A]:MET:HE2	1.99	0.44
1:A:46:LEU:HD23	1:A:55[B]:ARG:HD2	1.99	0.44
2:B:36:TRP:CG	2:B:81[A]:MET:HE3	2.55	0.42
1:A:161:ASN:ND2	1:A:177:SER:OG	2.53	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/212 (100%)	210 (99%)	3 (1%)	0	100	100
2	B	221/225 (98%)	218 (99%)	3 (1%)	0	100	100
All	All	434/437 (99%)	428 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/189 (102%)	182 (95%)	10 (5%)	23	10
2	B	190/188 (101%)	183 (96%)	7 (4%)	34	22
All	All	382/377 (101%)	365 (96%)	17 (4%)	30	15

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	LEU
1	A	38[A]	HIS
1	A	38[B]	HIS
1	A	60	SER
1	A	108	ARG
1	A	123	GLU
1	A	142	LYS
1	A	157	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	199	LYS
1	A	206	VAL
2	B	4	LEU
2	B	12	VAL
2	B	61	ASN
2	B	65	LYS
2	B	109	SER
2	B	182	GLN
2	B	224	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	157	ASN
1	A	161	ASN
2	B	61	ASN
2	B	182	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	229	-	4,4,4	0.27	0	6,6,6	0.29	0
5	GOL	B	227	-	5,5,5	0.65	0	5,5,5	1.58	1 (20%)
3	SO4	B	228	-	4,4,4	0.55	0	6,6,6	0.84	0
5	GOL	B	226	-	5,5,5	0.66	0	5,5,5	1.51	1 (20%)
3	SO4	A	213	-	4,4,4	0.33	0	6,6,6	1.05	0
3	SO4	A	214	-	4,4,4	0.44	0	6,6,6	0.23	0
3	SO4	B	230	-	4,4,4	0.31	0	6,6,6	0.53	0
3	SO4	B	231	-	4,4,4	0.49	0	6,6,6	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	227	-	-	3/4/4/4	-
5	GOL	B	226	-	-	0/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	226	GOL	C3-C2-C1	-2.56	101.77	111.70
5	B	227	GOL	O2-C2-C3	2.06	118.20	109.12

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	227	GOL	O1-C1-C2-C3
5	B	227	GOL	C1-C2-C3-O3
5	B	227	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	230	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	212/212 (100%)	-0.25	1 (0%) 91 94	24, 37, 52, 58	2 (0%)
2	B	223/225 (99%)	-0.32	2 (0%) 84 89	25, 35, 54, 73	0
All	All	435/437 (99%)	-0.29	3 (0%) 87 92	24, 36, 53, 73	2 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	202	THR	3.0
2	B	142	ALA	2.6
2	B	42	VAL	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	B	227	6/6	0.86	0.19	55,59,59,60	0
3	SO4	A	214	5/5	0.89	0.27	90,91,93,93	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	B	231	5/5	0.90	0.34	91,93,93,94	0
5	GOL	B	226	6/6	0.92	0.15	30,44,49,56	0
3	SO4	B	230	5/5	0.93	0.30	93,93,94,94	0
3	SO4	B	228	5/5	0.93	0.22	70,72,73,75	0
3	SO4	A	213	5/5	0.95	0.12	50,60,60,62	0
3	SO4	B	229	5/5	0.96	0.23	76,76,76,76	5
4	CL	A	215	1/1	0.99	0.04	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.