



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 06:50 am BST

PDB ID : 4IBQ  
Title : Human p53 core domain with hot spot mutation R273C  
Authors : Rozenberg, H.; Eldar, A.; Diskin-Posner, Y.; Shakked, Z.  
Deposited on : 2012-12-09  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	200	<div> <div>76%</div> <div>20%</div> <div>.</div> </div>
1	B	200	<div> <div>81%</div> <div>15%</div> <div>..</div> </div>
1	C	200	<div> <div>81%</div> <div>14%</div> <div>..</div> </div>
1	D	200	<div> <div>77%</div> <div>20%</div> <div>..</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7037 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellular tumor antigen p53.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	192	Total	C	N	O	S	0	7	0
			1540	946	280	293	21			
1	B	193	Total	C	N	O	S	0	5	0
			1529	943	275	292	19			
1	C	192	Total	C	N	O	S	0	6	0
			1518	937	270	291	20			
1	D	193	Total	C	N	O	S	0	2	0
			1502	927	270	287	18			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	273	CYS	ARG	ENGINEERED MUTATION	UNP P04637
B	273	CYS	ARG	ENGINEERED MUTATION	UNP P04637
C	273	CYS	ARG	ENGINEERED MUTATION	UNP P04637
D	273	CYS	ARG	ENGINEERED MUTATION	UNP P04637

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		
2	D	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	1
			4	2	2		
4	D	1	Total	C	O	0	1
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	263	Total 269	O 269	0	6
5	B	235	Total 240	O 240	0	6
5	C	230	Total 233	O 233	0	4
5	D	189	Total 190	O 190	0	4


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cellular tumor antigen p53




- Molecule 1: Cellular tumor antigen p53



- Molecule 1: Cellular tumor antigen p53



- Molecule 1: Cellular tumor antigen p53



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.73Å 69.72Å 83.51Å 90.00° 90.10° 90.00°	Depositor
Resolution (Å)	29.14 – 1.80 29.14 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.5 (29.14-1.80) 95.9 (29.14-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.79 (at 1.79Å)	Xtriage
Refinement program	PHENIX dev_1223	Depositor
R, $R_{free}$	0.134 , 0.163 0.134 , 0.162	Depositor DCC
$R_{free}$ test set	1861 reflections (2.59%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.6	Xtriage
Anisotropy	1.141	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 35.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.067 for -k,-h,-l 0.067 for k,h,-l 0.450 for h,-k,-l	Xtriage
Reported twinning fraction	0.480 for h,-k,-l	Depositor
Outliers	0 of 71928 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7037	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.35	0/1575	0.54	0/2141
1	B	0.34	0/1568	0.51	0/2130
1	C	0.33	0/1553	0.51	0/2115
1	D	0.33	0/1542	0.51	0/2094
All	All	0.34	0/6238	0.52	0/8480

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1540	0	1461	26	0
1	B	1529	0	1460	20	0
1	C	1518	0	1431	19	0
1	D	1502	0	1437	24	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	4	0	3	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	8	0	12	3	0
5	A	269	0	0	8	0
5	B	240	0	0	10	1
5	C	233	0	0	7	1
5	D	190	0	0	7	0
All	All	7037	0	5804	87	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:238:CYS:SG	5:B:520:HOH:O	2.42	0.76
1:D:220:TYR:O	5:D:487:HOH:O	2.08	0.72
1:C:272:VAL:O	5:C:501:HOH:O	2.07	0.71
1:B:241:SER:O	5:B:595:HOH:O	2.09	0.70
1:D:202:ARG:NH1	4:D:302[A]:EDO:O2	2.26	0.68
1:A:256:THR:HG22	1:A:267:ARG:HG3	1.78	0.66
1:C:256:THR:HG22	1:C:267:ARG:HG3	1.76	0.66
1:B:248:ARG:NH1	5:B:595:HOH:O	2.29	0.65
1:B:248:ARG:NH1	5:B:516:HOH:O	2.30	0.65
1:C:239:ASN:OD1	5:C:523:HOH:O	2.15	0.64
1:D:107:TYR:OH	5:D:522:HOH:O	2.13	0.63
1:A:277[A]:CYS:SG	5:A:620:HOH:O	2.38	0.63
1:C:150[A]:THR:OG1	5:C:589:HOH:O	2.15	0.63
1:C:174:ARG:NE	5:C:568:HOH:O	2.33	0.62
1:B:132:LYS:NZ	1:B:285:GLU:OE2	2.26	0.61
1:D:271:GLU:OE2	5:D:485:HOH:O	2.15	0.61
1:B:280:ARG:NH1	5:B:467:HOH:O	2.28	0.61
1:A:222:PRO:HB2	3:A:302:ACT:H1	1.85	0.59
1:C:164:LYS:HB2	1:C:252:LEU:HD23	1.85	0.59
1:C:107:TYR:CZ	1:C:152:PRO:HD3	2.38	0.58
1:C:188:LEU:O	5:C:550:HOH:O	2.16	0.58
1:A:242:CYS:SG	5:A:483[B]:HOH:O	2.57	0.58
1:A:174[A]:ARG:NH1	5:A:502:HOH:O	2.22	0.57
1:A:263:ASN:ND2	5:A:441:HOH:O	2.16	0.57
1:D:162:ILE:HD13	1:D:254:ILE:HD11	1.86	0.56
1:A:103:TYR:CZ	1:B:243[A]:MET:HB2	2.40	0.56
1:D:133[A]:MET:HB3	1:D:272:VAL:HG22	1.88	0.56
1:D:164:LYS:O	5:D:569:HOH:O	2.18	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:SER:HA	1:A:248:ARG:HG3	1.88	0.55
1:A:119:ALA:O	1:A:122:VAL:HG22	2.06	0.55
1:A:175:ARG:HD3	1:A:237:MET:HB2	1.88	0.54
1:A:182:CYS:O	5:A:487:HOH:O	2.18	0.54
1:B:210:ASN:OD1	5:B:526:HOH:O	2.18	0.54
1:A:103:TYR:CZ	1:B:243[B]:MET:HB2	2.43	0.54
1:D:97:VAL:HG11	1:D:169:MET:HG2	1.90	0.54
1:B:256[B]:THR:HG21	5:B:469:HOH:O	2.08	0.54
1:A:174[A]:ARG:NH2	5:A:513:HOH:O	2.41	0.52
1:B:256[B]:THR:HG22	1:B:267:ARG:HG3	1.93	0.50
1:C:271:GLU:OE1	5:C:502:HOH:O	2.19	0.50
1:C:280:ARG:NH1	5:C:451:HOH:O	2.43	0.49
1:C:179:HIS:O	1:C:182:CYS:HB2	2.11	0.49
1:D:175:ARG:HD3	1:D:237:MET:HB2	1.94	0.49
1:A:168:HIS:CD2	1:A:249:ARG:HH11	2.31	0.48
1:D:108:GLY:O	1:D:110:ARG:NH1	2.45	0.48
1:C:101:LYS:O	1:C:267:ARG:HD2	2.13	0.48
1:A:136:GLN:HB2	1:A:139:LYS:HG3	1.96	0.47
1:B:268:ASN:ND2	5:B:476:HOH:O	2.43	0.47
1:B:179:HIS:O	1:B:182:CYS:HB2	2.14	0.47
1:A:123:THR:HG23	1:A:139:LYS:HB3	1.97	0.47
1:A:283:ARG:HB3	4:D:302[A]:EDO:H22	1.97	0.46
1:A:202:ARG:NH2	1:A:221:GLU:OE2	2.48	0.46
1:D:181:ARG:HA	4:D:303[A]:EDO:H12	1.98	0.46
1:C:151:PRO:HD2	1:C:220:TYR:CE2	2.50	0.46
1:D:104:GLN:NE2	5:D:544:HOH:O	2.48	0.46
1:A:110:ARG:HG3	1:A:146:TRP:HB2	1.98	0.46
1:A:133[B]:MET:SD	1:A:141[B]:CYS:SG	3.11	0.46
1:D:96:SER:HA	1:D:97:VAL:HA	1.78	0.45
1:A:288:ASN:HB2	5:A:629:HOH:O	2.17	0.44
1:B:101:LYS:O	1:B:267:ARG:HD2	2.17	0.44
1:C:193:HIS:CE1	1:C:214:HIS:HB3	2.52	0.44
1:D:96:SER:HB2	1:D:170:THR:HG22	1.99	0.44
1:B:286:GLU:C	1:B:288:ASN:H	2.20	0.43
1:A:177:PRO:HA	1:A:180:GLU:HG2	2.01	0.43
1:D:190:PRO:HB2	1:D:193:HIS:CD2	2.53	0.43
1:B:190:PRO:HA	1:B:191:PRO:HD3	1.92	0.43
1:D:151:PRO:HA	1:D:152:PRO:HD3	1.83	0.42
1:A:190:PRO:HA	1:A:191:PRO:HD3	1.94	0.42
1:D:156:ARG:NH1	5:D:415:HOH:O	2.51	0.42
1:B:119:ALA:O	1:B:122:VAL:HG22	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:VAL:HG11	1:A:218:VAL:HG11	2.01	0.41
1:A:123:THR:HA	1:A:136:GLN:OE1	2.20	0.41
1:C:150[A]:THR:HA	1:C:151:PRO:HD3	1.81	0.41
1:B:100:GLN:OE1	5:B:528[B]:HOH:O	2.22	0.41
1:C:97:VAL:HG11	1:C:169:MET:HG2	2.03	0.41
1:A:205:TYR:OH	5:A:415:HOH:O	2.21	0.41
1:B:282:ARG:NH2	5:B:475:HOH:O	2.49	0.41
1:C:97:VAL:HA	1:C:98:PRO:HD3	1.94	0.41
1:D:133[B]:MET:SD	1:D:141:CYS:HB3	2.61	0.41
1:D:198:GLU:HB2	1:D:235:ASN:OD1	2.21	0.41
1:D:239:ASN:HA	1:D:274:VAL:HB	2.02	0.41
1:D:158:ARG:HB3	1:D:256[C]:THR:CG2	2.51	0.41
1:B:110:ARG:HA	1:B:268:ASN:ND2	2.36	0.41
1:D:150:THR:HG22	5:D:483:HOH:O	2.19	0.41
1:C:224:GLU:HG3	1:C:225:VAL:HG22	2.03	0.40
1:C:175:ARG:HD3	1:C:237:MET:HB2	2.03	0.40
1:D:119:ALA:O	1:D:122:VAL:HG22	2.22	0.40
1:D:204:GLU:HB3	1:D:217:VAL:HG13	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:444:HOH:O	5:C:608:HOH:O[1_655]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/200 (98%)	195 (99%)	2 (1%)	0	100	100
1	B	196/200 (98%)	192 (98%)	3 (2%)	1 (0%)	29	15

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	196/200 (98%)	193 (98%)	3 (2%)	0	100	100
1	D	194/200 (97%)	191 (98%)	3 (2%)	0	100	100
All	All	783/800 (98%)	771 (98%)	11 (1%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	97	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/179 (98%)	169 (97%)	6 (3%)	37	22
1	B	174/179 (97%)	165 (95%)	9 (5%)	23	10
1	C	170/179 (95%)	161 (95%)	9 (5%)	22	9
1	D	169/179 (94%)	164 (97%)	5 (3%)	41	27
All	All	688/716 (96%)	659 (96%)	29 (4%)	35	15

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	SER
1	A	114	LEU
1	A	198	GLU
1	A	217	VAL
1	A	238[A]	CYS
1	A	238[B]	CYS
1	B	110	ARG
1	B	114	LEU
1	B	149	SER
1	B	180	GLU
1	B	192[A]	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	192[B]	GLN
1	B	198	GLU
1	B	224	GLU
1	B	225	VAL
1	C	124[A]	CYS
1	C	124[B]	CYS
1	C	182	CYS
1	C	192[A]	GLN
1	C	192[B]	GLN
1	C	225	VAL
1	C	238[A]	CYS
1	C	238[B]	CYS
1	C	252	LEU
1	D	114	LEU
1	D	123	THR
1	D	192	GLN
1	D	217	VAL
1	D	238	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	168	HIS
1	C	115	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	D	303[A]	-	3,3,3	0.47	0	2,2,2	0.29	0
3	ACT	A	302	-	1,3,3	1.49	0	0,3,3	0.00	-
4	EDO	D	302[A]	-	3,3,3	0.44	0	2,2,2	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	D	303[A]	-	-	0/1/1/1	-
4	EDO	D	302[A]	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	303[A]	EDO	1	0
3	A	302	ACT	1	0
4	D	302[A]	EDO	2	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	192/200 (96%)	-0.61	2 (1%) 82 80	8, 15, 31, 51	0
1	B	193/200 (96%)	-0.67	0 100 100	8, 17, 32, 38	0
1	C	192/200 (96%)	-0.65	0 100 100	8, 17, 31, 53	0
1	D	193/200 (96%)	-0.62	0 100 100	10, 19, 31, 45	1 (0%)
All	All	770/800 (96%)	-0.64	2 (0%) 94 92	8, 17, 32, 53	1 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	153	PRO	2.5
1	A	226	GLY	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ACT	A	302	4/4	0.91	0.22	30,32,35,39	0
4	EDO	D	303[A]	4/4	0.95	0.11	19,23,25,27	4
4	EDO	D	302[A]	4/4	0.98	0.09	21,21,22,26	4
2	ZN	B	301	1/1	0.99	0.05	17,17,17,17	0
2	ZN	C	301	1/1	0.99	0.06	21,21,21,21	0
2	ZN	D	301	1/1	1.00	0.06	20,20,20,20	0
2	ZN	A	301	1/1	1.00	0.09	19,19,19,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.