



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 10:52 AM BST

PDB ID : 4IBU
Title : Human p53 core domain with hot spot mutation R273C and second-site suppressor mutation T284R in sequence-specific complex with DNA
Authors : Eldar, A.; Rozenberg, H.; Diskin-Posner, Y.; Shakked, Z.
Deposited on : 2012-12-09
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	200	<div> <div>2%</div> <div>86%</div> <div>12%</div> <div>••</div> </div>
1	B	200	<div> <div>4%</div> <div>91%</div> <div>8%</div> <div>••</div> </div>
1	C	200	<div> <div>3%</div> <div>88%</div> <div>10%</div> <div>•</div> </div>
1	D	200	<div> <div>3%</div> <div>89%</div> <div>9%</div> <div>••</div> </div>
2	E	12	<div> <div>8%</div> <div>50%</div> <div>42%</div> <div>8%</div> </div>
2	F	12	<div> <div>25%</div> <div>58%</div> <div>33%</div> <div>8%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	12	<div><div></div><div>33%</div><div>42%</div><div>58%</div></div>
2	H	12	<div><div></div><div>8%</div><div>50%</div><div>50%</div></div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellular tumor antigen p53.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	S	0	16	0
			1651	1017	308	306	20			
1	B	198	Total	C	N	O	S	0	9	0
			1611	988	302	301	20			
1	C	195	Total	C	N	O	S	0	10	0
			1578	975	290	292	21			
1	D	196	Total	C	N	O	S	0	9	0
			1582	971	292	300	19			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	273	CYS	ARG	ENGINEERED MUTATION	UNP P04637
A	284	ARG	THR	ENGINEERED MUTATION	UNP P04637
B	273	CYS	ARG	ENGINEERED MUTATION	UNP P04637
B	284	ARG	THR	ENGINEERED MUTATION	UNP P04637
C	273	CYS	ARG	ENGINEERED MUTATION	UNP P04637
C	284	ARG	THR	ENGINEERED MUTATION	UNP P04637
D	273	CYS	ARG	ENGINEERED MUTATION	UNP P04637
D	284	ARG	THR	ENGINEERED MUTATION	UNP P04637


- Molecule 2 is a DNA chain called DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	11	Total	C	N	O	P	0	0	0
			221	105	42	64	10			
2	F	11	Total	C	N	O	P	0	0	0
			227	106	44	66	11			
2	G	12	Total	C	N	O	P	0	2	0
			284	134	55	82	13			
2	H	12	Total	C	N	O	P	0	0	0
			243	115	47	70	11			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	1
			8	4	4		
4	A	1	Total	C	O	0	1
			8	4	4		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	H	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	259	Total 263	O 263	0	8
6	B	189	Total 191	O 191	0	5
6	C	259	Total 263	O 263	0	6
6	D	179	Total 186	O 186	0	5
6	E	30	Total 30	O 30	0	1
6	F	16	Total 16	O 16	0	0
6	G	16	Total 17	O 17	0	1
6	H	11	Total 12	O 12	0	1



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cellular tumor antigen p53




- Molecule 1: Cellular tumor antigen p53




- Molecule 1: Cellular tumor antigen p53




- Molecule 1: Cellular tumor antigen p53




- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')



- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')



- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')



- Molecule 2: DNA (5'-D(*CP*GP*GP*GP*CP*AP*TP*GP*CP*CP*CP*G)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.54Å 58.16Å 77.97Å 82.96° 87.78° 73.57°	Depositor
Resolution (Å)	30.12 – 1.70 37.24 – 1.70	Depositor EDS
% Data completeness (in resolution range)	95.3 (30.12-1.70) 95.4 (37.24-1.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.70 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.163 , 0.197 0.163 , 0.197	Depositor DCC
R_{free} test set	4788 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	19.2	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 50.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8459	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/1704	0.59	0/2308
1	B	0.33	0/1652	0.55	0/2237
1	C	0.39	0/1625	0.60	0/2201
1	D	0.32	0/1619	0.54	0/2193
2	E	0.67	0/247	1.34	1/379 (0.3%)
2	F	0.65	0/254	1.27	1/390 (0.3%)
2	G	0.58	0/317	1.29	2/485 (0.4%)
2	H	0.63	0/272	1.24	1/418 (0.2%)
All	All	0.40	0/7690	0.73	5/10611 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	10	DC	O4'-C1'-N1	-8.29	102.20	108.00
2	H	6	DA	O4'-C1'-N9	-7.46	102.78	108.00
2	G	6	DA	O4'-C1'-N9	-6.52	103.44	108.00
2	E	3	DG	O4'-C1'-N9	-5.69	104.02	108.00
2	G	7	DT	N3-C4-O4	5.15	122.99	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1651	0	1614	17	0
1	B	1611	0	1559	11	0
1	C	1578	0	1525	13	0
1	D	1582	0	1517	11	0
2	E	221	0	124	3	0
2	F	227	0	123	2	0
2	G	284	0	158	5	0
2	H	243	0	135	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	28	0	42	4	0
4	B	12	0	18	2	0
4	C	12	0	18	5	0
4	D	20	0	30	6	0
5	B	4	0	3	1	0
5	H	4	0	3	1	0
6	A	263	0	0	2	0
6	B	191	0	0	3	0
6	C	263	0	0	4	0
6	D	186	0	0	2	0
6	E	30	0	0	0	0
6	F	16	0	0	0	0
6	G	17	0	0	0	0
6	H	12	0	0	0	0
All	All	8459	0	6869	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:131:ASN:HB3	4:C:304:EDO:H22	1.61	0.82
1:A:209:ARG:HB3	4:A:306:EDO:H22	1.63	0.80
1:A:206:LEU:HD11	4:A:306:EDO:H12	1.65	0.79
1:A:284:ARG:NH2	2:F:7:DT:OP2	2.18	0.77
1:C:243[B]:MET:SD	6:D:431:HOH:O	2.46	0.73
1:C:114:LEU:HD11	4:C:302:EDO:H12	1.72	0.71
1:B:114:LEU:HD11	4:B:302:EDO:H12	1.75	0.69
2:H:1:DC:H2'	2:H:2:DG:C8	2.32	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:285:GLU:OE1	6:B:422:HOH:O	2.15	0.64
1:D:183[B]:SER:O	1:D:185:SER:N	2.32	0.63
4:C:304:EDO:O2	6:C:650:HOH:O	2.13	0.63
1:B:119:ALA:O	1:B:121:SER:N	2.32	0.62
1:B:198:GLU:OE1	1:B:235:ASN:ND2	2.22	0.62
1:D:139:LYS:HA	4:D:302:EDO:H12	1.81	0.62
2:F:2:DG:H2''	2:F:3:DG:H5'	1.83	0.61
1:A:166:SER:HA	1:A:169:MET:HG3	1.84	0.60
1:B:138:ALA:HB1	4:B:303:EDO:H12	1.83	0.58
6:A:464:HOH:O	2:G:12[A]:DG:H2''	2.04	0.57
1:A:290[B]:ARG:HH21	1:A:290[B]:ARG:HG3	1.69	0.56
1:D:219:PRO:HG3	4:D:303:EDO:H12	1.88	0.56
1:D:284:ARG:NH2	2:H:7:DT:OP2	2.40	0.54
1:C:256:THR:HG22	1:C:267:ARG:HG3	1.89	0.54
2:G:1:DC:H2'	2:G:2:DG:C8	2.43	0.54
1:A:120:LYS:HD3	2:E:2:DG:H2''	1.88	0.53
1:A:250:PRO:HA	4:A:303[B]:EDO:H11	1.91	0.53
1:D:140:THR:OG1	4:D:302:EDO:H21	2.09	0.51
1:A:256:THR:HG22	1:A:267:ARG:HG3	1.93	0.51
1:D:202:ARG:HG2	4:D:306:EDO:H22	1.93	0.51
1:D:132:LYS:NZ	1:D:285:GLU:OE2	2.34	0.50
1:B:289:LEU:O	1:B:291:LYS:N	2.38	0.50
2:E:10:DC:H2'	2:E:11:DC:C6	2.47	0.49
1:B:284[A]:ARG:NH2	2:E:7:DT:OP2	2.45	0.49
2:G:11[A]:DC:H2''	2:G:12[A]:DG:C8	2.48	0.48
2:H:11:DC:H2''	2:H:12:DG:C8	2.50	0.47
1:A:290[B]:ARG:NH2	1:A:290[B]:ARG:HG3	2.30	0.45
1:C:130:LEU:HD22	4:C:303:EDO:H12	1.96	0.45
2:G:10:DC:H2''	2:G:11[A]:DC:C5	2.52	0.45
1:A:175[A]:ARG:HD3	1:A:191:PRO:O	2.17	0.45
4:D:305:EDO:H12	6:D:472:HOH:O	2.16	0.45
6:C:499:HOH:O	5:H:101:ACT:H1	2.15	0.45
1:B:185:SER:O	5:B:305:ACT:H3	2.17	0.45
1:C:119:ALA:O	1:C:279:GLY:HA3	2.18	0.44
1:A:175[B]:ARG:HD3	1:A:237:MET:HB2	1.99	0.43
1:B:248[A]:ARG:NH1	6:B:495:HOH:O	2.50	0.43
1:C:120:LYS:HG3	1:C:280:ARG:HB2	1.99	0.43
1:A:103:TYR:OH	1:A:264:LEU:HB3	2.19	0.43
1:D:192[B]:GLN:HG3	1:D:214:HIS:CE1	2.53	0.43
1:A:131:ASN:HB3	4:A:304[B]:EDO:H11	1.99	0.43
1:D:175:ARG:HD3	1:D:191:PRO:O	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:165:GLN:O	1:D:169[A]:MET:HG3	2.18	0.42
1:A:107:TYR:CZ	1:A:152[B]:PRO:HD3	2.55	0.42
1:C:177:PRO:HA	1:C:180:GLU:HG2	2.01	0.42
1:C:142:PRO:HB3	4:C:302:EDO:H21	2.01	0.42
1:B:290[B]:ARG:HG3	6:B:583:HOH:O	2.20	0.42
2:G:11[B]:DC:H42	2:H:2:DG:H1	1.68	0.42
1:A:119:ALA:O	1:A:279:GLY:HA3	2.20	0.41
1:A:110:ARG:HA	1:A:268[B]:ASN:ND2	2.35	0.41
1:C:284:ARG:NH1	6:C:494:HOH:O	2.51	0.41
1:C:178:HIS:ND1	6:C:466:HOH:O	2.37	0.41
1:C:124:CYS:SG	1:C:133[B]:MET:SD	3.18	0.41
1:D:231:THR:HG21	4:D:305:EDO:H21	2.03	0.41
1:A:153[A]:PRO:HA	6:A:644:HOH:O	2.20	0.40
1:B:101:LYS:O	1:B:267:ARG:HD2	2.21	0.40
1:C:221:GLU:HA	1:C:222:PRO:HD3	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/200 (106%)	210 (100%)	1 (0%)	0	100	100
1	B	205/200 (102%)	200 (98%)	2 (1%)	3 (2%)	10	2
1	C	202/200 (101%)	200 (99%)	2 (1%)	0	100	100
1	D	203/200 (102%)	198 (98%)	1 (0%)	4 (2%)	7	1
All	All	821/800 (103%)	808 (98%)	6 (1%)	7 (1%)	29	5

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	120	LYS
1	D	183[A]	SER
1	D	183[B]	SER
1	D	184[A]	ASP
1	D	184[B]	ASP
1	B	290[A]	ARG
1	B	290[B]	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/179 (106%)	184 (97%)	6 (3%)	39	20
1	B	184/179 (103%)	182 (99%)	2 (1%)	73	63
1	C	179/179 (100%)	178 (99%)	1 (1%)	86	80
1	D	179/179 (100%)	175 (98%)	4 (2%)	52	34
All	All	732/716 (102%)	719 (98%)	13 (2%)	62	43

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100[A]	GLN
1	A	100[B]	GLN
1	A	110	ARG
1	A	169	MET
1	A	183	SER
1	A	291	LYS
1	B	110	ARG
1	B	289	LEU
1	C	110	ARG
1	D	192[A]	GLN
1	D	192[B]	GLN
1	D	207	ASP
1	D	289	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no

such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 24 ligands modelled in this entry, 4 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	C	303	-	3,3,3	0.42	0	2,2,2	0.48	0
4	EDO	A	303[B]	-	3,3,3	0.45	0	2,2,2	0.29	0
5	ACT	B	305	-	1,3,3	1.53	0	0,3,3	0.00	-
4	EDO	D	306	-	3,3,3	0.42	0	2,2,2	0.39	0
4	EDO	D	305	-	3,3,3	0.42	0	2,2,2	0.37	0
4	EDO	B	302	-	3,3,3	0.44	0	2,2,2	0.23	0
4	EDO	A	302	-	3,3,3	0.45	0	2,2,2	0.13	0
4	EDO	D	302	-	3,3,3	0.51	0	2,2,2	0.12	0
4	EDO	A	303[A]	-	3,3,3	0.47	0	2,2,2	0.20	0
4	EDO	A	306	-	3,3,3	0.50	0	2,2,2	0.13	0
5	ACT	H	101	-	1,3,3	0.98	0	0,3,3	0.00	-
4	EDO	A	304[A]	-	3,3,3	0.47	0	2,2,2	0.34	0
4	EDO	C	302	-	3,3,3	0.39	0	2,2,2	0.39	0
4	EDO	A	305	-	3,3,3	0.43	0	2,2,2	0.31	0
4	EDO	D	303	-	3,3,3	0.42	0	2,2,2	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	D	304	-	3,3,3	0.47	0	2,2,2	0.28	0
4	EDO	B	304	-	3,3,3	0.45	0	2,2,2	0.12	0
4	EDO	A	304[B]	-	3,3,3	0.49	0	2,2,2	0.10	0
4	EDO	C	304	-	3,3,3	0.45	0	2,2,2	0.46	0
4	EDO	B	303	-	3,3,3	0.47	0	2,2,2	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	303	-	-	0/1/1/1	-
4	EDO	A	303[B]	-	-	0/1/1/1	-
4	EDO	C	302	-	-	0/1/1/1	-
4	EDO	D	306	-	-	0/1/1/1	-
4	EDO	D	305	-	-	0/1/1/1	-
4	EDO	B	302	-	-	0/1/1/1	-
4	EDO	A	302	-	-	0/1/1/1	-
4	EDO	D	302	-	-	1/1/1/1	-
4	EDO	A	303[A]	-	-	1/1/1/1	-
4	EDO	A	306	-	-	1/1/1/1	-
4	EDO	A	304[A]	-	-	0/1/1/1	-
4	EDO	A	305	-	-	0/1/1/1	-
4	EDO	D	303	-	-	1/1/1/1	-
4	EDO	D	304	-	-	1/1/1/1	-
4	EDO	B	304	-	-	0/1/1/1	-
4	EDO	A	304[B]	-	-	0/1/1/1	-
4	EDO	C	304	-	-	1/1/1/1	-
4	EDO	B	303	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	302	EDO	O1-C1-C2-O2
4	A	306	EDO	O1-C1-C2-O2
4	D	303	EDO	O1-C1-C2-O2
4	C	304	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	D	304	EDO	O1-C1-C2-O2
4	A	303[A]	EDO	O1-C1-C2-O2

There are no ring outliers.

14 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	303	EDO	1	0
4	A	303[B]	EDO	1	0
5	B	305	ACT	1	0
4	D	306	EDO	1	0
4	D	305	EDO	2	0
4	B	302	EDO	1	0
4	D	302	EDO	2	0
4	A	306	EDO	2	0
5	H	101	ACT	1	0
4	C	302	EDO	2	0
4	D	303	EDO	1	0
4	A	304[B]	EDO	1	0
4	C	304	EDO	2	0
4	B	303	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/200 (98%)	-0.17	4 (2%) 65 69	9, 18, 35, 65	2 (1%)
1	B	198/200 (99%)	0.12	8 (4%) 38 42	12, 25, 49, 82	0
1	C	195/200 (97%)	-0.09	5 (2%) 56 60	7, 19, 38, 68	2 (1%)
1	D	196/200 (98%)	0.09	5 (2%) 56 60	13, 28, 47, 72	1 (0%)
2	E	11/12 (91%)	0.33	1 (9%) 9 10	22, 25, 65, 75	1 (9%)
2	F	11/12 (91%)	0.68	3 (27%) 0 0	21, 31, 61, 71	0
2	G	12/12 (100%)	0.99	4 (33%) 0 0	20, 31, 47, 58	0
2	H	12/12 (100%)	0.59	1 (8%) 11 13	23, 33, 69, 72	0
All	All	832/848 (98%)	0.02	31 (3%) 41 46	7, 23, 47, 82	6 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	289	LEU	11.8
1	D	289	LEU	5.7
1	B	290[A]	ARG	5.5
2	E	11	DC	4.7
1	B	291	LYS	4.3
1	C	289	LEU	4.3
1	C	225	VAL	4.0
1	B	225	VAL	3.8
2	F	3	DG	3.7
2	H	12	DG	3.7
1	A	96	SER	3.6
1	B	212	PHE	3.3
2	F	2	DG	3.2
1	D	290	ARG	3.1
2	G	3	DG	3.0
2	G	2	DG	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	288	ASN	2.8
1	C	153	PRO	2.8
2	F	4	DG	2.6
1	A	292	LYS	2.5
1	D	129	ALA	2.5
1	C	290	ARG	2.4
1	B	118	THR	2.4
1	B	224	GLU	2.4
2	G	4	DG	2.4
1	B	153	PRO	2.3
2	G	1	DC	2.3
1	C	226	GLY	2.1
1	D	228	ASP	2.1
1	A	291	LYS	2.1
1	A	225	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	D	304	4/4	0.69	0.17	42,43,44,52	4
4	EDO	C	304	4/4	0.74	0.28	18,25,27,28	4
4	EDO	A	303[A]	4/4	0.79	0.28	29,33,36,40	4
4	EDO	C	303	4/4	0.79	0.23	38,41,41,42	4
4	EDO	A	303[B]	4/4	0.79	0.28	35,36,38,40	4
4	EDO	B	304	4/4	0.80	0.24	27,27,28,40	4
4	EDO	A	304[B]	4/4	0.81	0.20	33,34,35,36	4
4	EDO	A	304[A]	4/4	0.81	0.20	32,34,35,36	4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	B	302	4/4	0.82	0.23	27,31,32,37	4
5	ACT	B	305	4/4	0.82	0.14	43,44,52,56	0
4	EDO	A	305	4/4	0.85	0.14	21,25,25,29	4
4	EDO	A	306	4/4	0.87	0.21	20,29,30,31	4
4	EDO	D	306	4/4	0.87	0.14	42,42,43,49	4
4	EDO	D	302	4/4	0.89	0.30	23,27,39,40	4
4	EDO	D	305	4/4	0.90	0.28	33,34,38,42	4
4	EDO	A	302	4/4	0.91	0.16	23,28,29,31	4
5	ACT	H	101	4/4	0.92	0.11	25,31,32,44	0
4	EDO	C	302	4/4	0.93	0.15	24,29,30,33	4
4	EDO	D	303	4/4	0.93	0.16	24,35,42,48	4
4	EDO	B	303	4/4	0.94	0.17	27,30,30,30	4
3	ZN	A	301	1/1	0.99	0.06	16,16,16,16	0
3	ZN	D	301	1/1	0.99	0.06	19,19,19,19	0
3	ZN	C	301	1/1	0.99	0.06	17,17,17,17	0
3	ZN	B	301	1/1	1.00	0.05	19,19,19,19	0

6.5 Other polymers ⓘ

There are no such residues in this entry.