



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 03:56 am BST

PDB ID : 6ICN
Title : Pseudomonas putida CBB5 NdmA with caffeine
Authors : Kim, J.H.; Kim, B.H.; Kang, S.Y.; Song, H.K.
Deposited on : 2018-09-06
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	<div> <div>7%</div> <div> <div></div> <div>86%</div> <div>8%</div> <div>7%</div> </div> </div>
1	B	369	<div> <div>6%</div> <div> <div></div> <div>88%</div> <div>5%</div> <div>7%</div> </div> </div>
1	C	369	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>7%</div> <div>7%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methylxanthine N1-demethylase NdmA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	344	Total	C	N	O	S	0	0	0
			2782	1771	476	522	13			
1	B	344	Total	C	N	O	S	0	0	0
			2782	1771	476	522	13			
1	C	344	Total	C	N	O	S	0	0	0
			2782	1771	476	522	13			

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	expression tag	UNP H9N289
A	-16	GLY	-	expression tag	UNP H9N289
A	-15	SER	-	expression tag	UNP H9N289
A	-14	SER	-	expression tag	UNP H9N289
A	-13	HIS	-	expression tag	UNP H9N289
A	-12	HIS	-	expression tag	UNP H9N289
A	-11	HIS	-	expression tag	UNP H9N289
A	-10	HIS	-	expression tag	UNP H9N289
A	-9	HIS	-	expression tag	UNP H9N289
A	-8	HIS	-	expression tag	UNP H9N289
A	-7	GLU	-	expression tag	UNP H9N289
A	-6	ASN	-	expression tag	UNP H9N289
A	-5	LEU	-	expression tag	UNP H9N289
A	-4	TYR	-	expression tag	UNP H9N289
A	-3	PHE	-	expression tag	UNP H9N289
A	-2	GLN	-	expression tag	UNP H9N289
A	-1	GLY	-	expression tag	UNP H9N289
A	0	SER	-	expression tag	UNP H9N289
B	-17	MET	-	expression tag	UNP H9N289
B	-16	GLY	-	expression tag	UNP H9N289
B	-15	SER	-	expression tag	UNP H9N289
B	-14	SER	-	expression tag	UNP H9N289
B	-13	HIS	-	expression tag	UNP H9N289

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	HIS	-	expression tag	UNP H9N289
B	-11	HIS	-	expression tag	UNP H9N289
B	-10	HIS	-	expression tag	UNP H9N289
B	-9	HIS	-	expression tag	UNP H9N289
B	-8	HIS	-	expression tag	UNP H9N289
B	-7	GLU	-	expression tag	UNP H9N289
B	-6	ASN	-	expression tag	UNP H9N289
B	-5	LEU	-	expression tag	UNP H9N289
B	-4	TYR	-	expression tag	UNP H9N289
B	-3	PHE	-	expression tag	UNP H9N289
B	-2	GLN	-	expression tag	UNP H9N289
B	-1	GLY	-	expression tag	UNP H9N289
B	0	SER	-	expression tag	UNP H9N289
C	-17	MET	-	expression tag	UNP H9N289
C	-16	GLY	-	expression tag	UNP H9N289
C	-15	SER	-	expression tag	UNP H9N289
C	-14	SER	-	expression tag	UNP H9N289
C	-13	HIS	-	expression tag	UNP H9N289
C	-12	HIS	-	expression tag	UNP H9N289
C	-11	HIS	-	expression tag	UNP H9N289
C	-10	HIS	-	expression tag	UNP H9N289
C	-9	HIS	-	expression tag	UNP H9N289
C	-8	HIS	-	expression tag	UNP H9N289
C	-7	GLU	-	expression tag	UNP H9N289
C	-6	ASN	-	expression tag	UNP H9N289
C	-5	LEU	-	expression tag	UNP H9N289
C	-4	TYR	-	expression tag	UNP H9N289
C	-3	PHE	-	expression tag	UNP H9N289
C	-2	GLN	-	expression tag	UNP H9N289
C	-1	GLY	-	expression tag	UNP H9N289
C	0	SER	-	expression tag	UNP H9N289

- Molecule 2 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			4	2	2		
2	B	1	Total	Fe	S	0	0
			4	2	2		
2	C	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 3 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Co	0	0
			1	1		
3	A	1	Total	Co	0	0
			1	1		
3	C	1	Total	Co	0	0
			1	1		

- Molecule 4 is CAFFEINE (three-letter code: CFF) (formula: C₈H₁₀N₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	4	2		
4	B	1	Total	C	N	O	0	0
			14	8	4	2		
4	C	1	Total	C	N	O	0	0
			14	8	4	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	471	Total	O	0	0
			471	471		
5	B	490	Total	O	0	0
			490	490		
5	C	461	Total	O	0	0
			461	461		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methylxanthine N1-demethylase NdmA



- Molecule 1: Methylxanthine N1-demethylase NdmA



- Molecule 1: Methylxanthine N1-demethylase NdmA



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	135.94Å 135.94Å 155.43Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.76 – 1.65 42.78 – 1.65	Depositor EDS
% Data completeness (in resolution range)	96.8 (33.76-1.65) 96.4 (42.78-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.25 (at 1.65Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.173 , 0.204 0.173 , 0.204	Depositor DCC
R_{free} test set	2006 reflections (1.01%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.012 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9825	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CFF, CO, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2867	0.57	0/3907
1	B	0.41	0/2867	0.57	0/3907
1	C	0.42	0/2867	0.59	1/3907 (0.0%)
All	All	0.42	0/8601	0.58	1/11721 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	113	ARG	NE-CZ-NH1	6.05	123.33	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	180	GLY	Peptide
1	B	180	GLY	Peptide
1	C	180	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2782	0	2640	19	0
1	B	2782	0	2640	15	0
1	C	2782	0	2640	17	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	14	0	10	1	0
4	B	14	0	10	1	0
4	C	14	0	10	1	0
5	A	471	0	0	9	2
5	B	490	0	0	9	2
5	C	461	0	0	6	0
All	All	9825	0	7950	50	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:322:ARG:NH1	5:B:501:HOH:O	1.91	1.02
1:A:348:ARG:NH1	5:A:501:HOH:O	1.93	1.00
1:A:322:ARG:NH2	5:A:502:HOH:O	2.03	0.88
1:B:336:ARG:NH1	5:B:502:HOH:O	2.05	0.88
1:A:196:ASP:OD2	1:A:348:ARG:NH2	2.17	0.77
1:B:336:ARG:NH2	5:B:504:HOH:O	2.23	0.62
1:C:184:ASP:HB3	1:C:186:ASN:ND2	2.16	0.61
1:B:80:GLN:NE2	5:B:510:HOH:O	2.35	0.59
1:B:7:ASN:N	5:B:511:HOH:O	2.36	0.57
1:C:14:ARG:NH1	1:C:44:GLU:OE1	2.30	0.56
1:C:10:ARG:NH1	1:C:162:GLU:OE2	2.37	0.56
1:C:12:TYR:CE1	1:C:325:LYS:HG2	2.42	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:TRP:CZ3	1:A:293:ILE:HG21	2.41	0.55
1:A:333:GLN:H	1:A:333:GLN:CD	2.09	0.55
1:B:10:ARG:NH2	5:B:508:HOH:O	2.33	0.55
1:C:189:GLU:OE2	5:C:502:HOH:O	2.18	0.55
1:B:298:PRO:HG2	1:B:302:PRO:HG3	1.90	0.54
1:C:322:ARG:NH2	5:C:512:HOH:O	2.43	0.52
1:C:102:ASN:OD1	5:C:504:HOH:O	2.20	0.50
1:B:157:TRP:CZ3	1:B:293:ILE:HG21	2.48	0.49
1:C:184:ASP:OD2	1:C:212:MET:HA	2.13	0.49
1:C:11:GLU:HG2	5:C:600:HOH:O	2.14	0.48
1:A:298:PRO:HG2	1:A:302:PRO:HG3	1.97	0.47
1:C:117:GLU:OE2	5:C:503:HOH:O	2.19	0.47
1:C:157:TRP:CZ3	1:C:293:ILE:HG21	2.50	0.47
1:B:282:ASN:OD1	4:B:402:CFF:H81	2.15	0.47
1:B:242:SER:HB3	5:B:506:HOH:O	2.16	0.46
1:A:282:ASN:OD1	4:A:402:CFF:H81	2.15	0.46
1:C:282:ASN:OD1	4:C:402:CFF:H81	2.16	0.46
1:A:157:TRP:CZ3	1:A:293:ILE:HD13	2.52	0.45
1:B:348:ARG:NH1	5:B:521:HOH:O	2.50	0.45
1:C:12:TYR:CD1	1:C:325:LYS:HG2	2.52	0.45
1:A:61:ARG:NH2	1:B:302:PRO:HG2	2.32	0.44
1:C:59:ARG:NH2	1:C:112:ASP:OD2	2.50	0.44
1:C:333:GLN:H	1:C:333:GLN:CD	2.21	0.44
1:A:81:CYS:SG	1:A:82:PRO:HD2	2.58	0.44
1:A:222:SER:OG	5:A:505:HOH:O	2.21	0.44
1:B:323:GLU:HG3	5:B:558:HOH:O	2.17	0.43
1:A:73:THR:HG22	5:A:833:HOH:O	2.17	0.43
1:A:77:ASN:ND2	5:A:519:HOH:O	2.51	0.43
1:A:43:ASN:HB2	5:A:841:HOH:O	2.17	0.43
1:B:297:TRP:HA	1:B:298:PRO:C	2.38	0.43
1:C:322:ARG:HD2	5:C:525:HOH:O	2.19	0.42
1:A:76:GLY:O	5:A:506:HOH:O	2.22	0.42
1:A:135:GLU:OE2	5:A:504:HOH:O	2.21	0.41
1:A:44:GLU:OE1	1:A:59:ARG:NH1	2.52	0.41
1:B:166:GLU:O	1:B:170:ASP:HB2	2.21	0.41
1:A:59:ARG:HD2	5:A:617:HOH:O	2.20	0.41
1:A:7:ASN:OD1	1:A:8:ASP:N	2.51	0.41
1:C:297:TRP:HA	1:C:298:PRO:C	2.41	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:579:HOH:O	5:B:798:HOH:O[4_556]	2.16	0.04
5:A:683:HOH:O	5:B:553:HOH:O[4_556]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/369 (93%)	336 (98%)	6 (2%)	0	100	100
1	B	342/369 (93%)	333 (97%)	9 (3%)	0	100	100
1	C	342/369 (93%)	332 (97%)	10 (3%)	0	100	100
All	All	1026/1107 (93%)	1001 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/328 (93%)	301 (98%)	5 (2%)	62	41
1	B	306/328 (93%)	303 (99%)	3 (1%)	76	62
1	C	306/328 (93%)	303 (99%)	3 (1%)	76	62
All	All	918/984 (93%)	907 (99%)	11 (1%)	71	53

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	TYR
1	A	113	ARG
1	A	164	ARG
1	A	202	PHE
1	A	350	TYR
1	B	83	TYR
1	B	164	ARG
1	B	202	PHE
1	C	83	TYR
1	C	164	ARG
1	C	202	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	216	ASN
1	C	186	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CFF	C	402	-	8,15,15	1.90	2 (25%)	8,23,23	1.21	1 (12%)
2	FES	B	400	1	0,4,4	0.00	-	-	-	-
4	CFF	B	402	-	8,15,15	1.87	2 (25%)	8,23,23	1.19	1 (12%)
2	FES	C	400	1	0,4,4	0.00	-	-	-	-
2	FES	A	400	1	0,4,4	0.00	-	-	-	-
4	CFF	A	402	-	8,15,15	1.76	2 (25%)	8,23,23	1.11	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CFF	C	402	-	-	-	0/2/2/2
2	FES	B	400	1	-	-	0/1/1/1
4	CFF	B	402	-	-	-	0/2/2/2
2	FES	C	400	1	-	-	0/1/1/1
2	FES	A	400	1	-	-	0/1/1/1
4	CFF	A	402	-	-	-	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	402	CFF	C5-C4	-3.52	1.34	1.39
4	B	402	CFF	C5-C4	-3.44	1.34	1.39
4	C	402	CFF	C6-N1	-3.28	1.33	1.38
4	A	402	CFF	C5-C4	-3.19	1.35	1.39
4	B	402	CFF	C6-N1	-3.13	1.33	1.38
4	A	402	CFF	C6-N1	-2.95	1.34	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	402	CFF	C14-N7-C8	-2.24	114.66	125.43
4	C	402	CFF	C14-N7-C8	-2.23	114.72	125.43
4	B	402	CFF	C14-N7-C8	-2.18	114.92	125.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	402	CFF	1	0
4	B	402	CFF	1	0
4	A	402	CFF	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	344/369 (93%)	0.10	27 (7%)	13 12	14, 22, 43, 65	0
1	B	344/369 (93%)	0.10	23 (6%)	17 16	15, 21, 41, 60	0
1	C	344/369 (93%)	0.01	19 (5%)	25 24	15, 22, 39, 59	0
All	All	1032/1107 (93%)	0.07	69 (6%)	17 16	14, 22, 42, 65	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	349	SER	7.9
1	A	350	TYR	7.7
1	A	186	ASN	5.8
1	C	350	TYR	5.8
1	C	7	ASN	5.7
1	C	186	ASN	5.3
1	B	350	TYR	5.0
1	B	7	ASN	5.0
1	A	348	ARG	4.8
1	B	349	SER	4.7
1	A	7	ASN	4.7
1	B	143	ASN	4.6
1	A	347	ASP	4.4
1	A	8	ASP	4.2
1	C	8	ASP	4.1
1	C	347	ASP	4.1
1	C	187	ASN	3.9
1	A	211	ASP	3.9
1	C	349	SER	3.7
1	A	216	ASN	3.7
1	C	143	ASN	3.7
1	B	243	SER	3.6
1	A	213	ALA	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	242	SER	3.4
1	C	213	ALA	3.3
1	C	348	ARG	3.2
1	B	102	ASN	3.2
1	A	9	GLU	3.2
1	B	242	SER	3.2
1	B	186	ASN	3.1
1	B	347	ASP	3.1
1	A	187	ASN	3.0
1	C	102	ASN	3.0
1	B	216	ASN	3.0
1	A	240	TYR	2.9
1	B	206	TYR	2.9
1	A	185	PRO	2.9
1	A	143	ASN	2.9
1	A	215	PRO	2.9
1	C	211	ASP	2.8
1	B	185	PRO	2.8
1	B	9	GLU	2.8
1	B	209	PRO	2.8
1	C	208	THR	2.6
1	C	240	TYR	2.5
1	B	211	ASP	2.5
1	B	8	ASP	2.5
1	C	210	GLU	2.5
1	A	214	VAL	2.4
1	B	208	THR	2.4
1	A	208	THR	2.4
1	A	221	GLY	2.4
1	B	187	ASN	2.3
1	B	258	HIS	2.3
1	A	210	GLU	2.3
1	C	330	GLU	2.3
1	C	9	GLU	2.3
1	A	243	SER	2.2
1	C	209	PRO	2.2
1	A	333	GLN	2.2
1	A	206	TYR	2.2
1	B	240	TYR	2.2
1	B	348	ARG	2.2
1	C	333	GLN	2.1
1	B	10	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	10	ARG	2.1
1	B	180	GLY	2.1
1	A	212	MET	2.0
1	A	102	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CFF	B	402	14/14	0.86	0.13	25,27,28,28	0
4	CFF	A	402	14/14	0.86	0.14	25,27,30,32	0
4	CFF	C	402	14/14	0.94	0.13	23,25,27,28	0
3	CO	C	401	1/1	0.99	0.07	20,20,20,20	0
2	FES	C	400	4/4	0.99	0.09	15,15,16,18	0
3	CO	B	401	1/1	0.99	0.09	20,20,20,20	0
3	CO	A	401	1/1	0.99	0.08	20,20,20,20	0
2	FES	B	400	4/4	0.99	0.09	16,16,16,17	0
2	FES	A	400	4/4	1.00	0.10	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.