



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 09:45 pm BST

PDB ID : 1IDU  
Title : CRYSTAL STRUCTURE OF THE PEROXIDE FORM OF THE  
VANADIUM-CONTAINING CHLOROPEROXIDASE FROM CURVU-  
LARIA INAEQUALIS  
Authors : Messerschmidt, A.; Prade, L.; Wever, R.  
Deposited on : 2001-04-05  
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	609	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5096 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called VANADIUM CHLOROPEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	576	4487	2856	772	846	13	58	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	185	ASN	HIS	CONFLICT	UNP P49053
A	544	GLU	PRO	SEE REMARK 999	UNP P49053

- Molecule 2 is VANADATE ION (three-letter code: VO4) (formula: O<sub>4</sub>V).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	V		
2	A	1	5	4	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	604	Total 604	O 604	3	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: VANADIUM CHLOROPEROXIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	129.95Å 129.95Å 111.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 2.24 33.84 – 2.24	Depositor EDS
% Data completeness (in resolution range)	88.3 (8.00-2.24) 88.3 (33.84-2.24)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.53 (at 2.24Å)	Xtriage
Refinement program	X-PLOR 3.843	Depositor
R, $R_{free}$	0.177 , (Not available) 0.171 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.1	Xtriage
Anisotropy	0.004	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 115.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.036 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5096	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/4620	0.75	4/6316 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	121	PRO	N-CA-CB	5.67	110.11	103.30
1	A	125	PRO	N-CA-CB	5.43	109.81	103.30
1	A	123	PRO	N-CA-CB	5.36	109.73	103.30
1	A	49	LEU	CA-CB-CG	-5.11	103.55	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4487	0	4265	74	0
2	A	5	0	0	0	0
3	A	604	0	0	12	0
All	All	5096	0	4265	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:576:GLN:HA	1:A:576:GLN:HE21	1.49	0.78
1:A:232:ARG:H	1:A:576:GLN:HE22	1.31	0.77
1:A:394:LYS:HE3	1:A:495:VAL:HG12	1.71	0.73
1:A:28:HIS:HE1	3:A:2144:HOH:O	1.73	0.72
1:A:449:ARG:HD3	1:A:461:ILE:HG13	1.72	0.70
1:A:519:VAL:CG1	1:A:523:GLY:HA2	2.27	0.65
1:A:519:VAL:HG13	1:A:523:GLY:HA2	1.80	0.64
1:A:116:LYS:HG2	1:A:134:TYR:CG	2.33	0.63
1:A:82:THR:HG23	1:A:88:ARG:HA	1.80	0.63
1:A:79:SER:O	1:A:82:THR:HG22	1.99	0.62
1:A:130:SER:HB2	3:A:1869:HOH:O	2.02	0.60
1:A:17:GLU:HB2	3:A:1799:HOH:O	2.01	0.60
1:A:63:ASP:O	1:A:67:SER:HB2	2.01	0.60
1:A:549:PRO:HB2	1:A:554:PRO:HB2	1.86	0.58
1:A:11:LYS:HE2	3:A:1868:HOH:O	2.04	0.57
1:A:241:LEU:HD21	1:A:555:LEU:HB2	1.87	0.57
1:A:4:VAL:HG12	1:A:132:ASN:ND2	2.21	0.55
1:A:74:PHE:HB2	3:A:1937:HOH:O	2.06	0.54
1:A:82:THR:CG2	1:A:88:ARG:HA	2.37	0.54
1:A:205:PRO:HA	1:A:215:LYS:HA	1.90	0.54
1:A:572:PRO:HG2	1:A:575:ILE:HG12	1.90	0.53
1:A:378:TRP:CZ3	1:A:380:THR:HG22	2.44	0.53
1:A:205:PRO:HA	1:A:214:PRO:O	2.09	0.53
1:A:47:PRO:HD2	1:A:48:PRO:HD2	1.91	0.52
1:A:305:ASN:O	1:A:309:ARG:HG3	2.09	0.51
1:A:438:MET:HA	1:A:473:VAL:CG1	2.40	0.51
1:A:297:ILE:HD11	1:A:448:ASN:ND2	2.26	0.51
1:A:82:THR:HG21	1:A:87:TYR:O	2.12	0.50
1:A:181:GLN:HG3	1:A:221:TYR:HB2	1.93	0.50
1:A:69:CYS:SG	3:A:2116:HOH:O	2.60	0.49
1:A:300:PRO:HB2	1:A:301:PRO:HD3	1.95	0.49
1:A:116:LYS:HG2	1:A:134:TYR:CD1	2.47	0.49
1:A:538:ARG:HG3	3:A:2170:HOH:O	2.14	0.47
1:A:248:ARG:HB3	1:A:542:GLU:HB2	1.96	0.47
1:A:66:PHE:HB3	1:A:71:PRO:HD3	1.96	0.47
1:A:139:LEU:O	1:A:143:ARG:HB2	2.14	0.47
1:A:190:ARG:HG2	1:A:191:TYR:CD1	2.50	0.46
1:A:390:ASP:HB3	3:A:1680:HOH:O	2.15	0.46
1:A:576:GLN:NE2	1:A:576:GLN:HA	2.24	0.46
1:A:219:ARG:HH11	1:A:219:ARG:HG2	1.80	0.45

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:572:PRO:HA	1:A:573:PRO:HD2	1.84	0.45
1:A:36:VAL:O	1:A:40:VAL:HG22	2.17	0.45
1:A:25:PHE:O	1:A:28:HIS:HB3	2.17	0.45
1:A:17:GLU:HG2	1:A:321:LEU:HD12	1.98	0.45
1:A:456:ASP:HA	1:A:457:PRO:HD2	1.87	0.44
1:A:42:GLY:HA2	1:A:129:ILE:HG23	2.00	0.44
1:A:191:TYR:HA	1:A:359:TRP:CE2	2.52	0.44
1:A:395:PRO:HB3	1:A:397:PHE:CZ	2.52	0.44
1:A:82:THR:HG21	3:A:1965:HOH:O	2.17	0.44
1:A:190:ARG:HG2	1:A:191:TYR:CE1	2.51	0.44
1:A:49:LEU:HD13	1:A:179:ALA:HB2	1.99	0.44
1:A:484:PHE:CD2	1:A:530:VAL:HG22	2.53	0.43
1:A:302:ARG:O	1:A:306:GLN:HG3	2.16	0.43
1:A:104:GLY:O	1:A:108:LYS:HB2	2.19	0.43
1:A:437:ASP:HA	1:A:475:HIS:HA	2.01	0.43
1:A:378:TRP:CH2	1:A:380:THR:HG22	2.54	0.42
1:A:535:TYR:CZ	1:A:553:VAL:HG11	2.54	0.42
1:A:207:ASP:HB3	1:A:210:ASN:HB3	2.00	0.42
1:A:479:ALA:HA	1:A:482:LEU:HD12	2.02	0.42
1:A:538:ARG:HB2	1:A:547:LEU:HB3	2.00	0.42
1:A:218:PHE:CE2	1:A:220:GLN:HB2	2.55	0.42
1:A:573:PRO:HA	1:A:576:GLN:HG2	2.02	0.41
1:A:400:TYR:HA	1:A:401:PRO:HA	1.88	0.41
1:A:519:VAL:HG13	1:A:523:GLY:CA	2.46	0.41
1:A:553:VAL:HB	1:A:554:PRO:HD3	2.01	0.41
1:A:241:LEU:HD11	1:A:401:PRO:HG3	2.03	0.41
1:A:307:ILE:O	1:A:311:ILE:HG13	2.20	0.41
1:A:394:LYS:HE2	1:A:394:LYS:HB3	1.90	0.41
1:A:568:LEU:HA	1:A:568:LEU:HD12	1.89	0.41
1:A:405:ALA:HB1	1:A:553:VAL:HG23	2.03	0.41
1:A:93:ASN:ND2	3:A:2035:HOH:O	2.53	0.41
1:A:46:GLY:HA2	3:A:2001:HOH:O	2.20	0.40
1:A:521:ASN:ND2	3:A:1903:HOH:O	2.48	0.40
1:A:299:THR:HB	1:A:300:PRO:HD2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	574/609 (94%)	545 (95%)	23 (4%)	6 (1%)	15 11

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	GLN
1	A	124	ASN
1	A	128	ASN
1	A	290	ALA
1	A	122	ASN
1	A	434	ILE

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	467/506 (92%)	445 (95%)	22 (5%)	26 27

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	VAL
1	A	49	LEU
1	A	59	LEU
1	A	67	SER
1	A	79	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	82	THR
1	A	96	ASN
1	A	108	LYS
1	A	192	LYS
1	A	196	GLU
1	A	235	THR
1	A	301	PRO
1	A	323	ASN
1	A	350	TRP
1	A	351	LYS
1	A	359	TRP
1	A	437	ASP
1	A	449	ARG
1	A	519	VAL
1	A	521	ASN
1	A	569	LYS
1	A	576	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	132	ASN
1	A	174	ASN
1	A	220	GLN
1	A	576	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	VO4	A	1579	1	1,4,4	5.18	1 (100%)	-		

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1579	VO4	O1-V	5.18	1.93	1.63

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	566/609 (92%)	-0.24	14 (2%) 57 58	8, 18, 40, 62	2 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	207	ASP	5.0
1	A	212	ASN	4.1
1	A	117	PRO	4.0
1	A	214	PRO	3.9
1	A	211	PRO	3.6
1	A	213	GLY	3.5
1	A	210	ASN	3.5
1	A	209	ASN	3.0
1	A	512	THR	2.9
1	A	93	ASN	2.8
1	A	82	THR	2.6
1	A	155	GLU	2.3
1	A	208	PRO	2.1
1	A	521	ASN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	VO4	A	1579	5/5	0.99	0.11	11,13,16,18	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.