



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 05:05 am BST

PDB ID : 4IFR  
Title : 2.40 Angstroms X-ray crystal structure of R239A 2-amino-3-carboxymuconate-6-semialdehyde decarboxylase from *Pseudomonas fluorescens*  
Authors : Huo, L.; Davis, I.; Chen, L.; Liu, A.  
Deposited on : 2012-12-14  
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	<div> <div>6%</div> <div>84%</div> <div>15%</div> <div>.</div> </div>
1	B	330	<div> <div>15%</div> <div>72%</div> <div>24%</div> <div>..</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5172 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-amino-3-carboxymuconate 6-semialdehyde decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	330	Total	C	N	O	S	1	0	0
			2572	1641	445	468	18			
1	B	330	Total	C	N	O	S	1	0	0
			2572	1641	445	468	18			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	239	ALA	ARG	engineered mutation	UNP Q83V25
B	239	ALA	ARG	engineered mutation	UNP Q83V25

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	15	Total	O	0	0
			15	15		
3	B	11	Total	O	0	0
			11	11		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2-amino-3-carboxymuconate 6-semialdehyde decarboxylase



- Molecule 1: 2-amino-3-carboxymuconate 6-semialdehyde decarboxylase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.16 Å 91.16 Å 167.99 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.52 – 2.39 31.52 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.4 (31.52-2.39) 95.5 (31.52-2.39)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 2.39 Å)	Xtriage
Refinement program	PHENIX 1.7.3 _928	Depositor
R, $R_{free}$	0.210 , 0.295 0.206 , 0.289	Depositor DCC
$R_{free}$ test set	1454 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	65.9	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 52.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5172	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/2639	0.60	0/3577
1	B	0.44	0/2639	0.61	0/3577
All	All	0.44	0/5278	0.60	0/7154

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2572	0	2512	25	0
1	B	2572	0	2512	49	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	15	0	0	1	0
3	B	11	0	0	0	0
All	All	5172	0	5024	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:156:ASP:CB	1:B:160:LEU:HD23	2.02	0.90
1:B:87:THR:HG23	1:B:151:GLY:HA3	1.63	0.81
1:A:271:VAL:O	3:A:506:HOH:O	2.03	0.77
1:B:156:ASP:HB2	1:B:160:LEU:HD23	1.66	0.75
1:B:124:GLN:HA	1:B:150:LEU:HD11	1.74	0.70
1:B:127:ASP:N	1:B:127:ASP:OD1	2.25	0.67
1:B:135:ARG:HH21	1:B:136:ALA:HA	1.60	0.66
1:B:156:ASP:HB3	1:B:160:LEU:HD23	1.77	0.65
1:B:73:ASP:HA	1:B:114:ARG:HH21	1.61	0.65
1:B:97:ALA:O	1:B:101:ASN:ND2	2.35	0.60
1:A:168:ALA:HB2	1:A:222:LEU:HB2	1.82	0.60
1:B:40:THR:OG1	1:B:41:GLY:N	2.38	0.57
1:B:178:PRO:HB3	1:B:202:THR:OG1	2.06	0.56
1:A:124:GLN:HA	1:A:150:LEU:HD21	1.90	0.54
1:B:122:PRO:HD2	1:B:129:ALA:HA	1.90	0.54
1:B:119:ALA:HB3	1:B:144:ILE:HD13	1.90	0.53
1:A:4:PRO:O	1:A:323:SER:HB2	2.08	0.53
1:B:218:ILE:HG21	1:B:222:LEU:HD22	1.90	0.53
1:A:119:ALA:HB3	1:A:144:ILE:HD13	1.90	0.52
1:B:178:PRO:HB2	1:B:198:MET:HG2	1.91	0.52
1:B:244:TRP:HA	1:B:250:VAL:HB	1.91	0.52
1:B:311:SER:O	1:B:318:LYS:NZ	2.31	0.52
1:A:309:LEU:O	1:A:318:LYS:NZ	2.42	0.51
1:B:27:ASN:O	1:B:28:HIS:ND1	2.43	0.51
1:B:135:ARG:HE	1:B:135:ARG:C	2.14	0.51
1:A:74:VAL:HG22	1:A:114:ARG:HA	1.92	0.51
1:A:148:ASN:OD1	1:A:178:PRO:HA	2.13	0.49
1:B:8:MET:HG3	1:B:291:LEU:O	2.12	0.49
1:B:112:PRO:O	1:B:116:LYS:NZ	2.31	0.48
1:B:198:MET:HB3	1:B:199:PRO:HD3	1.94	0.48
1:A:187:MET:HE2	1:A:193:PRO:HA	1.96	0.48
1:B:127:ASP:O	1:B:131:LYS:HD2	2.13	0.48
1:A:212:SER:HA	1:B:190:TRP:CZ2	2.48	0.47
1:A:286:GLU:HG3	1:A:320:LYS:HD2	1.95	0.47
1:A:181:MET:HG3	1:A:201:GLU:HG3	1.97	0.47
1:A:203:GLN:HG3	1:A:233:PHE:HA	1.96	0.47
1:A:187:MET:HE1	1:A:197:ALA:H	1.80	0.46
1:B:5:ARG:HG3	1:B:322:ILE:HG23	1.97	0.46
1:A:14:PRO:HG2	1:A:16:ILE:HD11	1.98	0.46
1:B:86:TYR:HB3	1:B:150:LEU:HA	1.98	0.46
1:B:157:ASP:HB2	1:B:158:ALA:H	1.63	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:203:GLN:HG3	1:B:233:PHE:HA	1.97	0.45
1:A:188:LYS:HE2	1:A:188:LYS:HB3	1.72	0.45
1:B:37:LYS:O	1:B:39:ASP:N	2.46	0.45
1:B:311:SER:C	1:B:313:LEU:H	2.20	0.45
1:A:63:ARG:HD3	1:A:75:GLN:OE1	2.17	0.45
1:B:134:SER:OG	1:B:170:GLU:OE1	2.32	0.44
1:B:39:ASP:O	1:B:55:GLN:HB2	2.17	0.44
1:B:16:ILE:HG12	1:B:103:PHE:CE1	2.53	0.44
1:B:37:LYS:C	1:B:39:ASP:H	2.20	0.44
1:B:80:THR:HG22	1:B:82:VAL:HG23	1.99	0.44
1:B:55:GLN:HA	1:B:58:TRP:CH2	2.53	0.43
1:A:192:LEU:HD13	1:B:204:LEU:HD11	2.00	0.43
1:B:168:ALA:HB2	1:B:222:LEU:HD13	2.01	0.43
1:A:236:LEU:HD11	1:B:232:SER:HB3	2.00	0.43
1:B:110:HIS:O	1:B:111:ASN:HB2	2.19	0.43
1:B:181:MET:O	1:B:182:MET:HB2	2.19	0.43
1:B:42:SER:OG	1:B:49:ASN:OD1	2.24	0.43
1:B:25:ASP:HB3	1:B:29:ALA:H	1.84	0.42
1:B:155:LEU:HA	1:B:155:LEU:HD23	1.77	0.42
1:A:37:LYS:HD3	1:A:39:ASP:OD1	2.20	0.42
1:A:8:MET:HA	1:A:76:VAL:HB	2.01	0.42
1:A:78:CYS:HB3	1:A:118:LEU:HB2	2.02	0.41
1:B:96:TRP:O	1:B:100:MET:HG2	2.20	0.41
1:B:161:GLU:HG3	1:B:218:ILE:HD11	2.02	0.41
1:B:263:ASP:OD2	1:B:264:ARG:HG2	2.20	0.41
1:A:291:LEU:HD22	1:A:322:ILE:HD12	2.01	0.41
1:A:5:ARG:HD3	1:A:71:GLY:O	2.21	0.41
1:B:88:TRP:HB2	1:B:93:ALA:HB2	2.02	0.41
1:B:109:ALA:HA	1:B:112:PRO:HG3	2.02	0.41
1:A:45:MET:HE3	1:A:45:MET:HB3	1.95	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/330 (99%)	313 (95%)	14 (4%)	1 (0%)	41	55
1	B	328/330 (99%)	294 (90%)	26 (8%)	8 (2%)	6	6
All	All	656/660 (99%)	607 (92%)	40 (6%)	9 (1%)	11	15

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	182	MET
1	A	252	GLU
1	B	40	THR
1	B	270	ALA
1	B	37	LYS
1	B	150	LEU
1	B	59	ASP
1	B	286	GLU
1	B	38	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/269 (100%)	257 (96%)	12 (4%)	27	44
1	B	269/269 (100%)	241 (90%)	28 (10%)	7	10
All	All	538/538 (100%)	498 (93%)	40 (7%)	13	22

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ARG
1	A	16	ILE
1	A	51	ARG
1	A	171	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	179	TRP
1	A	185	GLN
1	A	186	ARG
1	A	188	LYS
1	A	223	LYS
1	A	299	LEU
1	A	301	GLU
1	A	315	GLU
1	B	27	ASN
1	B	35	SER
1	B	40	THR
1	B	48	ASN
1	B	51	ARG
1	B	87	THR
1	B	92	LYS
1	B	103	PHE
1	B	117	VAL
1	B	127	ASP
1	B	131	LYS
1	B	135	ARG
1	B	148	ASN
1	B	150	LEU
1	B	157	ASP
1	B	160	LEU
1	B	171	ASP
1	B	179	TRP
1	B	181	MET
1	B	185	GLN
1	B	198	MET
1	B	217	ARG
1	B	245	ARG
1	B	251	ARG
1	B	256	ARG
1	B	282	SER
1	B	286	GLU
1	B	323	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	120	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	330/330 (100%)	0.32	19 (5%)	23 22	50, 70, 93, 107	1 (0%)
1	B	330/330 (100%)	0.89	50 (15%)	2 1	50, 82, 119, 140	0
All	All	660/660 (100%)	0.61	69 (10%)	6 5	50, 75, 111, 140	1 (0%)

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	150	LEU	21.6
1	B	151	GLY	21.1
1	B	152	ASP	13.7
1	B	175	LEU	4.4
1	B	333	ASN	4.2
1	B	78	CYS	4.2
1	B	313	LEU	4.1
1	B	177	HIS	3.9
1	B	34	VAL	3.8
1	B	176	VAL	3.7
1	B	145	GLN	3.7
1	B	227	GLY	3.6
1	B	118	LEU	3.5
1	B	84	PHE	3.5
1	A	78	CYS	3.4
1	B	178	PRO	3.4
1	B	26	ALA	3.4
1	B	47	LYS	3.3
1	B	183	GLY	3.3
1	B	206	ILE	3.2
1	B	15	ARG	3.2
1	B	21	ALA	3.1
1	B	83	MET	3.0
1	A	333	ASN	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	36	ALA	2.9
1	B	79	ALA	2.9
1	B	54	TYR	2.9
1	B	80	THR	2.9
1	B	92	LYS	2.8
1	B	185	GLN	2.7
1	B	10	SER	2.7
1	B	148	ASN	2.7
1	A	145	GLN	2.7
1	A	79	ALA	2.6
1	A	177	HIS	2.6
1	B	147	GLY	2.6
1	A	225	CYS	2.6
1	B	39	ASP	2.6
1	B	22	ALA	2.6
1	A	146	ILE	2.6
1	A	11	HIS	2.5
1	B	226	PHE	2.5
1	A	178	PRO	2.5
1	B	82	VAL	2.5
1	A	176	VAL	2.4
1	A	174	ILE	2.4
1	B	225	CYS	2.4
1	B	144	ILE	2.4
1	B	154	ASP	2.3
1	B	37	LYS	2.3
1	B	38	GLY	2.3
1	A	26	ALA	2.3
1	A	34	VAL	2.3
1	B	9	HIS	2.3
1	B	74	VAL	2.3
1	A	81	PRO	2.2
1	B	315	GLU	2.2
1	B	146	ILE	2.2
1	A	175	LEU	2.2
1	B	202	THR	2.1
1	A	80	THR	2.1
1	A	9	HIS	2.1
1	A	119	ALA	2.1
1	B	119	ALA	2.1
1	B	29	ALA	2.0
1	B	113	GLN	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	36	ALA	2.0
1	B	155	LEU	2.0
1	B	17	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	B	401	1/1	0.98	0.22	69,69,69,69	0
2	ZN	A	401	1/1	0.99	0.23	60,60,60,60	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.