



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 08:44 pm BST

PDB ID : 5IGY  
Title : Macrolide 2'-phosphotransferase type II - complex with GDP and erythromycin  
Authors : Berghuis, A.M.; Fong, D.H.  
Deposited on : 2016-02-28  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	<div> <div></div> <div>87%</div> <div>10%</div> <div>.</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5526 atoms, of which 2485 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Macrolide 2'-phosphotransferase II.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	295	4847	1566	2406	409	455	11	0	13	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>).



- 
- The image displays the chemical structure of Erythromycin (ERY), a 14-membered macrolide antibiotic. The structure is shown with stereochemistry and atom numbering. Key features include:
- Macrolide Ring:** A 14-membered ring containing an oxygen atom (O9) and a ketone group (C1=O1).
  - Cladinose Sugar:** A six-membered ring attached to the macrolide ring at C3, containing an oxygen atom (O4) and a hydroxyl group (OH6).
  - Desosamine Sugar:** A six-membered ring attached to the macrolide ring at C5, containing an oxygen atom (O3) and a hydroxyl group (OH06).
  - Phenylglycidyl Ether (PGE) Group:** A side chain attached to the desosamine ring at C2, consisting of a phenyl ring (C29-C33) and an epoxide ring (C34-C35).
  - Atom Numbering:** Carbons are labeled C1 through C37. Oxygen atoms are labeled O1 through O13. The nitrogen atom in the PGE group is labeled N1.
  - Stereochemistry:** Chiral centers are indicated with wedged and dashed bonds. For example, C1 is (R), C2 is (S), C3 is (R), C4 is (S), C5 is (R), C6 is (R), C7 is (S), C8 is (R), C9 is (S), C10 is (R), C11 is (S), C12 is (R), C13 is (S), C14 is (R), C15 is (S), C16 is (R), C17 is (S), C18 is (R), C19 is (S), C20 is (R), C21 is (S), C22 is (R), C23 is (S), C24 is (R), C25 is (S), C26 is (R), C27 is (S), C28 is (R), C29 is (S), C30 is (R), C31 is (S), C32 is (R), C33 is (S), C34 is (R), C35 is (S), C36 is (R), C37 is (S).


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	520	Total O 520 520	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Macrolide 2'-phosphotransferase II



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	39.87Å 80.52Å 97.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.82 – 1.45 41.82 – 1.45	Depositor EDS
% Data completeness (in resolution range)	100.0 (41.82-1.45) 100.0 (41.82-1.45)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.86 (at 1.45Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, $R_{free}$	0.170 , 0.196 0.171 , 0.196	Depositor DCC
$R_{free}$ test set	2835 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.2	Xtriage
Anisotropy	0.260	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 50.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5526	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, CA, ERY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/2555	0.58	0/3464

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2441	2406	2343	28	1
2	A	1	0	0	0	0
3	A	28	12	12	0	0
4	A	51	67	64	1	0
5	A	520	0	0	19	2
All	All	3041	2485	2419	29	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:GLU:OE1	5:A:501:HOH:O	1.84	0.96
1:A:129:ARG:NH1	5:A:503:HOH:O	2.01	0.93
1:A:213:ASN:OD1	5:A:502:HOH:O	1.93	0.86
1:A:290[A]:GLU:HG3	5:A:513:HOH:O	1.84	0.77
1:A:213:ASN:CG	5:A:502:HOH:O	2.25	0.73
1:A:290[A]:GLU:OE2	5:A:504:HOH:O	2.11	0.68
1:A:213:ASN:CB	5:A:502:HOH:O	2.46	0.63
1:A:213:ASN:HB3	5:A:502:HOH:O	1.97	0.63
1:A:54:GLU:OE1	5:A:506:HOH:O	2.16	0.61
1:A:51:PRO:HG3	1:A:56:VAL:HG21	1.81	0.61
1:A:129:ARG:HD2	5:A:503:HOH:O	2.01	0.59
1:A:292[B]:MET:HE3	5:A:604:HOH:O	2.01	0.59
1:A:53:ARG:O	1:A:59:ARG:NH2	2.39	0.56
1:A:129:ARG:NH1	5:A:519:HOH:O	2.41	0.54
1:A:228:VAL:HG21	5:A:626:HOH:O	2.08	0.54
1:A:290[A]:GLU:CG	5:A:513:HOH:O	2.50	0.54
1:A:136:SER:OG	5:A:505:HOH:O	2.15	0.52
1:A:290[B]:GLU:OE1	5:A:508:HOH:O	2.19	0.51
1:A:298:GLU:OE2	5:A:511:HOH:O	2.20	0.49
1:A:170:PHE:O	5:A:509:HOH:O	2.20	0.49
1:A:57:MET:HB2	1:A:58:PRO:HD3	1.96	0.48
1:A:4:ASP:OD1	5:A:510:HOH:O	2.20	0.47
1:A:15:HIS:CD2	1:A:86:THR:HG21	2.50	0.47
4:A:403:ERY:H343	4:A:403:ERY:H353	2.01	0.42
1:A:154:GLU:HA	1:A:157[B]:MET:HE3	2.02	0.41
1:A:104:ASP:OD1	1:A:104:ASP:C	2.59	0.41
1:A:36:VAL:HB	1:A:48:LEU:HB2	2.03	0.41
1:A:186:ASP:CG	1:A:189[A]:MET:HG3	2.42	0.40
1:A:64:LYS:HD3	1:A:82:TRP:CE3	2.56	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:926:HOH:O	5:A:927:HOH:O[2_554]	2.12	0.08
5:A:515:HOH:O	5:A:736:HOH:O[1_655]	2.12	0.08
1:A:73:TYR:OH	1:A:242:GLU:OE2[1_655]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/302 (101%)	298 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/256 (102%)	260 (100%)	1 (0%)	91	80

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	254	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	ERY	A	403	-	53,53,53	3.77	29 (54%)	82,82,82	1.96	26 (31%)
3	GDP	A	402	2	24,30,30	1.12	2 (8%)	31,47,47	1.98	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ERY	A	403	-	-	0/72/107/107	0/3/3/3
3	GDP	A	402	2	-	1/12/32/32	0/3/3/3

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	403	ERY	C10-C11	-10.71	1.40	1.54
4	A	403	ERY	C12-C13	-9.39	1.37	1.54
4	A	403	ERY	O10-C6	-7.63	1.31	1.44
4	A	403	ERY	O13-C12	-6.51	1.33	1.44
4	A	403	ERY	C35-C12	-6.49	1.41	1.52
4	A	403	ERY	O5-C16	-6.12	1.32	1.44
4	A	403	ERY	C23-C24	-5.59	1.41	1.53
4	A	403	ERY	C32-C6	-5.18	1.44	1.52
4	A	403	ERY	C16-C17	-5.09	1.41	1.53
4	A	403	ERY	O2-C1	5.00	1.46	1.34
4	A	403	ERY	C10-C9	-4.89	1.45	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	403	ERY	C4-C5	-4.65	1.43	1.55
4	A	403	ERY	O9-C26	-4.39	1.36	1.44
4	A	403	ERY	C33-C8	-4.28	1.38	1.52
4	A	403	ERY	C25-C24	-4.15	1.44	1.53
3	A	402	GDP	C6-C5	3.87	1.48	1.41
4	A	403	ERY	C19-C16	-3.87	1.43	1.52
4	A	403	ERY	C7-C6	-3.58	1.48	1.54
4	A	403	ERY	C25-C26	-3.45	1.45	1.51
4	A	403	ERY	O2-C13	-3.39	1.40	1.46
4	A	403	ERY	C18-C17	-3.28	1.45	1.53
4	A	403	ERY	O11-C9	3.10	1.26	1.21
4	A	403	ERY	C24-N1	-2.94	1.41	1.48
4	A	403	ERY	O7-C22	2.79	1.49	1.41
4	A	403	ERY	C8-C9	-2.79	1.43	1.52
4	A	403	ERY	C34-C10	-2.70	1.47	1.53
4	A	403	ERY	O6-C17	-2.32	1.38	1.42
4	A	403	ERY	C2-C3	-2.23	1.49	1.55
4	A	403	ERY	O5-C20	2.21	1.50	1.43
3	A	402	GDP	C5-C4	2.07	1.46	1.40
4	A	403	ERY	O4-C14	-2.06	1.37	1.42

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	GDP	C5-C6-N1	-6.67	114.31	123.43
3	A	402	GDP	C6-N1-C2	5.53	124.72	115.93
4	A	403	ERY	O13-C12-C35	-4.72	98.03	107.78
4	A	403	ERY	O5-C16-C17	4.41	110.35	103.81
4	A	403	ERY	O9-C26-C25	4.25	115.65	109.14
4	A	403	ERY	O11-C9-C8	-4.22	113.43	121.26
4	A	403	ERY	C30-C2-C3	3.81	121.56	112.92
4	A	403	ERY	C8-C9-C10	3.50	125.18	119.10
4	A	403	ERY	C22-O7-C5	3.41	122.18	116.25
4	A	403	ERY	O4-C18-C17	3.35	116.02	110.03
4	A	403	ERY	C11-C10-C9	3.32	114.94	109.57
3	A	402	GDP	C6-C5-C4	-3.14	117.80	120.80
4	A	403	ERY	C13-C12-C11	3.12	114.44	108.28
4	A	403	ERY	O5-C16-C19	-3.07	105.81	110.92
3	A	402	GDP	C4-C5-N7	-2.95	106.32	109.40
4	A	403	ERY	O2-C1-O1	-2.88	118.56	123.94
4	A	403	ERY	C3-C4-C5	2.77	117.16	111.19
4	A	403	ERY	C22-O9-C26	-2.64	108.73	112.91

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	ERY	C16-C17-C18	-2.54	107.24	111.14
4	A	403	ERY	O7-C5-C6	2.54	109.52	106.39
4	A	403	ERY	C20-O5-C16	2.53	122.84	117.55
4	A	403	ERY	C31-C4-C3	-2.47	106.96	111.40
4	A	403	ERY	C29-N1-C24	2.46	120.51	113.11
4	A	403	ERY	C7-C8-C9	2.29	117.25	113.32
4	A	403	ERY	C33-C8-C7	-2.28	105.61	109.81
4	A	403	ERY	C12-C11-C10	2.21	119.19	116.43
4	A	403	ERY	C25-C24-C23	-2.16	106.85	109.97
4	A	403	ERY	C25-C24-N1	-2.15	109.60	115.67
4	A	403	ERY	C36-C13-C12	2.15	119.26	115.20
4	A	403	ERY	O6-C17-C16	2.14	115.12	111.12
3	A	402	GDP	N3-C2-N1	-2.11	124.41	127.22

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	GDP	C4'-C5'-O5'-PA


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	ERY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

## Ligand ERY A 403





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	295/302 (97%)	-0.19	3 (1%) 82 84	6, 17, 37, 56	4 (1%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	107	ILE	7.0
1	A	282[A]	LEU	2.2
1	A	4	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ERY	A	403	51/51	0.95	0.07	5,10,20,23	0
3	GDP	A	402	28/28	0.98	0.06	7,12,17,19	0
2	CA	A	401	1/1	1.00	0.05	11,11,11,11	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.