



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 01:18 pm BST

PDB ID : 5III  
Title : Crystal structure of the pre-catalytic ternary complex of DNA polymerase lambda with a templating 8-oxo-dG and an incoming dATP  
Authors : Burak, M.J.; Guja, K.E.; Garcia-Diaz, M.  
Deposited on : 2016-03-01  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	<div> <div>10%</div> <div>89%</div> <div>5% • 6%</div> </div>
2	D	4	<div> <div>75%</div> <div>25%</div> </div>
3	P	6	<div> <div>100%</div> </div>
4	T	11	<div> <div>82%</div> <div>9%</div> <div>9%</div> </div>

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3173 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase lambda.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	0	0
			2381	1506	425	439	11			

- Molecule 2 is a DNA chain called DNA (5'-D(P\*GP\*CP\*CP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			

- Molecule 3 is a DNA chain called DNA (5'-D(\*CP\*AP\*GP\*TP\*AP\*(DOC))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	6	Total	C	N	O	P	0	0	0
			118	58	23	32	5			

- Molecule 4 is a DNA chain called DNA (5'-D(\*CP\*GP\*GP\*CP\*(8OG)P\*GP\*TP\*AP\*CP\*TP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	T	11	Total	C	N	O	P	0	0	0
			226	107	43	66	10			

- Molecule 5 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			30	10	5	12	3		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total Na 2 2	0	0

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Mg 1 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	243	Total O 243 243	0	0
8	D	3	Total O 3 3	0	0
8	P	28	Total O 28 28	0	0
8	T	58	Total O 58 58	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase lambda



- Molecule 2: DNA (5'-D(P\*GP\*CP\*CP\*G)-3')




- Molecule 3: DNA (5'-D(\*CP\*AP\*GP\*TP\*AP\*(DOC))-3')



There are no outlier residues recorded for this chain.

- Molecule 4: DNA (5'-D(\*CP\*GP\*GP\*CP\*(8OG)P\*GP\*TP\*AP\*CP\*TP\*G)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.85Å 62.68Å 140.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.70 – 1.80 41.70 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (41.70-1.80) 93.8 (41.70-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.98 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.201 , 0.225 0.203 , 0.223	Depositor DCC
$R_{free}$ test set	2259 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.0	Xtriage
Anisotropy	0.641	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3173	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, DTP, DOC, 8OG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/2431	0.48	0/3295
2	D	1.27	1/92 (1.1%)	0.83	0/138
3	P	0.79	0/112	0.95	0/171
4	T	0.82	0/226	1.17	3/345 (0.9%)
All	All	0.47	1/2861 (0.0%)	0.61	3/3949 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	DG	OP3-P	-10.34	1.48	1.61

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	T	7	DT	C1'-O4'-C4'	-5.61	104.49	110.10
4	T	7	DT	O5'-P-OP2	-5.50	100.75	105.70
4	T	7	DT	O4'-C4'-C3'	-5.06	102.48	104.50

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2381	0	2285	13	0
2	D	83	0	45	3	0
3	P	118	0	69	0	0
4	T	226	0	125	2	0
5	A	30	0	12	1	0
6	A	2	0	0	0	0
7	A	1	0	0	0	0
8	A	243	0	0	2	0
8	D	3	0	0	0	0
8	P	28	0	0	0	0
8	T	58	0	0	0	0
All	All	3173	0	2536	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:396:GLU:HG3	1:A:414:ALA:HB2	1.75	0.69
5:A:601:DTP:O2G	8:A:701:HOH:O	2.11	0.68
1:A:308:ARG:NH2	2:D:1:DG:OP2	2.33	0.62
1:A:253:ASN:N	1:A:253:ASN:OD1	2.39	0.56
1:A:290:HIS:HE1	1:A:291:LYS:HE2	1.72	0.55
1:A:521:LYS:NZ	4:T:4:DC:OP1	2.35	0.54
2:D:3:DC:H2'	2:D:4:DG:C8	2.44	0.53
2:D:2:DC:H2'	2:D:3:DC:C6	2.45	0.52
1:A:396:GLU:OE1	8:A:702:HOH:O	2.19	0.50
1:A:312:LYS:HG3	1:A:322:LEU:HD11	1.95	0.48
1:A:308:ARG:O	1:A:312:LYS:HD3	2.14	0.47
1:A:511:HIS:CE1	1:A:514:ARG:HH21	2.34	0.46
1:A:516:MET:HG2	1:A:527:LEU:CD2	2.45	0.46
1:A:290:HIS:CE1	1:A:291:LYS:HE2	2.52	0.43
1:A:528:SER:HB2	4:T:7:DT:H5'	2.02	0.41
1:A:554:PRO:HG2	1:A:558:ASP:OD2	2.22	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/334 (93%)	300 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/280 (84%)	232 (98%)	4 (2%)	60	51

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	293	VAL
1	A	396	GLU
1	A	506	PHE
1	A	516	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	8OG	T	5	4	18,25,26	5.03	11 (61%)	21,37,40	1.88	4 (19%)
3	DOC	P	6	3,4	14,19,20	4.45	10 (71%)	13,26,29	1.91	3 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	8OG	T	5	4	-	1/3/21/22	0/3/3/3
3	DOC	P	6	3,4	-	0/4/18/19	0/2/2/2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	T	5	8OG	C8-N7	13.17	1.48	1.34
4	T	5	8OG	C3'-C4'	-8.25	1.30	1.53
3	P	6	DOC	C3'-C2'	-7.89	1.32	1.54
4	T	5	8OG	O4'-C4'	7.81	1.62	1.45
3	P	6	DOC	O4'-C4'	-7.50	1.29	1.44
3	P	6	DOC	O4'-C1'	5.82	1.55	1.42
3	P	6	DOC	C1'-N1	-5.81	1.32	1.49
3	P	6	DOC	C6-N1	5.54	1.42	1.35
4	T	5	8OG	C4-N3	5.20	1.43	1.35
4	T	5	8OG	C6-N1	5.18	1.42	1.33
4	T	5	8OG	C2-N1	5.09	1.44	1.35
4	T	5	8OG	C6-C5	4.42	1.49	1.41
4	T	5	8OG	O4'-C1'	-4.32	1.32	1.42
3	P	6	DOC	C2-N3	4.10	1.46	1.38
3	P	6	DOC	C6-C5	3.95	1.46	1.38
4	T	5	8OG	C2-N2	3.94	1.41	1.33

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	6	DOC	C4-N3	3.56	1.41	1.35
4	T	5	8OG	O3'-C3'	2.80	1.49	1.43
3	P	6	DOC	C5-C4	2.26	1.46	1.41
4	T	5	8OG	C2-N3	2.19	1.45	1.34
3	P	6	DOC	C4-N4	2.19	1.41	1.35

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	T	5	8OG	N3-C2-N1	-5.25	120.22	127.22
3	P	6	DOC	C2-N3-C4	4.15	120.55	116.34
4	T	5	8OG	C2-N3-C4	3.89	119.80	115.36
4	T	5	8OG	C5-C6-N1	-3.10	119.19	123.43
3	P	6	DOC	C4'-O4'-C1'	-2.86	107.11	109.81
4	T	5	8OG	C6-N1-C2	2.61	120.08	115.93
3	P	6	DOC	C3'-C2'-C1'	2.39	105.54	102.78

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	T	5	8OG	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	DTP	A	601	7	26,32,32	3.20	11 (42%)	30,50,50	2.16	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DTP	A	601	7	-	3/18/34/34	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	601	DTP	C2'-C3'	-10.54	1.25	1.52
5	A	601	DTP	O4'-C4'	-6.81	1.29	1.45
5	A	601	DTP	C1'-N9	-5.49	1.33	1.49
5	A	601	DTP	C3'-C4'	4.37	1.65	1.53
5	A	601	DTP	O4'-C1'	3.51	1.50	1.42
5	A	601	DTP	O3'-C3'	3.14	1.50	1.43
5	A	601	DTP	C2-N3	3.13	1.37	1.32
5	A	601	DTP	C6-N6	2.34	1.42	1.34
5	A	601	DTP	C2-N1	2.21	1.38	1.33
5	A	601	DTP	C2'-C1'	2.07	1.58	1.52
5	A	601	DTP	C5-C4	-2.03	1.35	1.40

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	601	DTP	C5-C6-N6	7.89	132.35	120.35
5	A	601	DTP	N3-C2-N1	-5.82	119.58	128.68
5	A	601	DTP	N6-C6-N1	-5.06	108.08	118.57
5	A	601	DTP	PB-O3B-PG	-2.16	125.41	132.83
5	A	601	DTP	C2'-C1'-N9	-2.12	109.39	114.27

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	601	DTP	PB-O3B-PG-O2G
5	A	601	DTP	PB-O3B-PG-O3G
5	A	601	DTP	PA-O3A-PB-O2B

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	601	DTP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	314/334 (94%)	0.39	35 (11%) 5 4	18, 31, 65, 82	0
2	D	4/4 (100%)	-0.49	0 100 100	35, 35, 36, 38	0
3	P	5/6 (83%)	-0.30	0 100 100	19, 19, 21, 21	0
4	T	10/11 (90%)	-0.42	0 100 100	21, 25, 37, 39	0
All	All	333/355 (93%)	0.35	35 (10%) 6 4	18, 31, 64, 82	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	293	VAL	7.4
1	A	535	ALA	7.0
1	A	289	PHE	5.7
1	A	547	PRO	5.5
1	A	252	HIS	4.6
1	A	534	THR	4.6
1	A	406	PHE	4.5
1	A	254	LEU	4.4
1	A	317	LEU	4.1
1	A	253	ASN	4.0
1	A	255	HIS	3.9
1	A	299	ALA	3.8
1	A	290	HIS	3.7
1	A	319	SER	3.6
1	A	294	THR	3.6
1	A	301	SER	3.2
1	A	316	ILE	3.1
1	A	297	GLN	3.0
1	A	313	ILE	2.9
1	A	288	SER	2.8
1	A	296	TYR	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	567	TYR	2.7
1	A	575	TRP	2.6
1	A	292	PRO	2.6
1	A	522	THR	2.6
1	A	314	ILE	2.6
1	A	321	HIS	2.4
1	A	536	VAL	2.4
1	A	286	LEU	2.3
1	A	320	GLY	2.2
1	A	566	PRO	2.2
1	A	295	SER	2.1
1	A	303	PRO	2.1
1	A	310	ALA	2.1
1	A	285	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	8OG	T	5	23/24	0.93	0.11	23,27,43,47	0
3	DOC	P	6	18/19	0.98	0.14	18,21,24,27	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	NA	A	603	1/1	0.97	0.07	31,31,31,31	0
5	DTP	A	601	30/30	0.98	0.12	19,22,25,27	0


*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
7	MG	A	604	1/1	0.99	0.13	21,21,21,21	0
6	NA	A	602	1/1	0.99	0.07	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.