



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 04:00 am BST

PDB ID : 6IIK  
Title : USP14 catalytic domain with IU1  
Authors : Mei, Z.Q.; Wang, Y.W.; He, W.; Wang, F.  
Deposited on : 2018-10-06  
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	399	<div> <div>6%</div> <div> <div></div> <div>76%</div> <div>7%</div> <div>•</div> <div>16%</div> </div> </div>
1	B	399	<div> <div>14%</div> <div> <div></div> <div>71%</div> <div>12%</div> <div>•</div> <div>16%</div> </div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5754 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	0	0
			2700	1721	449	511	19			
1	B	334	Total	C	N	O	S	0	0	0
			2688	1712	448	509	19			

- Molecule 2 is 1-[1-(4-fluorophenyl)-2,5-dimethyl-1H-pyrrol-3-yl]-2-(pyrrolidin-1-yl)ethan-1-one (three-letter code: IU1) (formula: C<sub>18</sub>H<sub>21</sub>FN<sub>2</sub>O) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			22	18	1	2	1		
2	B	1	Total	C	F	N	O	0	0
			22	18	1	2	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	196	Total 196	O 196	0	0
3	B	126	Total 126	O 126	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 14



- Molecule 1: Ubiquitin carboxyl-terminal hydrolase 14



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.87Å 81.67Å 107.37Å 90.00° 93.68° 90.00°	Depositor
Resolution (Å)	29.22 – 1.97 29.22 – 1.97	Depositor EDS
% Data completeness (in resolution range)	96.7 (29.22-1.97) 96.7 (29.22-1.97)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.79 (at 1.96Å)	Xtriage
Refinement program	PHENIX phenix.refine: 1.8 _1069	Depositor
R, $R_{free}$	0.189 , 0.218 0.190 , 0.219	Depositor DCC
$R_{free}$ test set	3455 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.2	Xtriage
Anisotropy	0.530	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 51.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5754	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IU1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	0/2751	0.56	1/3704 (0.0%)
1	B	0.36	0/2738	0.52	0/3685
All	All	0.40	0/5489	0.54	1/7389 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	424	LEU	CA-CB-CG	6.98	131.35	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2700	0	2684	20	0
1	B	2688	0	2675	30	0
2	A	22	0	0	0	0
2	B	22	0	0	0	0
3	A	196	0	0	9	0
3	B	126	0	0	5	0
All	All	5754	0	5359	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:472:TRP:O	3:B:601:HOH:O	1.89	0.88
1:B:206:GLN:OE1	3:B:602:HOH:O	2.05	0.73
1:A:146:MET:SD	3:A:794:HOH:O	2.48	0.71
1:A:301:GLN:OE1	3:A:603:HOH:O	2.08	0.71
1:A:340:ASN:N	3:A:607:HOH:O	2.29	0.66
1:B:291:LYS:HG2	1:B:358:LEU:HD21	1.79	0.65
1:B:280:ASN:O	1:B:280:ASN:ND2	2.30	0.65
1:B:402:GLU:O	1:B:484:ARG:NH1	2.32	0.62
1:A:418:TYR:HB3	1:A:480:TYR:HB3	1.84	0.59
1:A:303:PRO:O	1:A:306:GLN:NE2	2.38	0.57
1:B:374:PHE:CE2	1:B:483:ARG:HD3	2.40	0.56
1:B:375:LYS:NZ	3:B:605:HOH:O	2.40	0.55
1:B:462:ASP:OD1	1:B:465:ARG:NH2	2.39	0.55
1:B:295:GLN:HE22	1:B:314:SER:HB3	1.73	0.54
1:B:418:TYR:HB3	1:B:480:TYR:HB3	1.91	0.52
1:B:259:GLU:OE2	1:B:307:ARG:NH2	2.43	0.51
1:B:280:ASN:O	1:B:282:GLU:N	2.41	0.51
1:B:352:MET:HE1	1:B:483:ARG:HD2	1.93	0.50
1:A:330:ARG:NH2	3:A:604:HOH:O	2.32	0.50
1:B:173:ILE:H	1:B:173:ILE:HD13	1.77	0.49
1:A:446:GLU:HG3	1:A:457:ILE:HG23	1.95	0.48
1:A:468:GLY:HA2	1:A:474:ILE:HG12	1.95	0.48
1:B:134:LYS:NZ	3:B:611:HOH:O	2.46	0.48
1:A:220:ASP:N	1:A:220:ASP:OD1	2.47	0.48
1:B:297:GLU:HB2	1:B:310:LEU:HD11	1.95	0.48
1:A:133:LEU:HD11	1:A:156:LEU:HD23	1.96	0.47
1:B:374:PHE:CE2	1:B:485:VAL:HB	2.49	0.47
1:A:139:ALA:HA	3:A:688:HOH:O	2.14	0.47
1:A:267:LYS:NZ	3:A:608:HOH:O	2.33	0.46
1:B:330:ARG:HB3	1:B:343:VAL:HG13	1.97	0.45
1:A:285:TYR:OH	3:A:601:HOH:O	1.81	0.45
1:A:330:ARG:O	1:A:332:PHE:N	2.50	0.44
1:B:284:LYS:HD3	1:B:345:LYS:HZ3	1.82	0.44
1:B:130:LYS:HB3	1:B:130:LYS:HE2	1.86	0.43
1:A:141:ARG:HA	1:B:173:ILE:HD11	1.99	0.43
1:B:468:GLY:HA2	1:B:474:ILE:HG12	2.01	0.43
1:B:447:TRP:CD2	1:B:463:ILE:HG13	2.54	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:129:LEU:HD23	1:B:129:LEU:HA	1.84	0.42
1:B:260:SER:HA	1:B:261:GLU:HA	1.90	0.42
1:A:447:TRP:CD2	1:A:463:ILE:HG13	2.55	0.42
1:B:371:ARG:O	1:B:375:LYS:HG2	2.20	0.42
1:A:177:GLN:NE2	3:A:602:HOH:O	2.06	0.42
1:B:442:ARG:HD3	1:B:446:GLU:OE2	2.20	0.41
1:A:330:ARG:NE	3:A:604:HOH:O	2.23	0.41
1:B:446:GLU:HG3	1:B:457:ILE:HG23	2.03	0.41
1:B:148:SER:HB2	3:B:696:HOH:O	2.21	0.41
1:A:257:CYS:HB2	1:A:311:TYR:CE2	2.56	0.40
1:A:443:LYS:HB2	1:A:446:GLU:HB3	2.02	0.40
1:B:263:GLU:OE2	1:B:303:PRO:HG2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/399 (82%)	313 (96%)	12 (4%)	2 (1%)	25	14
1	B	326/399 (82%)	311 (95%)	13 (4%)	2 (1%)	25	14
All	All	653/798 (82%)	624 (96%)	25 (4%)	4 (1%)	25	14

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	331	PHE
1	B	264	GLU
1	A	168	SER
1	B	260	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/358 (84%)	291 (97%)	8 (3%)	44	35
1	B	297/358 (83%)	287 (97%)	10 (3%)	37	25
All	All	596/716 (83%)	578 (97%)	18 (3%)	41	29

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	168	SER
1	A	202	GLU
1	A	220	ASP
1	A	287	PHE
1	A	332	PHE
1	A	424	LEU
1	A	443	LYS
1	A	484	ARG
1	B	173	ILE
1	B	189	LYS
1	B	241	SER
1	B	287	PHE
1	B	299	THR
1	B	304	THR
1	B	308	ASN
1	B	314	SER
1	B	343	VAL
1	B	352	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	301	GLN
1	B	295	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	IU1	A	501	-	23,24,24	1.94	1 (4%)	25,34,34	2.49	7 (28%)
2	IU1	B	501	-	23,24,24	1.93	1 (4%)	25,34,34	2.46	11 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IU1	A	501	-	-	0/8/19/19	0/3/3/3
2	IU1	B	501	-	-	2/8/19/19	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	IU1	C5-N8	-8.63	1.34	1.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	IU1	C5-N8	-8.58	1.34	1.46

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	IU1	C21-C9-C10	-6.88	116.11	128.72
2	B	501	IU1	C21-C9-C10	-6.25	117.27	128.72
2	A	501	IU1	C6-C5-C4	-4.59	114.54	121.33
2	A	501	IU1	C4-C5-N8	4.48	125.45	119.19
2	A	501	IU1	C22-C12-N8	4.36	127.86	122.37
2	B	501	IU1	C6-C5-C4	-4.11	115.23	121.33
2	B	501	IU1	C15-N16-C17	-3.86	108.13	113.25
2	B	501	IU1	C22-C12-N8	3.58	126.88	122.37
2	B	501	IU1	C19-C20-N16	3.52	108.03	103.92
2	A	501	IU1	C7-C6-C5	3.17	123.68	119.07
2	A	501	IU1	C7-C2-C3	-3.07	118.75	122.83
2	B	501	IU1	C17-N16-C20	2.83	106.79	104.04
2	B	501	IU1	C7-C6-C5	2.78	123.12	119.07
2	B	501	IU1	C7-C2-C3	-2.74	119.19	122.83
2	A	501	IU1	C3-C4-C5	2.73	123.04	119.07
2	B	501	IU1	C3-C4-C5	2.65	122.92	119.07
2	B	501	IU1	C4-C5-N8	2.36	122.50	119.19
2	B	501	IU1	C6-C5-N8	2.20	122.27	119.19

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	501	IU1	O14-C13-C15-N16
2	B	501	IU1	C11-C13-C15-N16

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	335/399 (83%)	0.36	24 (7%) 15 17	16, 36, 74, 98	0
1	B	334/399 (83%)	0.82	57 (17%) 1 1	27, 47, 97, 128	0
All	All	669/798 (83%)	0.59	81 (12%) 4 4	16, 43, 87, 128	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	260	SER	8.5
1	A	332	PHE	8.3
1	B	332	PHE	7.9
1	A	146	MET	6.5
1	B	331	PHE	6.5
1	B	259	GLU	5.6
1	B	485	VAL	5.2
1	A	377	LEU	4.8
1	B	220	ASP	4.8
1	B	193	GLY	4.7
1	B	305	LEU	4.6
1	B	401	TYR	4.5
1	A	145	GLU	4.5
1	B	146	MET	4.4
1	B	486	GLU	4.3
1	B	311	TYR	4.3
1	B	258	THR	4.1
1	A	220	ASP	4.0
1	B	145	GLU	3.8
1	B	102	GLU	3.7
1	B	309	ALA	3.7
1	B	307	ARG	3.6
1	B	477	VAL	3.5
1	B	191	GLU	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	444	GLN	3.5
1	A	340	ASN	3.4
1	A	102	GLU	3.4
1	B	310	LEU	3.4
1	B	304	THR	3.4
1	B	400	LYS	3.4
1	A	123	ILE	3.3
1	B	144	GLY	3.3
1	A	341	ALA	3.2
1	B	137	ALA	3.2
1	B	192	GLN	3.2
1	A	376	ASP	3.2
1	B	303	PRO	3.1
1	A	477	VAL	3.1
1	B	341	ALA	3.0
1	B	257	CYS	3.0
1	A	375	LYS	3.0
1	B	261	GLU	3.0
1	A	192	GLN	3.0
1	B	375	LYS	3.0
1	A	401	TYR	3.0
1	B	301	GLN	2.9
1	B	152	ILE	2.8
1	A	281	GLN	2.7
1	A	262	GLU	2.7
1	B	424	LEU	2.7
1	B	240	LYS	2.7
1	A	191	GLU	2.6
1	B	278	PHE	2.6
1	B	445	ASP	2.6
1	A	485	VAL	2.6
1	B	174	ILE	2.6
1	A	193	GLY	2.5
1	B	306	GLN	2.5
1	B	262	GLU	2.5
1	A	331	PHE	2.5
1	B	443	LYS	2.4
1	B	281	GLN	2.4
1	B	175	LEU	2.4
1	B	300	LYS	2.4
1	A	261	GLU	2.4
1	B	123	ILE	2.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	190	GLY	2.4
1	B	176	LEU	2.3
1	B	265	VAL	2.2
1	B	422	ALA	2.2
1	B	219	GLU	2.2
1	A	219	GLU	2.1
1	B	264	GLU	2.1
1	A	444	GLN	2.1
1	B	296	GLU	2.1
1	B	122	CYS	2.1
1	B	299	THR	2.0
1	B	263	GLU	2.0
1	A	424	LEU	2.0
1	B	478	LEU	2.0
1	B	267	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	IU1	B	501	22/22	0.90	0.16	29,43,72,75	0
2	IU1	A	501	22/22	0.92	0.13	28,37,52,54	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around IU1 B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around IU1 A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.