



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 14, 2020 – 10:58 am BST

PDB ID : 2IJX  
Title : Crystal structure of PCNA3 monomer from *Sulfolobus solfataricus*.  
Authors : Hlinkova, V.; Ling, H.  
Deposited on : 2006-10-01  
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	244	77% 21% ..
1	B	244	87% 11% .
1	C	244	80% 16% ..
1	D	244	85% 14% .

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8406 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase sliding clamp A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	244	Total	C	N	O	S	0	3	0
			1948	1239	314	391	4			
1	B	244	Total	C	N	O	S	0	6	0
			1954	1243	313	393	5			
1	C	244	Total	C	N	O	S	0	5	0
			1955	1243	312	395	5			
1	D	244	Total	C	N	O	S	0	0	0
			1934	1229	312	389	4			

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	123	Total	O	0	0
			123	123		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	150	Total 150	O 150	0	0
3	C	119	Total 119	O 119	0	0
3	D	143	Total 143	O 143	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase sliding clamp A




- Molecule 1: DNA polymerase sliding clamp A



- Molecule 1: DNA polymerase sliding clamp A



- Molecule 1: DNA polymerase sliding clamp A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.77Å 85.77Å 264.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.62 – 1.90 29.56 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.62-1.90) 96.3 (29.56-1.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.204 , 0.251 0.210 , 0.258	Depositor DCC
$R_{free}$ test set	1530 reflections (2.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.2	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 51.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	8406	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.70	0/1983	0.84	1/2671 (0.0%)
1	B	0.76	0/2002	0.80	2/2696 (0.1%)
1	C	0.78	2/1999 (0.1%)	0.86	5/2694 (0.2%)
1	D	0.72	0/1960	0.81	1/2641 (0.0%)
All	All	0.74	2/7944 (0.0%)	0.83	9/10702 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	159	GLU	CD-OE1	5.25	1.31	1.25
1	C	156	GLU	CD-OE2	5.13	1.31	1.25

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	159	GLU	OE1-CD-OE2	9.17	134.30	123.30
1	C	41	ARG	NE-CZ-NH1	7.11	123.85	120.30
1	C	41	ARG	NE-CZ-NH2	-6.73	116.94	120.30
1	D	18	LEU	CA-CB-CG	6.49	130.23	115.30
1	C	124	ILE	CG1-CB-CG2	5.94	124.48	111.40

There are no chirality outliers.



All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	103	GLY	Peptide
1	A	120	GLU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1948	0	1969	38	0
1	B	1954	0	1977	19	1
1	C	1955	0	1970	39	1
1	D	1934	0	1945	25	0
2	A	12	0	18	0	0
2	B	20	0	30	0	0
2	C	32	0	48	1	0
2	D	16	0	24	1	0
3	A	123	0	0	1	0
3	B	150	0	0	4	0
3	C	119	0	0	5	0
3	D	143	0	0	7	0
All	All	8406	0	7981	112	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 112 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:20:ARG:HB2	3:C:643:HOH:O	1.60	0.98
1:C:156:GLU:OE2	1:C:158:HIS:HE1	1.46	0.97
1:A:100:ASN:HD21	1:A:106:ASN:ND2	1.62	0.97
1:A:100:ASN:HD21	1:A:106:ASN:HD21	1.10	0.95
1:C:124:ILE:HG23	1:D:124:ILE:CG1	2.03	0.88

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:6:ASP:OD2	1:C:6:ASP:OD2[3_545]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/244 (100%)	239 (98%)	3 (1%)	3 (1%)	13	4
1	B	248/244 (102%)	244 (98%)	4 (2%)	0	100	100
1	C	247/244 (101%)	242 (98%)	4 (2%)	1 (0%)	34	24
1	D	242/244 (99%)	239 (99%)	1 (0%)	2 (1%)	19	9
All	All	982/976 (101%)	964 (98%)	12 (1%)	6 (1%)	25	15

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	105	THR
1	A	168	GLY
1	C	125	ASN
1	D	62	ASN
1	A	120	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/220 (101%)	209 (94%)	14 (6%)	18	8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	226/220 (103%)	214 (95%)	12 (5%)	22	13
1	C	225/220 (102%)	211 (94%)	14 (6%)	18	9
1	D	220/220 (100%)	209 (95%)	11 (5%)	24	15
All	All	894/880 (102%)	843 (94%)	51 (6%)	20	11

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	165	LYS
1	C	41	ARG
1	D	178	LYS
1	B	191	GLU
1	C	62	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	16	GLN
1	C	70	ASN
1	D	158	HIS
1	C	62	ASN
1	C	158	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	C	603	-	3,3,3	0.44	0	2,2,2	0.58	0
2	EDO	D	616	-	3,3,3	0.37	0	2,2,2	0.56	0
2	EDO	D	610	-	3,3,3	0.43	0	2,2,2	0.24	0
2	EDO	C	606	-	3,3,3	0.35	0	2,2,2	0.20	0
2	EDO	D	619	-	3,3,3	0.42	0	2,2,2	0.42	0
2	EDO	C	602	-	3,3,3	0.63	0	2,2,2	0.52	0
2	EDO	D	608	-	3,3,3	0.47	0	2,2,2	0.32	0
2	EDO	B	613	-	3,3,3	0.50	0	2,2,2	0.06	0
2	EDO	C	611	-	3,3,3	0.48	0	2,2,2	0.62	0
2	EDO	B	612	-	3,3,3	0.53	0	2,2,2	0.23	0
2	EDO	B	617	-	3,3,3	0.65	0	2,2,2	0.52	0
2	EDO	C	615	-	3,3,3	0.51	0	2,2,2	0.21	0
2	EDO	A	614	-	3,3,3	0.61	0	2,2,2	0.65	0
2	EDO	C	620	-	3,3,3	0.66	0	2,2,2	0.17	0
2	EDO	C	618	-	3,3,3	0.84	0	2,2,2	0.60	0
2	EDO	A	604	-	3,3,3	0.40	0	2,2,2	0.22	0
2	EDO	A	609	-	3,3,3	0.60	0	2,2,2	0.26	0
2	EDO	B	605	-	3,3,3	0.34	0	2,2,2	0.45	0
2	EDO	C	607	-	3,3,3	0.54	0	2,2,2	0.05	0
2	EDO	B	601	-	3,3,3	0.55	0	2,2,2	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	C	603	-	-	0/1/1/1	-
2	EDO	D	616	-	-	0/1/1/1	-
2	EDO	D	610	-	-	1/1/1/1	-
2	EDO	C	606	-	-	1/1/1/1	-
2	EDO	D	619	-	-	1/1/1/1	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	C	602	-	-	0/1/1/1	-
2	EDO	D	608	-	-	0/1/1/1	-
2	EDO	B	613	-	-	1/1/1/1	-
2	EDO	C	611	-	-	1/1/1/1	-
2	EDO	B	612	-	-	1/1/1/1	-
2	EDO	B	617	-	-	1/1/1/1	-
2	EDO	C	615	-	-	0/1/1/1	-
2	EDO	A	614	-	-	1/1/1/1	-
2	EDO	C	620	-	-	1/1/1/1	-
2	EDO	C	618	-	-	1/1/1/1	-
2	EDO	A	604	-	-	0/1/1/1	-
2	EDO	A	609	-	-	1/1/1/1	-
2	EDO	B	605	-	-	0/1/1/1	-
2	EDO	C	607	-	-	0/1/1/1	-
2	EDO	B	601	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	619	EDO	O1-C1-C2-O2
2	B	612	EDO	O1-C1-C2-O2
2	C	620	EDO	O1-C1-C2-O2
2	C	618	EDO	O1-C1-C2-O2
2	A	614	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	610	EDO	1	0
2	C	618	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.