



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 10:31 AM BST

PDB ID : 4IKM  
Title : X-ray structure of CARD8 CARD domain  
Authors : Jin, T.; Huang, M.; Smith, P.; Jiang, J.; Xiao, T.  
Deposited on : 2012-12-26  
Resolution : 2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	469	<div> <div>2%</div> <div> <div></div> <div>90%</div> <div>8%</div> </div> </div>
2	B	2	<div> <div>100%</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3703 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein, Caspase recruitment domain-containing protein 8.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	463	Total	C	I	N	O	S	2	0	0
			3601	2293	13	594	692	9			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP P0AEX9
A	83	ALA	ASP	conflict	UNP P0AEX9
A	84	ALA	LYS	conflict	UNP P0AEX9
A	173	ALA	GLU	conflict	UNP P0AEX9
A	174	ALA	ASN	conflict	UNP P0AEX9
A	240	ALA	LYS	conflict	UNP P0AEX9
A	360	ALA	GLU	conflict	UNP P0AEX9
A	363	ALA	LYS	conflict	UNP P0AEX9
A	364	ALA	ASP	conflict	UNP P0AEX9
A	368	ASN	-	expression tag	UNP P0AEX9
A	369	ALA	-	expression tag	UNP P0AEX9
A	370	VAL	-	expression tag	UNP P0AEX9
A	371	ASP	-	expression tag	UNP P0AEX9
A	459	ALA	-	expression tag	UNP Q9Y2G2
A	460	ALA	-	expression tag	UNP Q9Y2G2
A	461	ALA	-	expression tag	UNP Q9Y2G2
A	462	LEU	-	expression tag	UNP Q9Y2G2
A	463	GLU	-	expression tag	UNP Q9Y2G2
A	464	HIS	-	expression tag	UNP Q9Y2G2
A	465	HIS	-	expression tag	UNP Q9Y2G2
A	466	HIS	-	expression tag	UNP Q9Y2G2
A	467	HIS	-	expression tag	UNP Q9Y2G2
A	468	HIS	-	expression tag	UNP Q9Y2G2
A	469	HIS	-	expression tag	UNP Q9Y2G2

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	I	0	0
			2	2		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	69	Total	O	0	0
			69	69		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-binding periplasmic protein, Caspase recruitment domain-containing protein 8



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.88Å 94.88Å 219.39Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.44 – 2.46 47.44 – 2.46	Depositor EDS
% Data completeness (in resolution range)	99.3 (47.44-2.46) 96.5 (47.44-2.46)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.13 (at 2.45Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1217)	Depositor
R, $R_{free}$	0.201 , 0.253 0.202 , 0.252	Depositor DCC
$R_{free}$ test set	1122 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	64.2	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 51.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3703	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, GLC, IYR, TYI, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.27	0/3551	0.44	0/4800

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3601	0	3552	26	0
2	B	23	0	21	0	0
3	A	2	0	0	0	0
4	A	8	0	12	0	0
5	A	69	0	0	2	0
All	All	3703	0	3585	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:TYI:I2	1:A:174:ALA:HA	2.55	0.77
1:A:233:TRP:CD1	1:A:243:IYR:IE	3.26	0.58
1:A:73:GLN:HA	1:A:100:IYR:IE	2.73	0.58
1:A:243:IYR:IE	1:A:317:ARG:HD2	2.78	0.53
1:A:230:PRO:HG2	5:A:620:HOH:O	2.09	0.52
1:A:230:PRO:HA	1:A:233:TRP:CE2	2.45	0.52
1:A:241:VAL:O	1:A:243:IYR:HH	2.11	0.51
1:A:233:TRP:CG	1:A:243:IYR:IE	3.34	0.50
1:A:387:GLY:HA2	1:A:416:ARG:HD3	1.94	0.49
1:A:233:TRP:HB2	1:A:299:PRO:HG2	1.96	0.48
1:A:63:TRP:CD1	1:A:67:ARG:HG3	2.50	0.47
1:A:2:LYS:HA	1:A:2:LYS:HD2	1.73	0.47
1:A:44:LEU:HD13	1:A:61:ILE:HD11	1.96	0.46
1:A:406:GLU:HB3	1:A:426:MET:HE1	1.97	0.45
1:A:260:VAL:HB	1:A:330:ILE:HD13	1.97	0.45
1:A:131:GLU:O	1:A:134:PRO:HD2	2.18	0.43
1:A:308:TYI:I1	1:A:312:LEU:HD11	2.88	0.43
1:A:326:GLN:NE2	5:A:636:HOH:O	2.50	0.43
1:A:367:THR:OG1	1:A:454:ARG:NH1	2.51	0.43
1:A:69:GLY:HA3	1:A:333:ASN:O	2.18	0.42
1:A:383:GLN:HA	1:A:420:ASN:HB3	2.01	0.42
1:A:131:GLU:N	1:A:131:GLU:OE1	2.43	0.42
1:A:397:GLN:OE1	1:A:407:LYS:HD3	2.20	0.42
1:A:370:VAL:HG12	1:A:373:ALA:HB3	2.01	0.42
1:A:63:TRP:HB3	1:A:68:PHE:HE1	1.85	0.41
1:A:243:IYR:IE	1:A:245:VAL:CG2	3.39	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	453/469 (97%)	438 (97%)	15 (3%)	0	100	100



There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	362/368 (98%)	361 (100%)	1 (0%)	92 95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	259	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	IYR	A	452	1	12,13,14	0.87	0	14,17,19	0.70	0
1	TYI	A	172	1	13,14,15	0.90	0	16,19,21	0.67	0
1	TYI	A	448	1	13,14,15	0.85	0	16,19,21	0.68	0
1	IYR	A	100	1	12,13,14	0.79	0	14,17,19	1.06	2 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	IYR	A	243	1	12,13,14	0.89	0	14,17,19	0.97	1 (7%)
1	TYI	A	342	1	13,14,15	0.95	0	16,19,21	0.96	1 (6%)
1	TYI	A	308	1	13,14,15	0.80	0	16,19,21	0.84	0
1	TYI	A	211	1	13,14,15	0.85	0	16,19,21	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	IYR	A	452	1	-	0/5/6/8	0/1/1/1
1	TYI	A	172	1	-	0/5/6/8	0/1/1/1
1	TYI	A	448	1	-	0/5/6/8	0/1/1/1
1	IYR	A	100	1	-	0/5/6/8	0/1/1/1
1	IYR	A	243	1	-	0/5/6/8	0/1/1/1
1	TYI	A	342	1	-	0/5/6/8	0/1/1/1
1	TYI	A	308	1	-	1/5/6/8	0/1/1/1
1	TYI	A	211	1	-	2/5/6/8	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	342	TYI	CG-CB-CA	-3.12	107.79	114.10
1	A	100	IYR	CF-CE-IE	-2.35	117.48	119.81
1	A	243	IYR	CF-CE-IE	-2.19	117.64	119.81
1	A	100	IYR	CD-CE-IE	2.00	122.31	118.61

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	308	TYI	O-C-CA-CB
1	A	211	TYI	C-CA-CB-CG
1	A	211	TYI	N-CA-CB-CG

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	172	TYI	1	0
1	A	100	IYR	1	0
1	A	243	IYR	5	0
1	A	308	TYI	1	0

## 5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GLC	B	1	2	12,12,12	2.59	6 (50%)	17,17,17	1.31	2 (11%)
2	GLC	B	2	2	11,11,12	1.68	3 (27%)	15,15,17	0.93	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	2/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	GLC	C3-C2	-4.68	1.40	1.52
2	B	1	GLC	C6-C5	-4.62	1.36	1.51
2	B	1	GLC	O5-C5	3.31	1.52	1.44
2	B	2	GLC	O5-C1	3.03	1.48	1.43
2	B	2	GLC	C2-C3	-2.93	1.48	1.52
2	B	1	GLC	O2-C2	2.73	1.49	1.43
2	B	1	GLC	C1-C2	-2.57	1.46	1.52
2	B	1	GLC	O3-C3	2.40	1.48	1.43
2	B	2	GLC	O2-C2	-2.31	1.38	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	C1-C2-C3	2.56	115.62	110.31
2	B	1	GLC	O5-C1-C2	2.46	114.67	110.28
2	B	2	GLC	O5-C5-C6	2.23	110.71	107.20

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	GLC	C4-C5-C6-O6
2	B	1	GLC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	A	504	-	3,3,3	0.47	0	2,2,2	0.32	0
4	EDO	A	505	-	3,3,3	0.45	0	2,2,2	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	504	-	-	0/1/1/1	-
4	EDO	A	505	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	455/469 (97%)	-0.07	10 (2%) 62 58	42, 74, 120, 178	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	142	ALA	4.6
1	A	122	LEU	3.5
1	A	241	VAL	3.2
1	A	123	LEU	3.1
1	A	141	LYS	3.0
1	A	462	LEU	2.9
1	A	236	ILE	2.8
1	A	380	ARG	2.4
1	A	463	GLU	2.2
1	A	146	SER	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	IYR	A	243	13/14	0.88	0.21	81,94,103,108	1
1	TYI	A	211	14/15	0.90	0.14	62,111,136,179	0
1	IYR	A	100	13/14	0.94	0.14	51,62,74,190	1
1	IYR	A	452	13/14	0.96	0.14	87,93,116,133	5
1	TYI	A	308	14/15	0.97	0.12	70,82,91,118	0
1	TYI	A	448	14/15	0.98	0.11	77,93,103,106	0
1	TYI	A	342	14/15	0.98	0.11	53,69,98,122	0

*Continued on next page...*

*Continued from previous page...*


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	TYI	A	172	14/15	0.99	0.12	72,85,93,122	5

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GLC	B	1	12/12	0.96	0.17	45,54,63,71	0
2	GLC	B	2	11/12	0.98	0.15	41,50,54,55	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	IOD	A	503	1/1	0.89	0.08	108,108,108,108	1
4	EDO	A	504	4/4	0.92	0.15	77,82,82,83	0
3	IOD	A	502	1/1	0.93	0.10	117,117,117,117	1
4	EDO	A	505	4/4	0.94	0.13	72,74,76,77	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.