



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 09:11 pm BST

PDB ID : 5INW
Title : Structure of reaction loop cleaved lamprey angiotensinogen
Authors : Wei, H.; Zhou, A.
Deposited on : 2016-03-08
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	<div> <div></div> <div>73% 6% 20%</div> </div>
1	B	421	<div> <div>4%</div> <div>72% 6% 21%</div> </div>
2	C	35	<div> <div>74% 14% 11%</div> </div>
2	D	35	<div> <div>83% 9% 9%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative angiotensinogen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	0	0
			2616	1673	447	484	12			
1	B	331	Total	C	N	O	S	0	0	0
			2578	1646	440	480	12			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP C0MNC6
A	416A	ALA	-	insertion	UNP C0MNC6
A	416B	ALA	-	insertion	UNP C0MNC6
A	418	ARG	ILE	engineered mutation	UNP C0MNC6
B	0	GLY	-	expression tag	UNP C0MNC6
B	416A	ALA	-	insertion	UNP C0MNC6
B	416B	ALA	-	insertion	UNP C0MNC6
B	418	ARG	ILE	engineered mutation	UNP C0MNC6

- Molecule 2 is a protein called C-terminal peptide of Putative angiotensinogen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	31	Total	C	N	O	S	0	0	0
			248	162	42	43	1			
2	D	32	Total	C	N	O	S	0	0	0
			255	167	43	44	1			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	O	0	0
			2	2		



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	42.40Å 119.36Å 141.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	91.26 – 2.70 91.26 – 2.70	Depositor EDS
% Data completeness (in resolution range)	90.6 (91.26-2.70) 90.7 (91.26-2.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.204 , 0.277 0.206 , 0.273	Depositor DCC
R_{free} test set	930 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	62.3	Xtriage
Anisotropy	0.399	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5704	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	1/2668 (0.0%)	0.87	2/3596 (0.1%)
1	B	0.69	0/2629	0.88	5/3547 (0.1%)
2	C	0.73	0/253	0.96	0/343
2	D	0.72	0/261	0.89	0/355
All	All	0.73	1/5811 (0.0%)	0.88	7/7841 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	182	TRP	CB-CG	5.13	1.59	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	79	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	B	128	ARG	NE-CZ-NH1	6.11	123.36	120.30
1	B	271	TYR	CA-CB-CG	5.93	124.67	113.40
1	B	90	ARG	NE-CZ-NH2	-5.65	117.47	120.30
1	A	79	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	B	380	ARG	NE-CZ-NH1	5.36	122.98	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	380	ARG	NE-CZ-NH1	5.12	122.86	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	267	VAL	Peptide
1	A	341	ASN	Peptide
1	B	267	VAL	Peptide
1	B	341	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2616	0	2658	14	0
1	B	2578	0	2610	14	0
2	C	248	0	261	6	0
2	D	255	0	268	5	0
3	B	5	0	0	0	0
4	A	2	0	0	0	0
All	All	5704	0	5797	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:315:THR:HG21	2:C:429:PRO:HD2	1.87	0.57
1:B:172:LYS:HE3	1:B:221:VAL:HG11	1.88	0.56
1:A:172:LYS:HE3	1:A:221:VAL:HG11	1.87	0.56
1:A:300:PRO:HA	1:A:306:SER:HB3	1.88	0.56
1:B:266:TRP:CZ3	1:B:272:SER:HB3	2.42	0.56
1:A:266:TRP:CZ3	1:A:272:SER:HB3	2.43	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:VAL:O	1:A:221:VAL:HG12	2.06	0.53
1:B:300:PRO:HA	1:B:306:SER:HB3	1.90	0.53
1:B:221:VAL:HG12	1:B:221:VAL:O	2.09	0.52
1:A:309:VAL:HG13	2:C:430:PHE:CD1	2.46	0.51
1:B:253:ASN:HB2	1:B:280:GLN:HE22	1.77	0.50
1:A:300:PRO:HA	1:A:306:SER:CB	2.44	0.47
1:B:315:THR:HG21	2:D:429:PRO:HD2	1.96	0.47
1:A:219:GLU:HG3	1:A:224:LYS:HA	1.96	0.47
1:B:92:TYR:HB2	2:D:445:MET:HE1	1.97	0.45
1:B:219:GLU:HG3	1:B:224:LYS:HA	1.98	0.45
1:B:92:TYR:HB2	2:D:445:MET:CE	2.47	0.45
1:B:300:PRO:HA	1:B:306:SER:CB	2.47	0.44
1:A:288:GLN:OE1	1:A:313:HIS:NE2	2.50	0.43
1:A:105:ILE:O	2:C:446:GLY:HA3	2.19	0.43
1:A:92:TYR:HB2	2:C:445:MET:HE1	2.00	0.43
1:A:107:PHE:CZ	2:C:445:MET:HE2	2.54	0.42
1:B:107:PHE:CZ	2:D:445:MET:HE2	2.55	0.42
1:B:309:VAL:HG13	2:D:430:PHE:CD1	2.56	0.41
1:A:92:TYR:HB2	2:C:445:MET:CE	2.50	0.41
1:B:253:ASN:HB2	1:B:280:GLN:NE2	2.34	0.41
1:A:245:ILE:CG2	1:A:409:ALA:HB3	2.51	0.40
1:B:288:GLN:OE1	1:B:313:HIS:NE2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/421 (79%)	320 (97%)	9 (3%)	2 (1%)	25	50
1	B	327/421 (78%)	317 (97%)	9 (3%)	1 (0%)	41	66
2	C	29/35 (83%)	27 (93%)	2 (7%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	30/35 (86%)	29 (97%)	1 (3%)	0	100	100
All	All	717/912 (79%)	693 (97%)	21 (3%)	3 (0%)	34	60

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	74	LYS
1	A	223	SER
1	B	223	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/357 (80%)	285 (99%)	2 (1%)	84	94
1	B	283/357 (79%)	280 (99%)	3 (1%)	73	90
2	C	28/32 (88%)	27 (96%)	1 (4%)	35	64
2	D	29/32 (91%)	29 (100%)	0	100	100
All	All	627/778 (81%)	621 (99%)	6 (1%)	76	91

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	287	THR
1	A	347	SER
1	B	170	MET
1	B	287	THR
1	B	347	SER
2	C	428	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	GLN
1	B	280	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	501	-	4,4,4	0.26	0	6,6,6	0.70	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	335/421 (79%)	-0.18	4 (1%) 79 80	35, 62, 104, 139	0
1	B	331/421 (78%)	0.22	15 (4%) 33 31	45, 81, 119, 180	0
2	C	31/35 (88%)	-0.40	0 100 100	42, 60, 86, 87	0
2	D	32/35 (91%)	-0.14	0 100 100	57, 76, 106, 120	0
All	All	729/912 (79%)	-0.01	19 (2%) 56 57	35, 70, 115, 180	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	340	SER	11.1
1	B	163	ARG	5.4
1	B	338	GLY	4.9
1	B	164	SER	4.3
1	B	152	TYR	4.3
1	B	165	ALA	3.4
1	B	270	SER	3.2
1	B	286	TYR	2.8
1	B	339	PRO	2.6
1	B	167	PHE	2.6
1	A	281	ARG	2.5
1	A	374	TYR	2.3
1	B	264	ARG	2.3
1	A	271	TYR	2.2
1	B	268	ASN	2.2
1	B	266	TRP	2.1
1	B	166	GLY	2.1
1	A	152	TYR	2.1
1	B	271	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	B	501	5/5	0.99	0.11	53,54,57,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.