



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 08:17 am BST

PDB ID : 1IRJ  
Title : Crystal Structure of the MRP14 complexed with CHAPS  
Authors : Itou, H.; Yao, M.; Watanabe, N.; Nishihira, J.; Tanaka, I.  
Deposited on : 2001-10-09  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	113	58% 17% • 25%
1	B	113	58% 13% • 26%
1	C	113	46% 25% •• 27%
1	D	113	50% 22% • 27%
1	E	113	48% 25% • 27%
1	F	113	34% 36% • 26%
1	G	113	61% 12% • 26%
1	H	113	60% 14% • 25%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5871 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Migration Inhibitory Factor-Related Protein 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	85	Total	C	N	O	S	0	0	0
			707	445	121	136	5			
1	B	84	Total	C	N	O	S	0	0	0
			700	441	120	134	5			
1	C	83	Total	C	N	O	S	0	0	0
			694	438	119	133	4			
1	D	82	Total	C	N	O	S	0	0	0
			685	432	117	132	4			
1	E	83	Total	C	N	O	S	0	0	0
			694	438	119	133	4			
1	F	84	Total	C	N	O	S	0	0	0
			700	441	120	134	5			
1	G	84	Total	C	N	O	S	0	0	0
			700	441	120	134	5			
1	H	85	Total	C	N	O	S	0	0	0
			707	445	121	136	5			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	2	Total	Ca	0	0
			2	2		
2	D	2	Total	Ca	0	0
			2	2		
2	E	2	Total	Ca	0	0
			2	2		
2	H	2	Total	Ca	0	0
			2	2		
2	B	2	Total	Ca	0	0
			2	2		
2	C	2	Total	Ca	0	0
			2	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		
2	F	2	Total	Ca	0	0
			2	2		

- Molecule 3 is 3-[(3-CHOLAMIDOPROPYL)DIMETHYLAMMONIO]-1-PROPANESULFO NATE (three-letter code: CPS) (formula: C<sub>32</sub>H<sub>58</sub>N<sub>2</sub>O<sub>7</sub>S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			32	27	1	4		
3	D	1	Total	C	N	O	0	0
			32	27	1	4		
3	D	1	Total	C	N	O	0	0
			32	27	1	4		
3	G	1	Total	C	N	O	0	0
			32	27	1	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	19	Total	O	0	0
			19	19		
4	B	28	Total	O	0	0
			28	28		
4	C	6	Total	O	0	0
			6	6		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	20	Total 20	O 20	0	0
4	E	9	Total 9	O 9	0	0
4	F	2	Total 2	O 2	0	0
4	G	30	Total 30	O 30	0	0
4	H	26	Total 26	O 26	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: Migration Inhibitory Factor-Related Protein 14

Chain A: 




#### • Molecule 1: Migration Inhibitory Factor-Related Protein 14

Chain B: 




#### • Molecule 1: Migration Inhibitory Factor-Related Protein 14

Chain C: 




#### • Molecule 1: Migration Inhibitory Factor-Related Protein 14

Chain D: 



#### • Molecule 1: Migration Inhibitory Factor-Related Protein 14



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.59 Å 178.44 Å 61.23 Å 90.00° 113.17° 90.00°	Depositor
Resolution (Å)	10.00 – 2.10	Depositor
% Data completeness (in resolution range)	99.8 (10.00-2.10)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.245 , 0.274	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5871	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CPS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	0/716	0.58	0/957
1	B	0.43	0/709	0.64	1/947 (0.1%)
1	C	0.40	0/703	0.55	0/939
1	D	0.40	0/694	0.60	0/928
1	E	0.35	0/703	0.58	0/939
1	F	0.32	0/709	0.49	0/947
1	G	0.46	0/709	0.66	0/947
1	H	0.43	0/716	0.66	0/957
All	All	0.41	0/5659	0.60	1/7561 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	32	LEU	CA-CB-CG	5.36	127.64	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	707	0	709	18	0
1	B	700	0	702	20	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	694	0	697	33	0
1	D	685	0	684	21	0
1	E	694	0	697	51	0
1	F	700	0	702	77	0
1	G	700	0	702	15	0
1	H	707	0	709	12	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
2	G	2	0	0	0	0
2	H	2	0	0	0	0
3	B	32	0	42	5	0
3	D	64	0	84	12	0
3	G	32	0	42	6	0
4	A	19	0	0	1	0
4	B	28	0	0	0	0
4	C	6	0	0	0	0
4	D	20	0	0	0	0
4	E	9	0	0	0	0
4	F	2	0	0	0	0
4	G	30	0	0	2	0
4	H	26	0	0	0	0
All	All	5871	0	5770	241	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 241 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:189:CPS:C5	3:B:189:CPS:C10	1.76	1.63
3:D:289:CPS:C10	3:D:289:CPS:C5	1.75	1.62
3:G:489:CPS:C5	3:G:489:CPS:C10	1.75	1.62
3:D:389:CPS:C5	3:D:389:CPS:C10	1.76	1.61
1:E:5:MET:CE	1:F:47:ASN:H	1.76	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	83/113 (74%)	80 (96%)	3 (4%)	0	100	100
1	B	82/113 (73%)	81 (99%)	1 (1%)	0	100	100
1	C	81/113 (72%)	74 (91%)	6 (7%)	1 (1%)	13	8
1	D	80/113 (71%)	80 (100%)	0	0	100	100
1	E	81/113 (72%)	76 (94%)	5 (6%)	0	100	100
1	F	82/113 (73%)	66 (80%)	13 (16%)	3 (4%)	3	1
1	G	82/113 (73%)	81 (99%)	1 (1%)	0	100	100
1	H	83/113 (74%)	83 (100%)	0	0	100	100
All	All	654/904 (72%)	621 (95%)	29 (4%)	4 (1%)	25	21

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	43	LYS
1	F	45	LEU
1	C	51	LYS
1	F	52	GLU

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/102 (79%)	80 (99%)	1 (1%)	71	77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	80/102 (78%)	76 (95%)	4 (5%)	24	23
1	C	79/102 (78%)	76 (96%)	3 (4%)	33	34
1	D	78/102 (76%)	77 (99%)	1 (1%)	69	75
1	E	79/102 (78%)	78 (99%)	1 (1%)	69	75
1	F	80/102 (78%)	77 (96%)	3 (4%)	33	34
1	G	80/102 (78%)	78 (98%)	2 (2%)	47	52
1	H	81/102 (79%)	80 (99%)	1 (1%)	71	77
All	All	638/816 (78%)	622 (98%)	16 (2%)	47	52

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	57	LYS
1	D	32	LEU
1	F	55	ASN
1	C	53	ASN
1	G	4	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	7	GLN
1	E	73	GLN
1	H	61	HIS
1	E	28	HIS
1	F	17	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 16 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CPS	D	389	-	35,35,45	4.68	20 (57%)	54,54,70	3.92	29 (53%)
3	CPS	D	289	-	35,35,45	4.62	20 (57%)	54,54,70	3.75	29 (53%)
3	CPS	B	189	-	35,35,45	4.66	21 (60%)	54,54,70	3.67	26 (48%)
3	CPS	G	489	-	35,35,45	4.46	20 (57%)	54,54,70	3.73	28 (51%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CPS	D	389	-	-	4/13/78/90	0/4/4/4
3	CPS	D	289	-	-	0/13/78/90	0/4/4/4
3	CPS	B	189	-	-	2/13/78/90	0/4/4/4
3	CPS	G	489	-	-	0/13/78/90	0/4/4/4

The worst 5 of 81 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	389	CPS	C10-C5	13.21	1.76	1.54
3	B	189	CPS	C10-C5	13.18	1.76	1.54
3	D	389	CPS	O4-C4	-13.10	1.21	1.43
3	G	489	CPS	C10-C5	12.79	1.75	1.54
3	D	289	CPS	C10-C5	12.73	1.75	1.54

The worst 5 of 112 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	389	CPS	C6-C18-C17	12.57	128.47	111.81
3	D	289	CPS	C6-C18-C17	11.91	127.60	111.81
3	G	489	CPS	C6-C18-C17	11.39	126.91	111.81
3	B	189	CPS	C6-C18-C17	10.95	126.33	111.81
3	D	389	CPS	C19-C3-C4	9.49	126.84	114.30

There are no chirality outliers.

5 of 6 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
3	D	389	CPS	C23-C24-N1-C25
3	D	389	CPS	O1-C24-N1-C25
3	B	189	CPS	N1-C25-C26-C27
3	D	389	CPS	C22-C23-C24-O1
3	B	189	CPS	C23-C24-N1-C25


There are no ring outliers.

4 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	389	CPS	6	0
3	D	289	CPS	6	0
3	B	189	CPS	5	0
3	G	489	CPS	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.