



wwPDB X-ray Structure Validation Summary Report ⓘ

May 18, 2020 – 07:55 am BST

PDB ID : 6IS1
Title : Crystal Structure of Staphylococcus aureus response regulator ArlR receiver domain in complex with BeF3 and Mg
Authors : Wen, Y.; Ouyang, Z.
Deposited on : 2018-11-15
Resolution : 1.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	128	<div> <div>2%</div> <div>85% 9% 6%</div> </div>
1	B	128	<div> <div>81% 13% 5%</div> </div>
1	C	128	<div> <div>2%</div> <div>83% 10% 7%</div> </div>
1	D	128	<div> <div>2%</div> <div>78% 13% 7%</div> </div>
1	E	128	<div> <div>3%</div> <div>78% 14% 8%</div> </div>
1	F	128	<div> <div>84% 9% 5%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	128	 2% 84% 9% 6%
1	H	128	 86% 8% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	IMD	H	303	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 16897 atoms, of which 7828 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Response regulator ArlR.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	120	Total	C	H	N	O	S	0	3	0
			1973	629	982	164	195	3			
1	C	119	Total	C	H	N	O	S	0	1	0
			1931	615	962	160	192	2			
1	B	121	Total	C	H	N	O	S	0	4	0
			2017	642	1006	170	196	3			
1	D	119	Total	C	H	N	O	S	0	1	0
			1950	620	973	164	191	2			
1	E	118	Total	C	H	N	O	S	0	0	0
			1920	610	959	160	189	2			
1	F	121	Total	C	H	N	O	S	0	1	0
			1972	629	980	167	194	2			
1	G	120	Total	C	H	N	O	S	0	0	0
			1942	617	970	161	192	2			
1	H	120	Total	C	H	N	O	S	0	1	0
			1963	624	981	163	192	3			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	initiating methionine	UNP Q2YY03
A	-5	GLY	-	expression tag	UNP Q2YY03
A	-4	HIS	-	expression tag	UNP Q2YY03
A	-3	HIS	-	expression tag	UNP Q2YY03
A	-2	HIS	-	expression tag	UNP Q2YY03
A	-1	HIS	-	expression tag	UNP Q2YY03
A	0	HIS	-	expression tag	UNP Q2YY03
A	1	HIS	-	expression tag	UNP Q2YY03
C	-6	MET	-	initiating methionine	UNP Q2YY03
C	-5	GLY	-	expression tag	UNP Q2YY03
C	-4	HIS	-	expression tag	UNP Q2YY03
C	-3	HIS	-	expression tag	UNP Q2YY03
C	-2	HIS	-	expression tag	UNP Q2YY03

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	HIS	-	expression tag	UNP Q2YY03
C	0	HIS	-	expression tag	UNP Q2YY03
C	1	HIS	-	expression tag	UNP Q2YY03
B	-6	MET	-	initiating methionine	UNP Q2YY03
B	-5	GLY	-	expression tag	UNP Q2YY03
B	-4	HIS	-	expression tag	UNP Q2YY03
B	-3	HIS	-	expression tag	UNP Q2YY03
B	-2	HIS	-	expression tag	UNP Q2YY03
B	-1	HIS	-	expression tag	UNP Q2YY03
B	0	HIS	-	expression tag	UNP Q2YY03
B	1	HIS	-	expression tag	UNP Q2YY03
D	-6	MET	-	initiating methionine	UNP Q2YY03
D	-5	GLY	-	expression tag	UNP Q2YY03
D	-4	HIS	-	expression tag	UNP Q2YY03
D	-3	HIS	-	expression tag	UNP Q2YY03
D	-2	HIS	-	expression tag	UNP Q2YY03
D	-1	HIS	-	expression tag	UNP Q2YY03
D	0	HIS	-	expression tag	UNP Q2YY03
D	1	HIS	-	expression tag	UNP Q2YY03
E	-6	MET	-	initiating methionine	UNP Q2YY03
E	-5	GLY	-	expression tag	UNP Q2YY03
E	-4	HIS	-	expression tag	UNP Q2YY03
E	-3	HIS	-	expression tag	UNP Q2YY03
E	-2	HIS	-	expression tag	UNP Q2YY03
E	-1	HIS	-	expression tag	UNP Q2YY03
E	0	HIS	-	expression tag	UNP Q2YY03
E	1	HIS	-	expression tag	UNP Q2YY03
F	-6	MET	-	initiating methionine	UNP Q2YY03
F	-5	GLY	-	expression tag	UNP Q2YY03
F	-4	HIS	-	expression tag	UNP Q2YY03
F	-3	HIS	-	expression tag	UNP Q2YY03
F	-2	HIS	-	expression tag	UNP Q2YY03
F	-1	HIS	-	expression tag	UNP Q2YY03
F	0	HIS	-	expression tag	UNP Q2YY03
F	1	HIS	-	expression tag	UNP Q2YY03
G	-6	MET	-	initiating methionine	UNP Q2YY03
G	-5	GLY	-	expression tag	UNP Q2YY03
G	-4	HIS	-	expression tag	UNP Q2YY03
G	-3	HIS	-	expression tag	UNP Q2YY03
G	-2	HIS	-	expression tag	UNP Q2YY03
G	-1	HIS	-	expression tag	UNP Q2YY03
G	0	HIS	-	expression tag	UNP Q2YY03

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	1	HIS	-	expression tag	UNP Q2YY03
H	-6	MET	-	initiating methionine	UNP Q2YY03
H	-5	GLY	-	expression tag	UNP Q2YY03
H	-4	HIS	-	expression tag	UNP Q2YY03
H	-3	HIS	-	expression tag	UNP Q2YY03
H	-2	HIS	-	expression tag	UNP Q2YY03
H	-1	HIS	-	expression tag	UNP Q2YY03
H	0	HIS	-	expression tag	UNP Q2YY03
H	1	HIS	-	expression tag	UNP Q2YY03

- Molecule 2 is BERYLLIUM TRIFLUORIDE ION (three-letter code: BEF) (formula: BeF₃) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Be	F	0	0
			4	1	3		
2	C	1	Total	Be	F	0	0
			4	1	3		
2	B	1	Total	Be	F	0	0
			4	1	3		
2	D	1	Total	Be	F	0	0
			4	1	3		
2	E	1	Total	Be	F	0	0
			4	1	3		
2	F	1	Total	Be	F	0	0
			4	1	3		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	G	1	Total 4	Be 1	F 3	0	0
2	H	1	Total 4	Be 1	F 3	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total 1	Mg 1	0	0
3	D	1	Total 1	Mg 1	0	0
3	E	1	Total 1	Mg 1	0	0
3	H	1	Total 1	Mg 1	0	0
3	B	1	Total 1	Mg 1	0	0
3	C	1	Total 1	Mg 1	0	0
3	A	1	Total 1	Mg 1	0	0
3	F	1	Total 1	Mg 1	0	0

- Molecule 4 is IMIDAZOLE (three-letter code: IMD) (formula: C₃H₅N₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	N	0	0
			10	3	5	2		
4	F	1	Total	C	H	N	0	0
			10	3	5	2		
4	H	1	Total	C	H	N	0	0
			10	3	5	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	174	Total	O	0	0
			174	174		
5	C	83	Total	O	0	0
			83	83		
5	B	195	Total	O	0	0
			195	195		
5	D	146	Total	O	0	0
			146	146		
5	E	86	Total	O	0	0
			86	86		
5	F	184	Total	O	0	0
			184	184		
5	G	109	Total	O	0	0
			109	109		
5	H	182	Total	O	0	0
			182	182		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Response regulator ArlR




- Molecule 1: Response regulator ArlR




- Molecule 1: Response regulator ArlR




- Molecule 1: Response regulator ArlR



- Molecule 1: Response regulator ArlR




- Molecule 1: Response regulator ArlR

Chain F:  84% 9% • 5%




- Molecule 1: Response regulator ArlR

Chain G:  84% 9% • 6%



- Molecule 1: Response regulator ArlR

Chain H:  86% 8% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	61.11Å 61.74Å 66.06Å 90.36° 105.45° 92.01°	Depositor
Resolution (Å)	31.77 – 1.59 43.95 – 1.59	Depositor EDS
% Data completeness (in resolution range)	93.7 (31.77-1.59) 93.7 (43.95-1.59)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.59Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.159 , 0.189 0.159 , 0.189	Depositor DCC
R_{free} test set	2020 reflections (1.72%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtriage
Anisotropy	0.098	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for -h,k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	16897	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 70.33 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.1805e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BEF, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/1014	0.73	2/1373 (0.1%)
1	B	0.52	0/1039	0.69	1/1407 (0.1%)
1	C	0.41	0/985	0.67	0/1335
1	D	0.47	0/995	0.69	2/1348 (0.1%)
1	E	0.43	0/975	0.63	0/1321
1	F	0.53	0/1011	0.73	1/1371 (0.1%)
1	G	0.47	0/986	0.70	1/1337 (0.1%)
1	H	0.51	0/1000	0.76	1/1355 (0.1%)
All	All	0.48	0/8005	0.70	8/10847 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	52	ASP	CB-CG-OD2	-6.71	112.26	118.30
1	A	9	ASP	CB-CG-OD1	6.65	124.28	118.30
1	H	9	ASP	CB-CG-OD1	5.67	123.40	118.30
1	G	52	ASP	CB-CG-OD2	-5.18	113.64	118.30
1	B	52	ASP	CB-CG-OD2	-5.18	113.64	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	1	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	991	982	993	10	0
1	B	1011	1006	1018	14	0
1	C	969	962	966	6	0
1	D	977	973	977	16	0
1	E	961	959	962	10	1
1	F	992	980	990	9	0
1	G	972	970	972	7	0
1	H	982	981	986	7	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
2	D	4	0	0	0	0
2	E	4	0	0	0	0
2	F	4	0	0	0	0
2	G	4	0	0	0	0
2	H	4	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
4	B	5	5	5	0	0
4	F	5	5	5	3	0
4	H	5	5	5	4	0
5	A	174	0	0	8	2
5	B	195	0	0	8	4
5	C	83	0	0	2	2
5	D	146	0	0	9	3
5	E	86	0	0	2	3
5	F	184	0	0	4	3

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	G	109	0	0	3	2
5	H	182	0	0	4	2
All	All	9069	7828	7879	81	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 81 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:18:GLU:OE1	5:H:401:HOH:O	1.84	0.94
1:G:87:LYS:NZ	5:G:401:HOH:O	2.00	0.94
1:F:54:MET:SD	5:F:569:HOH:O	2.25	0.94
1:A:106:GLU:OE2	5:A:401:HOH:O	1.98	0.81
1:B:119:GLN:NE2	5:B:401:HOH:O	2.16	0.77

The worst 5 of 11 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:565:HOH:O	5:F:548:HOH:O[1_556]	1.64	0.56
5:C:483:HOH:O	5:E:468:HOH:O[1_655]	1.89	0.31
5:D:524:HOH:O	5:E:454:HOH:O[1_656]	1.98	0.22
5:B:547:HOH:O	5:F:474:HOH:O[1_556]	2.04	0.16
5:G:467:HOH:O	5:H:568:HOH:O[1_455]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	121/128 (94%)	120 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	123/128 (96%)	120 (98%)	3 (2%)	0	100	100
1	C	118/128 (92%)	116 (98%)	2 (2%)	0	100	100
1	D	118/128 (92%)	116 (98%)	2 (2%)	0	100	100
1	E	116/128 (91%)	114 (98%)	2 (2%)	0	100	100
1	F	120/128 (94%)	118 (98%)	2 (2%)	0	100	100
1	G	118/128 (92%)	116 (98%)	2 (2%)	0	100	100
1	H	119/128 (93%)	118 (99%)	1 (1%)	0	100	100
All	All	953/1024 (93%)	938 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/115 (96%)	110 (100%)	0	100	100
1	B	113/115 (98%)	113 (100%)	0	100	100
1	C	106/115 (92%)	104 (98%)	2 (2%)	57	34
1	D	108/115 (94%)	105 (97%)	3 (3%)	43	18
1	E	106/115 (92%)	102 (96%)	4 (4%)	33	10
1	F	110/115 (96%)	109 (99%)	1 (1%)	78	65
1	G	107/115 (93%)	105 (98%)	2 (2%)	57	34
1	H	109/115 (95%)	109 (100%)	0	100	100
All	All	869/920 (94%)	857 (99%)	12 (1%)	67	47

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	1	HIS
1	E	10	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	93	TYR
1	D	12	ASN
1	E	44	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	119	GLN
1	B	70	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 8 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BEF	H	301	1	0,3,3	0.00	-	-		
4	IMD	B	303	-	3,5,5	0.42	0	4,5,5	0.64	0
4	IMD	F	303	-	3,5,5	0.44	0	4,5,5	0.61	0
2	BEF	A	301	1	0,3,3	0.00	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BEF	E	301	1	0,3,3	0.00	-	-		
2	BEF	B	301	1	0,3,3	0.00	-	-		
2	BEF	C	301	1	0,3,3	0.00	-	-		
2	BEF	F	301	1	0,3,3	0.00	-	-		
2	BEF	G	301	1	0,3,3	0.00	-	-		
2	BEF	D	301	1	0,3,3	0.00	-	-		
4	IMD	H	303	-	3,5,5	0.46	0	4,5,5	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IMD	B	303	-	-	-	0/1/1/1
4	IMD	F	303	-	-	-	0/1/1/1
4	IMD	H	303	-	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


2 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	303	IMD	3	0
4	H	303	IMD	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	120/128 (93%)	-0.30	1 (0%) 86 86	12, 18, 38, 57	0
1	B	121/128 (94%)	-0.30	0 100 100	11, 17, 34, 71	0
1	C	119/128 (92%)	0.13	3 (2%) 57 55	16, 31, 51, 71	0
1	D	119/128 (92%)	-0.29	2 (1%) 70 69	13, 22, 43, 63	0
1	E	118/128 (92%)	0.10	4 (3%) 45 42	15, 32, 54, 72	0
1	F	121/128 (94%)	-0.30	0 100 100	11, 17, 34, 54	0
1	G	120/128 (93%)	-0.07	3 (2%) 57 55	14, 26, 49, 65	0
1	H	120/128 (93%)	-0.30	0 100 100	12, 18, 36, 67	0
All	All	958/1024 (93%)	-0.17	13 (1%) 75 75	11, 21, 48, 72	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	85	TYR	3.9
1	E	85	TYR	3.7
1	G	120	PRO	3.3
1	A	45	TYR	3.0
1	G	85	TYR	3.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.




6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	IMD	F	303	5/5	0.26	0.28	30,45,59,60	0
4	IMD	H	303	5/5	0.46	0.32	35,44,69,77	0
2	BEF	E	301	4/4	0.85	0.10	29,30,32,37	0
3	MG	G	302	1/1	0.85	0.08	22,22,22,22	0
2	BEF	C	301	4/4	0.88	0.08	30,31,35,36	0
4	IMD	B	303	5/5	0.88	0.15	30,43,53,55	0
3	MG	C	302	1/1	0.90	0.06	33,33,33,33	0
2	BEF	G	301	4/4	0.93	0.10	19,21,21,24	0
3	MG	E	302	1/1	0.93	0.07	40,40,40,40	0
2	BEF	D	301	4/4	0.94	0.09	17,20,21,22	0
3	MG	D	302	1/1	0.95	0.06	20,20,20,20	0
2	BEF	A	301	4/4	0.96	0.12	14,15,15,17	0
2	BEF	B	301	4/4	0.97	0.10	13,14,16,16	0
2	BEF	F	301	4/4	0.98	0.10	12,12,13,13	0
3	MG	A	302	1/1	0.98	0.07	15,15,15,15	0
2	BEF	H	301	4/4	0.98	0.10	15,15,16,16	0
3	MG	B	302	1/1	0.98	0.07	13,13,13,13	0
3	MG	F	302	1/1	0.99	0.06	12,12,12,12	0
3	MG	H	302	1/1	1.00	0.04	13,13,13,13	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG G 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BEF C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG C 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BEF G 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG E 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BEF D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG D 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BEF A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BEF B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BEF F 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BEF H 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG F 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG H 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.